

## Effects of Dietary Energy and Protein Levels on Growth Curve Parameters of Khazak Native Chickens

### Research Article

H. Faraji-Arough<sup>1\*</sup>, M. Ghazaghi<sup>2</sup>, F. Bagherzadeh Kasmani<sup>2</sup> and M. Rokouei<sup>2</sup>

<sup>1</sup> Department of Ostrich, Special Domestic Animals Institute, Research Institute of Zabol, Zabol, Iran

<sup>2</sup> Department of Animal Science, Faculty of Agriculture, University of Zabol, Zabol, Iran

Received on: 26 Oct 2022

Revised on: 6 Jan 2023

Accepted on: 25 Feb 2023

Online Published on: Sep 2023

\*Correspondence E-mail: [hadifaraji@uoz.ac.ir](mailto:hadifaraji@uoz.ac.ir)

© 2010 Copyright by Islamic Azad University, Rasht Branch, Rasht, Iran

Online version is available on: [www.ijas.ir](http://www.ijas.ir)

### ABSTRACT

This study was conducted to evaluate the effect of dietary energy (ME) and protein (CP) on growth curve parameters and absolute growth rates in the different ages of the Khazak chicks. A total of 360 one-day-old Khazak chicks were obtained from a local hatchery and in a  $3 \times 3$  factorial experiment with completely randomized design, chicks were randomly allocated to experimental diets including 2600, 2800, and 3000 kcal of ME/kg, and each containing 17, 19, and 21% CP from 7 to 98 days of age. Four growth model (Gompertz, Logistic, Lopez, and Richards) were fitted on weekly body weight data and the best model were selected by the goodness of fit criteria. Growth curve parameters were predicted for all chicks using the best model and other parameters including age ( $T_i$ ) and weight ( $W_i$ ) at the inflection point and absolute growth rate (AGR) in different ages were calculated from growth curve parameters. Based on goodness of fit criteria, the Richards model had the lowest Akaike's Information Criteria (AIC), root mean square error (RMSE), and highest adjusted determination coefficient ( $R^2_{Adj}$ ) than other models and was selected as the best model. The effect of ME was significant on the mature index ( $k$ ),  $W_i$ ,  $T_i$ , and all AGR parameters ( $P < 0.05$ ) while CP levels were significant on final weight ( $W_f$ ),  $W_i$ , and AGR parameters ( $P < 0.05$ ). The chicks fed with a diet containing 2600 kcal of ME/kg and 17% CP had the higher  $k$  parameter, and lower  $W_i$ ,  $T_i$ , and AGR than those fed with other diets ( $P < 0.05$ ). Considering that the level of 2800 kcal of ME/kg and 19% CP had no significant difference with the level of 3000 kcal of ME/kg and 21% CP, therefore diet with 2800 kcal of ME/kg and 19% CP was suggested as optimum levels for change the growth curve parameter and having best performance for Khazak chickens during 7 to 98 days of old.

**KEY WORDS** inflection point, growth, mathematical model, native chicken.

### INTRODUCTION

In tropical rural areas, rearing native poultry plays a pivotal role in the production of high-quality animal protein resources with organic properties and income generation (Norris *et al.* 2007; Padhi, 2016). Recently, public concern was increased about the use of modern broiler genotypes for the production of chicken meat, and the tendency is increasing toward the consumption of meat that produced from slow-growing broilers instead of fast-growing

(Dyubele *et al.* 2010). Furthermore, meat produced by modern broiler genotypes is less palatable than native breeds (Wattanachant, 2008). Rural poultry has significant prospects for future development due to the easy and abundant availability of all necessities input including land, labor, and feed resources in rural areas. This section can help in increasing household income and improving family health through best nutrition (Shehbaz and Khan, 2008). Knowledge of bird requirements to access their maximum production capacity is one of the solutions to increasing

poultry production in the countries. It is very important to consider the main components of feed due to the high cost of feed in poultry production farms (Wijten *et al.* 2004). Energy and protein are the main nutrients in most livestock diets that allotted about 90% of the total feed cost. Therefore, these nutrients must be used most efficiently for the formulation of poultry rations and the profitability of production (Durunna *et al.* 2005). There is little information about nutritional improvement in native breed than commercial broiler chickens. In this regard, determining the optimal energy and protein levels of the diet is very important to maximize the production parameters and carcass characteristics of native chickens (Miah *et al.* 2014). In some studies, the effect of different levels of energy and protein on growth performance in Korat (Maliwan *et al.* 2019), Venaraja (Perween *et al.* 2016), Desi (Miah *et al.* 2014), Arabi (Al-Khalifa and Al-Nasser, 2012), Assel (Haunshi *et al.* 2012) Betong (Nguyen *et al.* 2010), chickens were reported. However, in these studies, the effect of energy and protein was investigated on body weight gain, feed intake, and feed conversion ratio.

The growth changes of an animal can be measured as body weight in regular intervals and summarized by mathematical models fitted to growth curves (Aggrey, 2002). The growth curve can be useful in describing the production of animals, especially when they can be estimated using the number of daily feed requirements (Abbas *et al.* 2014). It has been shown that the shape of the growth curve was affected by the composition of the diet (Mohammed, 2015). Prediction of production, as well as the nutritional requirements of birds of different ages, can result in a restriction on the level of *ad libitum* access to feed (Lopez *et al.* 2000).

Studies on long-term growth curves of animals can be helpful in the dynamically understanding of growth patterns and responses to dietary nutrient density as well (Russo *et al.* 2009; Yun *et al.* 2015). In a study by Nahashon *et al.* (2010) reported that dietary protein and energy affect the growth parameters of the French guinea fowl by using of Gompertz-Laired model. Growth curve parameters of commercial broiler and native chickens that fed by different energy levels by four growth models were studied and a significant effect of energy level was reported on some growth parameters (Moharrery and Mirzaei, 2014).

The Khazak breed is one of the small native breeds in the Sistan (Sistan region, Iran), with short legs relatively high potential for egg production. A tendency toward the native chicken consumption in this region was more than industrial chickens, so an improvement of the performance for this breed in terms of growth by demining appropriate

levels of energy and protein could be useful (Faraji-Arough *et al.* 2019). Therefore, this study aimed to evaluate the growth curve parameters of Khazak chickens fed diets containing different levels of energy and protein, and to select the appropriate levels of energy and protein to improve the growth curve parameters.

## MATERIALS AND METHODS

### Experiment design and bird management

The present study was performed in an experiment poultry farm of the Research Center of Domestic Animals Institute, Research Institute of Zabol, Zabol, Iran. Animal handling and experimental procedures of the study were conducted following approved guidelines of the Research Animal Committee of the Research Institute and Iranian Council of Animal Care (1995). A total of 360 one-day-old Khazak chicks were wing-banded and weighed individually. Chicks were raised together until 7 days of age in floor pens containing litter composed of wood shaving. At one week of age, chicks were weighed and randomly distributed into nine groups. Each group had 40 chicks that were allocated into 4 replicates with 10 birds in each. The chicks were fed with a maize-soybean meal-based diet supplying three levels of metabolizable energy (2600, 2800, and 3000 kcal/kg) and three levels of crude protein (17%, 19%, and 21%) in a 3 × 3 factorial experiment with a completely randomized design (Table 1). Feed and water were provided *ad libitum* during the experiment (7 to 98 days of age). A brooding temperature was maintained at 35 °C from day 1 to 7 and then gradually reduced 2 °C per week until 21 °C. The chicks were individually weighed weekly until the 14<sup>th</sup> week of age.

### Mathematical models

Four non-linear mathematical models including Gompertz, Logistic, Lopez, and Richards were fitted to the body weight data to recognize the best model for predicting growth curve parameters. The age and weight at the inflection point and Absolute growth weight in different ages for each model were calculated based on the model parameters. The equations of fitted models and biological parameters are shown in Table 2. In all models,  $W$  is the body weight of a bird at age  $t$ ,  $W_0$ ,  $W_f$ , and  $k$  are initial and final weights, and coefficient of relative growth or maturing index, respectively. The parameter  $b$  indicates the age at approximately half the maximum body weight, and  $m$  represents the shape parameter. The fitting of models on body weight data was performed by nlme package of R software (Pinheiro *et al.* 2014).

**Table 1** Ingredients and nutrient composition of the experimental diets

Item	Diets								
	1	2	3	4	5	6	7	8	9
<b>Ingredient</b>									
Corn grain	57.92	55.36	52.81	66.87	64.31	61.82	73.79	67.66	61.43
Soybean meal (44% CP)	18.07	23.45	28.87	19.76	25.15	30.64	21.29	26.54	31.80
Wheat bran	19.46	16.94	14.29	8.87	6.35	3.43	-	-	-
Dicalcium phosphate	1.64	1.62	1.60	1.75	1.73	1.72	1.84	1.80	1.76
Calcium CO <sub>3</sub>	1.47	1.45	1.43	1.43	1.41	1.39	1.40	1.38	1.37
Sodium bicarbonate	0.54	0.32	0.17	0.41	0.19	0.17	0.30	0.18	0.17
NaCl	0.29	0.29	0.29	0.30	0.30	0.30	0.21	0.30	0.30
Trace mineral premix <sup>1</sup>	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25
Vitamin premix <sup>2</sup>	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25
DL-methionine	0.11	0.06	0.05	0.10	0.06	0.03	0.10	0.05	0.03
Oil	-	-	-	-	-	-	0.57	1.59	2.63
<b>Calculated chemical composition</b>									
Metabolizable energy (kcal/kg)	2600	2600	2600	2800	2800	2800	3000	3000	3000
Crude protein (%)	17.00	19.00	21.00	17.00	19.00	21.00	17.00	19.00	21.00
Lysine (%)	0.78	0.91	1.05	0.79	0.92	1.06	0.79	0.93	1.07
Methionine (%)	0.33	0.35	0.36	0.33	0.35	0.35	0.33	0.35	0.35
Methionine + cysteine (%)	0.62	0.63	0.66	0.62	0.63	0.66	0.62	0.63	0.65
Calcium (%)	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Available phosphorus (%)	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50

<sup>1</sup> Vitamin premix provided per kilogram of diet: vitamin A (from vitamin A acetate): 11500 U; Cholecalciferol: 2100 U; vitamin E (from DL- $\alpha$ -tocopheryl acetate): 22 U; vitamin B<sub>12</sub>: 0.60 mg; Riboflavin: 4.4 mg; Nicotinamide: 40 mg; Calcium pantothenate: 35 mg; Menadione (from menadione dimethyl-pyrimidinol): 1.50 mg; Folic acid: 0.80 mg; Thiamine: 3 mg; Pyridoxine: 10 mg; Biotin: 1 mg; Choline chloride: 560 mg and Ethoxyquin: 125 mg.

<sup>2</sup> Mineral premix provided per kilogram of diet: Mn (from MnSO<sub>4</sub>·H<sub>2</sub>O): 65 mg; Zn (from ZnO): 55 mg; Fe (from FeSO<sub>4</sub>·7H<sub>2</sub>O): 50 mg; Cu (from CuSO<sub>4</sub>·5H<sub>2</sub>O): 8 mg; I (from Ca (IO<sub>3</sub>)<sub>2</sub>·H<sub>2</sub>O): 1.8 mg; Se: 0.30 mg; Co (from Co<sub>2</sub>O<sub>3</sub>): 0.20 mg and Mo: 0.16 mg.

<sup>3</sup> DEB: dietary electrolyte balance represents dietary Na + K - Cl in mEq/kg of diet.

**Table 2** Equation of mathematical models and biological parameters

Model <sup>1</sup>	Equation	T <sub>i</sub>	W <sub>i</sub>	Absolute growth rate
Gompertz	$W = W_0 \exp \left\{ \left[ 1 - \exp(-k \times t) \right] \ln \left( \frac{W_f}{W_0} \right) \right\}$	$\frac{1}{k} \left[ \ln \left( \ln \left( \frac{W_f}{W_0} \right) \right) \right]$	$\frac{W_f}{e}$	$kW \ln \left( \frac{W_f}{W} \right)$
Richards	$W = \frac{W_0 W_f}{[W_0^m + (W_f^m - W_0^m) e^{-kt}]^{1/m}}$	$\frac{1}{k} \times \ln \left( \frac{m}{(W_f^m - W_0^m)/W_0^m} \right)$	$\frac{W_f}{\sqrt[m]{m+1}}$	$kW \left( \frac{W_f^m - W^m}{mW_f^m} \right)$
Logistic	$W = \frac{W_0 W_f}{[W_0 + (W_f - W_0) \exp(-k \times t)]}$	$\frac{1}{k} \ln \left( \frac{W_f - W_0}{W_0} \right)$	$\frac{W_f}{2}$	$kW \left( 1 - \frac{W_f}{W} \right)$
Lopez	$W = \frac{(W_0 \times b^k) + (W_f \times t^k)}{(b^k + t^k)}$	$b \left( \frac{k-1}{k+1} \right)^{1/2}$	$\frac{\left[ \left( 1 + \frac{1}{k} \right) W_0 + \left( 1 - \frac{1}{k} \right) W_f \right]}{2}$	$k \left( \frac{t^{k-1}}{b^k + t^k} \right) (W_f - W)$

<sup>1</sup> W: body weight of bird at age t; W<sub>0</sub>: initial weight; W<sub>f</sub>: final weight; k: coefficient of relative growth or maturing index; b: age at approximately half maximum body weight and m: shape parameter.

After fitting the models, four goodness of fit criteria were used to compare the models and selection of the best model for studied populations (Teleken *et al.* 2017):

1) Adjusted determination coefficient ( $R^2_{Adj}$ ):

$$R^2_{Adj} = 1 - \left[ \left( \frac{n-1}{n-k} \right) * (1 - R^2_{model}) \right]$$

2) Akaike's Information Criteria (AIC) =  $n \ln(SSE/n) + 2k$

3) Bayesian Information Criterion (BIC) =  $n \ln(SSE/n) + k \ln(n)$

4) Root mean square error (RMSE):

$$RMSE = \sqrt{\frac{SSE}{n-k}}$$

Where:

n and k: number of observation and parameters, respectively.

$R^2_{model}$ : determination coefficient that is equal to  $1 - (SSE/SST)$ .

SSE and SST: sum of square errors and total sum of squares, respectively. Smaller value for AIC, BIC, RMSE and the highest value for  $R^2_{Adj}$  a model indicate the best model.

The growth curve parameters for each bird were obtained by the nlsList package of R software using the best model. Then age and weight at the inflection point and absolute growth rate in different ages for each bird have calculated as the model parameters.

### Statistical analysis

Data were analyzed using the general linear model procedure of SAS (SAS, 2008). A two-way ANOVA was performed to test the main effects of different levels of energy and protein and their interaction effects on model parameters, age ( $T_i$ ) and weight ( $W_i$ ) at the inflection point, and absolute growth rate at ages 14, 21, 28, 35, 42, 49, 56, 63, 70, 77, 84, 91 and 98 days of age. The differences among treatment means were compared by Tukey's test procedure and means considered significant when  $P < 0.05$ .

## RESULTS AND DISCUSSION

The mean and standard error of growth parameters of studied models for all populations is shown in Table 3. Logistic and Lopez models overestimated  $W_0$  and  $W_f$  parameters than other models, respectively. The highest and lowest value of the k parameter was estimated by Lopez and Gompertz models, respectively.

The values of age ( $T_i$ ) and weight ( $W_i$ ) at the inflection point in the Lopez model were lower than other models despite having higher  $W_f$  and k parameters. However, the highest value for mentioned parameter was estimated by the Logistic model (56.965 days and 447.600g, respectively). Regarding the goodness of fit criteria, four models were fitted on body weight data and are suitable to describe the growth curve of this population. However, the smallest AIC and RMSE were calculated for the Richards model among all models, and the value  $R^2_{Adj}$  for Richards was higher than other models. Therefore, the Richards model was selected as the best model to describe the growth curve in this study, and the effect of different levels of energy and protein on growth curve parameters were evaluated with this model.

The mean growth curve parameters and absolute growth rate in different ages for ME and CP levels are presented in Table 4. The initial body weight ( $W_0$ ) was high in birds fed a diet containing 3000 kcal of ME/kg than other levels of ME, but the difference between ME levels was not significant ( $P > 0.05$ ). A similar trend was observed for final body weight ( $W_f$ ) so that the final body weight was high in higher levels of ME ( $P > 0.05$ ). However, the predicted maturing index (k) was significantly lower ( $P < 0.05$ ) in birds fed with a diet containing 3000 kcal of ME/kg compared with those fed the 2600 and 2800 kcal of ME/kg diets.

The difference between shape parameter (m) among dietary ME levels was not significant ( $P > 0.05$ ), but the difference of age ( $T_i$ ) and weight ( $W_i$ ) at the inflection point among ME levels were significant ( $P < 0.05$ ) so that birds fed with a diet containing 3000 kcal of ME/kg arrived at age at the inflection point in higher ages than other ME levels. For as much as the birds fed with higher levels (3000 and 2800 kcal/kg) of ME had a higher  $T_i$  so these birds showed a higher weight at the inflection point ( $W_i$ ) than those fed diet containing 2600 kcal of ME/kg ( $P < 0.05$ ).

The effect of diet CP levels was significant on the  $W_f$  and  $W_i$  parameters were significant ( $P < 0.05$ , Table 4) so that the final weight and weight at the inflection point of birds fed with a diet containing 21 and 19% CP was higher than those fed with the 17% CP diet. Although, diets containing high CP had higher  $W_0$ , m, and  $T_i$  values and lower k values than other diets, the difference between CP levels was not significant ( $P > 0.05$ ).

The mean absolute growth rate in different ages for various levels of ME and CP are presented in Table 4. The effect of ME and CP levels was significant on absolute growth rate in different ages ( $P > 0.05$ ). The highest absolute growth rate was observed for birds fed a diet containing 3000 kcal of ME/kg and 21% CP, while the lowest absolute growth rate was related to the diet containing 2600 kcal of ME/kg and 17% CP.

**Table 3** Estimated parameters and goodness of fit criteria for each model

Parameter <sup>1</sup>	Gompertz	Logistic	Lopez	Richards
W <sub>0</sub> ±SE	32.198±2.254	51.740±2.024	45.190±4.813	38.580±3.803
W <sub>f</sub> ±SE	1145.270±29.826	895.200±11.640	1520.000±117.200	1032.000±5.029
k±SE	0.024±0.0007	0.049±0.0010	1.891±0.085	0.031±0.004
m±SE	-	-	-	0.270±0.143
b±SE	-	-	94.100±7.469	-
W <sub>i</sub>	421.321	447.600	392.640	425.816
T <sub>i</sub>	53.041	56.965	50.498	53.745
RMSE	86.538	86.755	86.767	86.497
AIC	31319.37	31332.69	31332.56	31317.83
BIC	31342.92	31356.24	31361.99	31343.26
R <sup>2</sup> <sub>Adj</sub>	89.822	89.771	89.775	89.832

<sup>1</sup> W<sub>0</sub>: initial weight (g); W<sub>f</sub>: final weight (g); k: coefficient of relative growth or maturing index (g per d); b: age at approximately half maximum body weight; m: shape parameter; SE: Standard error; W<sub>i</sub>: weight at the inflection point (g); T<sub>i</sub>: age at the inflection point (d); RSME: root mean square error; BIC: bayesian information criterion; AIC: akaike information criterion and R<sup>2</sup><sub>Adj</sub>: adjusted coefficient of determination.

**Table 4** Effect of energy and protein levels on growth curve parameters and absolute growth rate in different ages

Parameters <sup>1</sup>	ME(k cal/kg)			P-value	CP (%)			P-value	ME × CP	SEM
	2600	2800	3000		17	19	21		P-value	
W <sub>0</sub>	38.745	39.297	40.894	0.6577	39.175	38.856	40.906	0.6641	0.9422	13.338
W <sub>f</sub>	1070.95	1156.59	1193.28	0.1107	1043.26 <sup>b</sup>	1156.66 <sup>ab</sup>	1219.91 <sup>a</sup>	0.0120	0.0934	325.667
k	0.042 <sup>a</sup>	0.036 <sup>ab</sup>	0.029 <sup>b</sup>	0.0018	0.40	0.035	0.032	0.0890	0.0075	0.019
m	0.474	0.438	0.337	0.5181	0.426	0.379	0.444	0.8630	0.8892	0.676
W <sub>i</sub>	407.19 <sup>b</sup>	461.12 <sup>a</sup>	462.06 <sup>a</sup>	0.0233	407.42 <sup>b</sup>	436.29 <sup>ab</sup>	486.66 <sup>a</sup>	0.0024	0.0064	124.100
T <sub>i</sub>	53.601 <sup>b</sup>	62.082 <sup>ab</sup>	72.245 <sup>a</sup>	0.0067	62.200	62.726	63.002	0.9902	0.0493	31.852
AGR <sub>14</sub>	4.792 <sup>b</sup>	4.970 <sup>ab</sup>	5.513 <sup>a</sup>	0.0079	4.598 <sup>b</sup>	5.071 <sup>ab</sup>	5.606 <sup>a</sup>	0.0004	0.0002	0.539
AGR <sub>21</sub>	5.365 <sup>b</sup>	5.456 <sup>ab</sup>	6.041 <sup>a</sup>	0.0485	5.172 <sup>b</sup>	5.586 <sup>ab</sup>	6.195 <sup>a</sup>	0.0029	0.0216	0.569
AGR <sub>28</sub>	6.408 <sup>b</sup>	6.713 <sup>ab</sup>	7.336 <sup>a</sup>	0.0116	6.358 <sup>b</sup>	6.595 <sup>b</sup>	7.503 <sup>a</sup>	0.0013	0.0027	0.712
AGR <sub>35</sub>	6.901 <sup>b</sup>	7.469 <sup>ab</sup>	7.865 <sup>a</sup>	0.0184	6.982 <sup>b</sup>	7.216 <sup>b</sup>	8.037 <sup>a</sup>	0.0066	0.0100	0.778
AGR <sub>42</sub>	7.729 <sup>b</sup>	8.187 <sup>ab</sup>	8.818 <sup>a</sup>	0.0521	7.746 <sup>b</sup>	8.040 <sup>ab</sup>	8.948 <sup>a</sup>	0.0232	0.0595	1.042
AGR <sub>49</sub>	8.289 <sup>b</sup>	8.830 <sup>ab</sup>	9.932 <sup>a</sup>	0.0141	8.243 <sup>b</sup>	9.011 <sup>ab</sup>	9.797 <sup>a</sup>	0.0236	0.0345	1.296
AGR <sub>56</sub>	8.825 <sup>b</sup>	9.674 <sup>ab</sup>	10.208 <sup>a</sup>	0.0343	8.744 <sup>b</sup>	9.535 <sup>ab</sup>	10.428 <sup>a</sup>	0.0093	0.0254	1.234
AGR <sub>63</sub>	8.997 <sup>b</sup>	9.838 <sup>ab</sup>	10.564 <sup>a</sup>	0.0430	8.950 <sup>b</sup>	9.773 <sup>ab</sup>	10.675 <sup>a</sup>	0.0241	0.0492	1.443
AGR <sub>70</sub>	8.206 <sup>b</sup>	8.605 <sup>ab</sup>	10.219 <sup>a</sup>	0.0141	8.239 <sup>b</sup>	8.828 <sup>ab</sup>	9.963 <sup>a</sup>	0.0487	0.0339	1.649
AGR <sub>77</sub>	6.488 <sup>b</sup>	8.218 <sup>ab</sup>	8.541 <sup>a</sup>	0.0368	6.527 <sup>b</sup>	8.093 <sup>ab</sup>	8.627 <sup>a</sup>	0.0392	0.0889	1.977
AGR <sub>84</sub>	5.957 <sup>b</sup>	7.441 <sup>ab</sup>	7.824 <sup>a</sup>	0.0411	5.818 <sup>b</sup>	7.333 <sup>ab</sup>	8.070 <sup>a</sup>	0.0155	0.0498	1.800
AGR <sub>91</sub>	5.392 <sup>b</sup>	6.765 <sup>ab</sup>	7.222 <sup>a</sup>	0.0479	5.157 <sup>b</sup>	6.853 <sup>ab</sup>	7.369 <sup>a</sup>	0.0139	0.0495	1.787
AGR <sub>98</sub>	4.594 <sup>b</sup>	6.041 <sup>ab</sup>	6.691 <sup>a</sup>	0.0185	4.746 <sup>b</sup>	6.057 <sup>ab</sup>	6.523 <sup>a</sup>	0.0474	0.0631	1.726

<sup>1</sup> W<sub>0</sub>: initial weight (g); W<sub>f</sub>: final weight (g); k: coefficient of relative growth or maturing index (g per d); b: age at approximately half maximum body weight; m: shape parameter; SE: Standard error; W<sub>i</sub>: weight at the inflection point (g); T<sub>i</sub>: age at the inflection point (d) and AGR: absolute growth rate in different age (g/d). Means within a row with different superscript are significantly different (P<0.05). SEM: standard error of the means.

However, the mean absolute growth rate for birds that received a diet with 2800 kcal of ME/kg and 19% CP was not significant with a high level of energy (3000 kcal/kg) and protein (21%).

Table 5 shows the interaction effect of ME and CP on

growth curve parameters and Absolute growth rate in different ages. The interaction effect of ME and CP was significant on the k and W<sub>i</sub> parameters (P<0.05) and the differences of W<sub>0</sub>, W<sub>f</sub>, m, and T<sub>i</sub> were not significant (P>0.05).

**Table 5** Interaction effect of energy and protein on growth curve parameters and Absolute growth rate in different ages

ME (kcal/kg)	2600			2800			3000		
	17	19	21	17	19	21	17	19	21
CP (%)	17	19	21	17	19	21	17	19	21
<b>Parameters<sup>1</sup></b>									
W <sub>0</sub>	39.481	38.663	38.092	37.768	37.773	42.352	40.275	40.134	42.274
W <sub>f</sub>	969.6	1101.7	1141.6	1040.4	1177.1	1252.4	1119.8	1194.2	1265.8
k	0.0490 <sup>a</sup>	0.0420 <sup>ab</sup>	0.0352 <sup>ab</sup>	0.0349 <sup>ab</sup>	0.0360 <sup>ab</sup>	0.0363 <sup>ab</sup>	0.0358 <sup>ab</sup>	0.0271 <sup>b</sup>	0.0248 <sup>b</sup>
m	0.5515	0.4670	0.4041	0.3621	0.3506	0.6012	0.3650	0.3200	0.3270
W <sub>i</sub>	379.51 <sup>b</sup>	411.75 <sup>ab</sup>	430.30 <sup>ab</sup>	415.49 <sup>ab</sup>	438.94 <sup>ab</sup>	528.94 <sup>a</sup>	427.26 <sup>ab</sup>	458.18 <sup>ab</sup>	500.73 <sup>ab</sup>
T <sub>i</sub>	50.47 <sup>b</sup>	54.55 <sup>ab</sup>	55.79 <sup>ab</sup>	53.67 <sup>ab</sup>	66.43 <sup>ab</sup>	66.15 <sup>ab</sup>	82.47 <sup>a</sup>	67.20 <sup>ab</sup>	67.07 <sup>ab</sup>
AGR <sub>14</sub>	4.244 <sup>b</sup>	4.980 <sup>b</sup>	5.152 <sup>b</sup>	4.907 <sup>b</sup>	4.981 <sup>b</sup>	5.023 <sup>b</sup>	4.644 <sup>b</sup>	5.251 <sup>b</sup>	6.643 <sup>a</sup>
AGR <sub>21</sub>	4.788 <sup>b</sup>	5.305 <sup>ab</sup>	6.001 <sup>ab</sup>	5.266 <sup>ab</sup>	5.434 <sup>ab</sup>	5.937 <sup>ab</sup>	5.461 <sup>ab</sup>	6.017 <sup>ab</sup>	6.646 <sup>a</sup>
AGR <sub>28</sub>	5.832 <sup>b</sup>	5.906 <sup>b</sup>	7.487 <sup>ab</sup>	6.036 <sup>b</sup>	6.814 <sup>ab</sup>	7.288 <sup>ab</sup>	7.206 <sup>ab</sup>	7.066 <sup>ab</sup>	7.735 <sup>a</sup>
AGR <sub>35</sub>	6.408 <sup>b</sup>	6.685 <sup>b</sup>	7.610 <sup>ab</sup>	7.519 <sup>ab</sup>	7.197 <sup>ab</sup>	7.691 <sup>ab</sup>	7.020 <sup>ab</sup>	7.766 <sup>ab</sup>	8.809 <sup>a</sup>
AGR <sub>42</sub>	7.321 <sup>b</sup>	7.381 <sup>b</sup>	8.486 <sup>ab</sup>	7.918 <sup>ab</sup>	8.218 <sup>ab</sup>	8.425 <sup>ab</sup>	7.999 <sup>ab</sup>	8.521 <sup>ab</sup>	9.934 <sup>a</sup>
AGR <sub>49</sub>	7.778 <sup>b</sup>	7.980 <sup>ab</sup>	9.110 <sup>ab</sup>	8.016 <sup>ab</sup>	9.230 <sup>ab</sup>	9.243 <sup>ab</sup>	8.934 <sup>ab</sup>	9.824 <sup>ab</sup>	11.038 <sup>a</sup>
AGR <sub>56</sub>	7.911 <sup>b</sup>	9.002 <sup>ab</sup>	9.561 <sup>ab</sup>	8.702 <sup>ab</sup>	10.086 <sup>ab</sup>	10.233 <sup>ab</sup>	9.617 <sup>ab</sup>	9.517 <sup>ab</sup>	11.489 <sup>a</sup>
AGR <sub>63</sub>	7.845 <sup>b</sup>	9.251 <sup>ab</sup>	9.896 <sup>ab</sup>	9.360 <sup>ab</sup>	10.070 <sup>ab</sup>	10.084 <sup>ab</sup>	9.646 <sup>ab</sup>	9.999 <sup>ab</sup>	12.047 <sup>a</sup>
AGR <sub>70</sub>	7.047 <sup>b</sup>	8.314 <sup>ab</sup>	9.257 <sup>ab</sup>	8.658 <sup>ab</sup>	8.407 <sup>ab</sup>	8.751 <sup>ab</sup>	9.013 <sup>ab</sup>	9.762 <sup>ab</sup>	11.882 <sup>a</sup>
AGR <sub>77</sub>	5.419	6.716	7.331	7.335	8.731	8.588	6.826	8.834	9.964
AGR <sub>84</sub>	4.547 <sup>b</sup>	6.166 <sup>ab</sup>	7.158 <sup>ab</sup>	6.732 <sup>ab</sup>	7.482 <sup>ab</sup>	8.109 <sup>ab</sup>	6.176 <sup>ab</sup>	8.351 <sup>ab</sup>	8.945 <sup>a</sup>
AGR <sub>91</sub>	3.885 <sup>b</sup>	5.559 <sup>ab</sup>	6.734 <sup>ab</sup>	5.881 <sup>ab</sup>	6.737 <sup>ab</sup>	7.676 <sup>ab</sup>	5.706 <sup>ab</sup>	8.263 <sup>a</sup>	7.698 <sup>ab</sup>
AGR <sub>98</sub>	3.679	4.525	5.577	4.971	6.090	7.062	5.587	7.557	6.930

<sup>1</sup> W<sub>0</sub>: initial weight (g); W<sub>f</sub>: final weight (g); k: coefficient of relative growth or maturing index (g per d); b: age at approximately half maximum body weight; m: shape parameter; SE: Standard error; W<sub>i</sub>: weight at the inflection point (g); T<sub>i</sub>: age at the inflection point (d) and AGR: absolute growth rate in different age (g/d). Means within a row with different superscript are significantly different (P<0.05). SEM: standard error of the means.

The mean of the mature index (k) for birds fed with a diet containing 2600 kcal of ME/kg and 16% CP was higher than other treatments, but only its difference was significant than diets with 3000 kcal of ME/kg and 19 and 21% CP. The highest value of weight at the inflection point was birds fed with a diet containing 3000 kcal of ME/kg and 21% CP that was significantly higher than the 2600 kcal of ME/kg and 17% CP diet. Interaction of ME and CP had a significant effect on absolute growth rate in different ages except for absolute growth rate at 77 and 98 days of age (P<0.05). In all ages, the mean absolute growth rate for birds fed with a diet containing 2600 kcal of ME/kg and 17% CP was lower than other diets.

Mathematical models are used for describing the growth curve, have the main role in poultry improvement programs. The importance of these models and use of models in poultry have been reported in other studies (Anthony *et al.* 1991; Aggrey, 2002) especially when the growth is related to the nutritional components (Aggrey, 2004).

Cumulative feed consumption until slaughter weight in poultry depends on growth rate and the shape of the growth curve, therefore the use of a mathematical model in combination with feed intake data can be useful in bioeconomic studies (Pasternak and Shalev, 1983).

In this study, the Richards model was selected as the best model to describe the growth curve of Khazak chickens. In agreement with this finding, the superiority of the Richards models to describe the growth curve of broiler and native chickens that fed with different levels of ME was reported (Tompic *et al.* 2011; Moharrery and Mirzaei, 2014).

Overestimation of W<sub>0</sub> parameter with Logistic model and k, W<sub>f</sub> parameters with Lopez model was reported in other studies of the growth curve in broiler chickens (Moharrery and Mirzaei, 2014; Masoudi and Azarfah, 2017; Faraji-Arough *et al.* 2019) which was in agreement with our results. In growth curve studies in native chickens, various models were introduced as the best model to describe the growth curve such as the Gompertz model for Nigerian

native chickens (Adenaike *et al.* 2017), slow-growing chickens in China (Zhao *et al.* 2015), Poland medium-growing chickens (Michalczyk *et al.* 2016), Logistic model for slow-growing chickens in the organic system (Eleroglu *et al.* 2014), Von Bertalanffy for Korean native chickens (Manjula *et al.* 2016), and Lopez for Khazak chickens (Faraji-Arough *et al.* 2019). Although, Faraji-Arough *et al.* (2019) introduced the Lopez model for Khazak chickens that in contrast with present results. The data of body weight in this study was related to chicks that fed with diets with different levels of ME and CP, but the used data in a study by Faraji-Arough *et al.* (2019) was collected from chicks that used the same diet (2800 kcal of ME/kg and 16% CP). Difference in the feeding of chickens in the present study and study by Faraji-Arough *et al.* (2019) can be a reason for the report of different appropriate models to describe the growth curve in Khazak chickens.

Our findings indicate that the maturity index ( $k$ ) decreased linearly with increasing of ME and CP levels (Table 4) whereas the final weight ( $W_f$ ) increased with the increment of ME and CP levels. Inverse association between  $W_f$  and  $k$  parameter was reported in other studies (Adenaike *et al.* 2017; Faraji-Arough *et al.* 2019) which indicate early maturing in birds result in the slow rate of growth in the first weeks of age. Furthermore, a correlation between  $W_f$  and  $k$  was reported negative ( $>-0.90$ , Masoudi and Azarfar, 2017; Faraji-Arough *et al.* 2019), therefore the chicks with a lower  $k$  value will have a higher  $W_f$ .

Although, the effect of different levels of ME and CP on growth performance of native chickens was reported in some studies (Nguyen *et al.* 2010; Haunshi *et al.* 2012; Al-Khalifa and Al-Nasser, 2012; Miah *et al.* 2014; Maliwan *et al.* 2019). However, in these studies, growth performance was considered as body weight gain, feed intake, and feed conversion ratio. In a study on broiler chickens, some of the growth curve parameters especially  $W_i$ ,  $W_0$ , growth rate on days 28, 35, and 42 were significantly affected by different levels of corn bran of diet (Masoudi and Azarfar, 2017).

Effect of ME and CP on growth curve parameter of French guinea fowl by Gompertz model was studied by Nahashon *et al.* (2010) and no significant effect of ME on growth curve parameter except for  $W_0$  was reported that was contrary to the results of the present study, but similar to our result, the effect of CP was significant on  $W_i$  and growth rate. It has also been reported that the final body weight of guinea fowl broiler fed diet containing 3100 or 3150 kcal of ME /kg was significantly higher than those fed the 3,050 kcal of ME /kg diet (Nahashon *et al.* 2005) that was opposite with our finding.

Based on the Richards model, the decreased dietary ME concentration cause a linear decrease in  $W_f$  in broiler chickens ( $P<0.05$ ) (Moharrery and Mirzaei, 2014) but no effect of ME on  $W_f$  and other growth parameters in native chickens was reported that was similar with present results. Adaptation of native chickens in consuming feed of lower ME concentration can be a reason for the lack of significant effect of dietary ME on  $W_f$ . On the other hand, the native chickens do not need high energy concentration in their feed due to having a slower growth rate and small size at maturity. Therefore, their feed intake was reduced with increasing ME in diet, which can be another reason for the non-significant effect of ME on  $W_f$  (Moharrery and Mirzaei, 2014).

After hatch, the growth of birds is accelerated till a certain age which this time showed by age at the inflection point ( $T_i$ ) and growth rate of bird in this maximum in this phase. The growth of birds decreased gradually after this age to achieve their mature weight (Sakomura and Rostagno, 2016). Our finding shows that this age for chicks fed with a diet containing a high level of ME significantly increased than a low level of ME. The increase of  $T_i$  provides an opportunity for the chicks to gain more body weight ( $W_i$ ). Similar to the present results, a significant effect of ME and CP on the growth curve parameter of Broiler chickens by Logistic and Gompertz was reported (Koushandeh *et al.* 2019). An increasing trend of  $W_i$ ,  $T_i$ , and absolute growth rate and decreasing trend of  $k$  parameter was reported with increasing the dietary CP level in gibel crap (Yun *et al.* 2015) that was similar with the present result.

## CONCLUSION

Based on goodness of fit criteria, the Richards model was selected as the best model to predict the growth curve parameters of Khazak chicks fed with a diet containing different levels of ME and CP. The ME and CP levels and their interaction had a significant effect on  $W_i$  and absolute growth rate in different ages. Also, the  $k$  and  $T_i$  parameter was affected significantly by ME levels and the final weight ( $W_f$ ) of chicks fed with a diet containing a high level of CP was significantly higher than those fed with a low level of CP diet. The difference of many studied parameters was not significant between 2800 and 3000 kcal of ME /kg and 19 with 21% CP, thus diet with 2800 kcal of ME/kg and 19% CP can be suggested for Khazak chickens during 7 to 98 days of age to the optimum improvement of the growth curve parameters.

On other hand, the results of this study can be useful in nutritional management and can help the producer to formulate the best diet when the growth rate is at its maximum.

## ACKNOWLEDGEMENT

The authors would like to thank the Center of Domestic Animals Institute, Research Institute of Zabol, Zabol, Iran for cooperation in performing this research.

## REFERENCES

- Abbas A.A., Yosif A.A., Shukur A.M. and Ali F.H. (2014). Effect of Genotypes, Storage periods and feed additives in broiler breeder diets on embryonic and hatching indicators and chicks blood parameters. *Sci. Agric.* **7**, 44-48.
- Adenaike A., Akpan U., Udoh J.E., Wheto M., Durosaro S.O., Sanda A.J. and Ikeobi C.O.N. (2017). Comparative evaluation of growth functions in three broiler strains of Nigerian chickens. *Pertanika J. Trop. Agric. Sci.* **40**, 611-620.
- Aggrey S.E. (2002). Comparison of three nonlinear and spline regression models for describing chicken growth curves. *Poult. Sci.* **81**, 1782-1788.
- Aggrey S.E. (2004). Modeling the effect of nutritional status on pre-asymptotic and relative growth rates in a random bred chicken population. *J. Anim. Breed. Genet.* **121**, 260-268.
- Al-Khalifa H. and Al-Nasser A. (2012). Effect of different protein levels on Arabi chicken performance. *Int. J. Poult. Sci.* **11**, 706-709.
- Anthony N.B., Emmerson D.A., Nestor K.E. and Bacon W.L. (1991). Comparison of growth curves of weight selected populations of turkey, quail and chickens. *Poult. Sci.* **70**, 13-19.
- Durunna C.S., Udedibie A.B.I. and Uchegbu M.C. (2005). Effect of dietary inclusion of Anthonata macrophyla meal on the performance of broiler starter chicks. *Nigerian J. Anim. Prod.* **32**, 268-273.
- Dyubele N.L., Muchenje V., Nkukwana T.T. and Chimonyo M. (2010). Consumer sensory characteristics of broiler and indigenous chicken meat: A South African example. *Food Qual. Prefer.* **21**, 815-819.
- Eleroglu H., Yıldırım A., Şekeroğlu A., Çoksöyler F.N. and Duman M. (2014). Comparison of growth curves by growth models in slow-growing chicken genotypes raised the organic system. *Int. J. Agric. Biol.* **16**, 529-535.
- Faraji-Arough H., Rokouei M., Maghsoudi A. and Mehri M. (2019). Evaluation of non-linear growth curves models for native slow-growing Khazak Chickens. *Poult. Sci. J.* **7**, 25-32.
- Haunshi S., Panda A.K., Rajkumar U., Padhi M.K., Niranjana M. and Chatterjee R.N. (2012). Effect of feeding different levels of energy and protein on performance of Aseel breed of chicken during juvenile phase. *Trop. Anim. Health Prod.* **44**, 1653-1658.
- Iranian Council of Animal Care. (1995). Guide to the Care and Use of Experimental Animals, vol. 1. Isfahan University of Technology, Isfahan, Iran.
- Koushandeh A., Chamani M., Yaghobfar A., Sadeghi A.A. and Baneh H. (2019). Comparison of the accuracy of nonlinear models and artificial neural network in the performance prediction of Ross 308 broiler chickens. *Poult. Sci. J.* **7**, 151-161.
- Lopez S., France J., Gerrits W.J.J., Dhanoa M.S., Humphries D.J. and Dijkstra J. (2000). A generalized michaelis-menten equation for the analysis of growth. *J. Anim. Sci.* **78**, 1816-1828.
- Maliwan P., Molee W. and Khempaka S. (2019). Response of Thai indigenous crossbred chickens to various dietary protein levels at different ages. *Trop. Anim. Health Prod.* **51**, 1427-1439.
- Manjula P., Park H.B., Yoo J., Wickramasuriya S., Seo D.W., Choi N.R., Kim C.D., Kang B.S., Oh K.S. and Sohn S.H. (2016). Comparative study on growth patterns of 25 commercial strains of Korean native chicken. *Korean J. Poult. Sci.* **43**, 1-14.
- Masoudi A. and Azarfar A. (2017). Comparison of nonlinear models describing growth curves of broiler chickens fed on different levels of corn bran. *Int. J. Avian Wildl. Biol.* **2**, 1-7.
- Miah M.Y., Chowdhury S.D., Bhuiyan A.K.F.H. and Ali M.S. (2014). Effect of different levels of dietary energy on growth performance of indigenous Desi chicks reared in confinement up to target weight of 950 g. *Livest. Res. Rural Dev.* Available at: <http://www.lrrd.org/lrrd26/7/miah26124.html>.
- Michalczuk M., Damaziak K. and Goryl A. (2016). Sigmoid models for the growth curves in medium-growing meat type chickens, raised under semi-confined conditions. *Ann. Anim. Sci.* **16**, 65-77.
- Mohammed F.A. (2015). Comparison of three nonlinear functions for describing chicken growth curves. *Sci. Agric.* **9**, 120-123.
- Moharrery A. and Mirzaei M. (2014). Growth characteristics of commercial broiler and native chickens as predicted by different growth functions. *J. Anim. Feed Sci.* **23**, 82-89.
- Nahashon S.N., Adefope N., Amenyenu A. and Wright D. (2005). Effect of dietary metabolizable energy and crude protein concentrations on growth performance and carcass characteristics of French guinea broilers. *Poult. Sci.* **84**, 337-344.
- Nahashon S.N., Aggrey S.E., Adefope N.A., Amenyenu A. and Wright D. (2010). Gompertz-Laird model prediction of optimum utilization of crude protein and metabolizable energy by French guinea fowl broilers. *Poult. Sci.* **89**, 52-57.
- Nguyen T.V., Bunchasak C. and Chantsavang S. (2010). Effects of dietary protein and energy on growth performance and carcass characteristics of Betong chickens (*Gallus domesticus*) during growing period. *Int. J. Poult. Sci.* **9**, 468-472.
- Norris D., Ngambi J.W., Benyi K., Makgahlele M.L., Shimelis H.A. and Nesamvuni E.A. (2007). Analysis of growth curves of indigenous male Venda and Naked Neck chickens. *South African J. Anim. Sci.* **37**, 21-26.
- Padhi M.K. (2016). Importance of indigenous breeds of chicken for rural economy and their improvements for higher production performance. *Scientifica.* **6**, 1-9.
- Pasternak H. and Shalev B.A. (1983). Genetic-economic evaluation of traits in a broiler enterprise: Reduction of food intake due to increased growth rate. *British Poult. Sci.* **24**, 531-536.
- Perween S., Kumar K., Chandramoni S.K., Singh P.K., Kumar M.



- and Dey A. (2016). Effect of feeding different dietary levels of energy and protein on growth performance and immune status of Vanaraja chicken in the tropic. *Vet. World*, **9**, 893-899.
- Pinheiro J., Bates D., DebRoy S., Sarkar D. and Core Team R. (2014) *Nlme: Linear and Nonlinear Mixed Effects Models*. R package version 3.1-117. Available at: <http://cran.r-project.org/web/packages/nlme/index.html>.
- Russo T., Baldi P., Parisi A., Magnifico G., Mariani S. and Cautaudella S. (2009). Levy processes and stochastic von Bertalanffy models of growth, with application to fish population analysis. *J. Theor. Biol.* **258**, 521-529.
- Sakomura N.K. and Rostagno H.S. (2016). Métodos de Pesquisa em Nutrição de Monogástricos. Funep-Unesp, Jaboticabal.
- SAS Institute. (2008). SAS<sup>®</sup>/STAT Software, Release 9.2. SAS Institute, Inc., Cary, NC, USA.
- Shehbaz M. and Khan S.H. (2008). Effects of different energy protein ratio on the performance of desi native chickens during growing phase. *Asian J. Poult. Sci.* **2**, 42-47.
- Teleken J.T., Galvão A.C. and Robazza W.D.S. (2017). Comparing non-linear mathematical models to describe growth of different animals. *Acta Sci.* **39**, 73-81.
- Tompic T., Dobša J., Legen S., Tompic N. and Medic H. (2011). Modeling the growth pattern of in-season and off-season Ross 308 broiler breeder flocks. *Poult. Sci.* **90**, 2879-2887.
- Wattanachant S. (2008). Factors affecting the quality characteristics of Thai indigenous chicken meat. *Suranaree J. Sci. Technol.* **15**, 317-322.
- Wijten P.J.A., Lemme A. and Langhout D.J. (2004). Effects of different dietary ideal protein levels on male and female broiler performance during different phases of life: Single phase effects, carryover effects, and interactions between phases. *Poult. Sci.* **83**, 2005-2015.
- Yun B., Yu X., Xue M., Liu Y., Wang J., Wu X., Han F. and Liang X. (2015). Effects of dietary protein levels on the long-term growth response and fitting growth models of gibel carp (*Carassius auratus gibelio*). *Anim. Nutr.* **1**, 70-76.
- Zhao Z., Li S., Huang H., Li C., Wang Q. and Xue L. (2015). Comparative study on growth and developmental model of indigenous chicken breeds in China. *Open J. Anim. Sci.* **5**, 219-223.
-