

## The Review of Almost Periodic Solutions to a Stochastic Differential Equation

S. Ahmadi <sup>a,\*</sup>

<sup>a</sup>*Tehran North Branch, Azad University, Tehran, Iran.*

---

**Abstract.** This paper proves the existence and uniqueness of quadratic mean almost periodic mild solutions for a class of stochastic differential equations in a real separable Hilbert space. The main technique is based upon an appropriate composition theorem combined with the Banach contraction mapping principle and an analytic semigroup of linear operators.

---

Received: 12 August 2015, Revised: 20 November 2015, Accepted: 16 December 2015.

**Keywords:** Stochastic differential equations, Quadratic mean almost periodic, Analytic semigroups of linear operators.

### Index to information contained in this paper

- 1 Introduction
- 2 Preliminaries
- 3 Main Results
- 4 Conclusion

### 1. Introduction

The history of stochastic differential equations (SDEs) can be seen to have started from the classic paper of Einstein, where he presented a mathematical connection between microscopic random motion of particles and the macroscopic diffusion equation. Qualitative properties such as existence, uniqueness, controllability and stability for various stochastic differential systems have been extensively studied by many researchers, see for instance [1] and the references therein. On the other hand, the existence of almost periodic solutions for deterministic differential equations have been considerably investigated in lots of publications because of its significance and applications in physics, mechanics and mathematical biology, see for example [2-4] and the references therein. Recently, the concept of quadratic mean almost periodicity was introduced by Bezandry and Diagana [5]. In [5], such a concept was

---

\* Corresponding author. Email: s.ahmadi.ch@gmail.com

subsequently applied to proving the existence and uniqueness of a quadratic mean almost periodic solution to the following stochastic differential equations

$$dy(t) = Mx(t)dt + N(t, y(t))dt + S(t, y(t))dw(t), t \in \mathbb{R}$$

where  $M : D(M) \subset L^2(\Omega; \mathbb{H}) \rightarrow L^2(\Omega; \mathbb{H})$  for  $t \in \mathbb{R}$  is a densely defined closed linear operators,  $N : \mathbb{R} \times L^2(\Omega; \mathbb{H}) \rightarrow L^2(\Omega; \mathbb{H})$  and  $S : \mathbb{R} \times L^2(\Omega; \mathbb{H}) \rightarrow L^2(\Omega; L_0^2)$  are jointly continuous satisfying some additional conditions, and  $w(t)$  is a Wiener process.

Bezandry and Diagana [6] have studied the existence and uniqueness of a quadratic mean almost periodic solution to a non-autonomous semi-linear stochastic differential equations such as

$$dy(t) = M(t)y(t)dt + N(t, y(t))dt + S(t, y(t))dw(t), t \in \mathbb{R}$$

where  $A(t)$  for  $t \in \mathbb{R}$  is a family of densely defined closed linear operators satisfying the so-called Acquistapace-Terreni condition in [7],  $N : \mathbb{R} \times L^2(\Omega; \mathbb{H}) \rightarrow L^2(\Omega; \mathbb{H})$  and  $S : \mathbb{R} \times L^2(\Omega; \mathbb{H}) \rightarrow L^2(\Omega; L_0^2)$  are jointly continuous satisfying some additional conditions, and  $w(t)$  is a Wiener process. And Bezandry in [8] has considered the existence of quadratic mean almost periodic solutions to a semi-linear functional stochastic integro-differential equations in the form

$$y'(t) = My(t) + \int_{-\infty}^t C(t-u)S(u, y(u))dw(u) + \int_{-\infty}^t B(t-u)N_2(u, y(u))du + N_1(t, y(t)),$$

where  $t \in \mathbb{R}$ ,  $M : D(M) \subset L^2(\Omega; \mathbb{H}) \rightarrow L^2(\Omega; \mathbb{H})$  is a densely defined closed (possibly unbounded) linear operator;  $B$  and  $C$  are convolution-type kernels in  $L^1(0, \infty)$  and  $L^2(0, \infty)$ , respectively, satisfying Assumptions 3.2 in [1];  $N_1, N_2 : \mathbb{R} \times L^2(\Omega; \mathbb{H}) \rightarrow L^2(\Omega; \mathbb{H})$  and  $G : \mathbb{R} \times L^2(\Omega; \mathbb{H}) \rightarrow L^2(\Omega; L_0^2)$  are jointly continuous functions. For more results on this topic, we refer the reader to the papers [9-12] and the references therein.

Motivated by the above mentioned works [5,13,8], the main purpose of this paper is to deal with the existence and uniqueness of quadratic mean almost periodic solutions to a class of neutral stochastic functional differential equations in the abstract form

$$d[y(t) - h(t, x(t))] = My(t)dt + S(t, y(t))dw(t), t \in \mathbb{R} \quad (1)$$

where  $M : D(M) \subset L^2(\Omega; \mathbb{H}) \rightarrow L^2(\Omega; \mathbb{H})$  is the infinitesimal generator of an analytic semigroup of linear operators  $\{T(t)\}_{t \geq 0}$  on  $L^2(\Omega; \mathbb{H})$ ,  $g : \mathbb{R} \times L^2(\Omega; \mathbb{H}) \rightarrow L^2(\Omega; \mathbb{H}_\alpha)$  and  $G : \mathbb{R} \times L^2(\Omega; \mathbb{H}) \rightarrow L^2(\Omega; L_2^0)$  are jointly continuous functions,  $w(t)$  is a Brownian motion. The main technique is based upon an appropriate composition theorem combined with the Banach contraction mapping principle and an analytic semigroup of linear operators. The obtained result can be seen as a contribution to this emerging field.

The rest of this paper is organized as follows: In Sect. 2, in this paper some basic definitions, lemmas and preliminary facts which will be need in the sequel. my main result and its proofs are arranged in Sect. 3. In the last section, conclusions and an example have been given to illustrate main result.

## 2. Preliminaries

This section is mainly concerned with some definitions, lemmas and preliminary facts which are used in what follows. For more details on this section, we refer the reader to [5,6,9,10]. Throughout the paper,  $(\mathbb{H}, \|\cdot\|)$  and  $(\mathbb{K}, \|\cdot\|)$  denote two real Hilbert spaces. Let  $\mathbb{P}$  be a complete probability space. We let  $L_2(\mathbb{K}, \mathbb{H})$  denote the space of all Hilbert–Schmidt operators  $\Phi : \mathbb{K} \rightarrow \mathbb{H}$ , equipped with the Hilbert–Schmidt norm  $\|\cdot\|_2$ . For a symmetric nonnegative operator  $Q \in L_2(\mathbb{K}, \mathbb{H})$  with finite trace we suppose that  $\{\omega(t) : t \in \mathbb{R}\}$  is a  $Q$ -Wiener process defined on  $(\Omega, \mathcal{F}, \mathbb{P})$  and with values in  $\mathbb{K}$ . So, actually,  $\omega$  can be obtained as follows: let  $\omega_i(t), t \in \mathbb{R}, i = 1, 2$ , be independent  $\mathbb{K}$ -valued  $Q$ -Wiener processes, then

$$\omega(t) = \begin{cases} \omega_1(t) & \text{if } t \geq 0 \\ \omega_2(-t) & \text{if } t \leq 0 \end{cases}$$

is  $Q$ -Wiener process with  $\mathbb{R}$  as time parameter.  $\mathcal{N}_t = \sigma\{\omega(s) : s \leq t\}$  is the  $\sigma$ -algebra generated by  $\omega$ .

The collection of all strongly measurable, square integrable,  $\mathbb{H}$ -valued random variable, denoted by  $L^2(\Omega; \mathbb{R})$ , is a Banach space equipped with norm  $\|y\|_{L^2(\Omega; \mathbb{H})} = (E\|y\|^2)^{\frac{1}{2}}$ , where the expectation  $E$  is defined  $E[y] = \int_{\Omega} x(\omega) dP(\omega)$ .

Let  $\mathbb{K}_0 = Q^{\frac{1}{2}}\mathbb{K}$  and  $L_0^2 = L_2(\mathbb{K}_0, \mathbb{H})$  with respect to the norm

$$\|\Phi\|_{L_0^2}^2 = \|\phi Q^{\frac{1}{2}}\|_2^2 = \text{Tr}(\phi Q \phi^*).$$

Let  $0 \in \rho(A)$  where  $\rho(M)$  is the resolvent of  $M$ . Then for  $0 < \alpha \leq 1$ , it is possible to define the fractional power  $(-M)^{\alpha}$ , as a closed linear operator on its domain  $D((-M)^{\alpha})$ . Furthermore, the subspace  $D((-M)^{\alpha})$  is dense in  $L^2(\Omega; \mathbb{H})$  and the expression

$$\|y\|_{\alpha} = \|(-M)^{\alpha}x\|_{L^2(\Omega; \mathbb{H})}, x \in D((-M)^{\alpha})$$

defines a norm on  $D((-M)^{\alpha})$ . we denote by  $L^2(\Omega; \mathbb{H}_{\alpha})$  the Banach space  $D((-M)^{\alpha})$  with norm  $\|y\|_{\alpha}$ .

The following properties hold by [14].

**Proposition 2.1.** suppose  $0 < \gamma \leq \mu \leq 1$ . Then the following properties hold:

- (i)  $L^2(\Omega; \mathbb{H}_{\mu})$  is a Banach space and  $L^2(\Omega; \mathbb{H}_{\mu}) \rightarrow L^2(\Omega; \mathbb{H}_{\gamma})$  is continuous.
- (ii) the function  $s \rightarrow (-M)^{\mu}T(s)$  is continuous on  $(0, \infty)$  and there exists  $A_{\mu} > 0$  such that  $\|(-M)^{\mu}T(t)\| \leq A_{\mu}e^{-\delta t}t^{-\mu}$  for each  $t > 0$ .
- (iii) For each  $y \in D((-M)^{\mu})$  and  $t \geq 0$ ,  $(-M)^{\mu}T(t)x = T(t)(-M)^{\mu}x$ .
- (iv)  $(-M)^{-\mu}$  is a bounded linear operator in  $L^2(\Omega; \mathbb{H})$  with  $D((-M)^{\mu}) = \text{Im}((-M)^{-\mu})$ .

In the following results and definitions, we let  $(\mathbb{Y}, \|\cdot\|_{\mathbb{Y}})$ ,  $(\mathbb{X}, \|\cdot\|_{\mathbb{X}})$  and  $(\mathbb{Z}, \|\cdot\|_{\mathbb{Z}})$  be Banach spaces and let  $L^2(\Omega; \mathbb{Y}), L^2(\Omega; \mathbb{X})$  and  $L^2(\Omega; \mathbb{Z})$  be their corresponding  $L^2$ -spaces, respectively.

**Definition 2.1.** [5] A stochastic process  $y : \mathbb{R} \rightarrow L^2(\Omega; \mathbb{Y})$  is said to be continuous whenever

$$\lim_{t \rightarrow s} E\|y(t) - y(s)\|_{\mathbb{Y}}^2 = 0.$$

**Definition 2.2.** [5] A continuous stochastic process  $y : \mathbb{R} \rightarrow L^2(\Omega; \mathbb{Y})$  is said to be quadratic mean almost periodic if for each  $\epsilon > 0$  there exists  $l(\epsilon) > 0$  such that any interval of length  $l(\epsilon)$  contains at least a number  $\tau$  for which

$$\sup_{t \in \mathbb{R}} E \|y(t + \tau) - y(t)\|_{\mathbb{Y}}^2 < \epsilon.$$

The collection of all stochastic processes  $y : \mathbb{R} \rightarrow L^2(\Omega; \mathbb{Y})$  which are quadratic mean almost periodic is then denoted by  $MP(\mathbb{R}; L^2(\Omega; \mathbb{Y}))$ .

**Proposition 2.2.** If  $y$  belongs to  $MP(\mathbb{R}; L^2(\Omega; \mathbb{X}))$ , then the following hold true:

- (i) the mapping  $t \rightarrow E \|y(t)\|_{\mathbb{Y}}^2$  is uniformly continuous,
- (ii) there exists a constant  $N > 0$ , such that  $E \|y(t)\|_{\mathbb{Y}}^2 \leq N$ , for each  $t \in \mathbb{R}$ ,
- (iii)  $y$  is stochastically bounded.

Let  $C(\mathbb{R}; L^2(\Omega; \mathbb{X}))$  denote the space of all continuous stochastic processes  $y : \mathbb{R} \rightarrow L^2(\Omega; \mathbb{X})$ . The notation  $CUB(\mathbb{R}; L^2(\Omega; \mathbb{Y}))$  stands for the collection of all stochastic processes  $x : \mathbb{R} \rightarrow L^2(\Omega; \mathbb{Y})$ , which are continuous and uniformly bounded. we know from [2] that  $CUB(\mathbb{R}; L^2(\Omega; \mathbb{X}))$  is a Banach space endowed with the norm:

$$\|y\|_{\infty} = \sup_{t \in \mathbb{R}} (E \|y(t)\|_{\mathbb{Y}}^2)^{\frac{1}{2}}.$$

**Proposition 2.3.**  $MP(\mathbb{R}; L^2(\Omega; \mathbb{Y})) \subset CUB(\mathbb{R}; L^2(\Omega; \mathbb{Y}))$  is a closed subspace.

**Proposition 2.4.**  $(MP(\mathbb{R}; L^2(\Omega; \mathbb{Y})), \|\cdot\|_{MP(\mathbb{R}; L^2(\Omega; \mathbb{Y}))})$  is a Banach space endowed with the norm:

$$\|y\|_{AP(\mathbb{R}; L^2(\Omega; \mathbb{Y}))} = \sup_{t \in \mathbb{R}} (E \|y(t)\|_{\mathbb{Y}}^2)^{\frac{1}{2}}.$$

**Definition 2.3.** [2] The function  $N : \mathbb{R} \times L^2(\Omega; \mathbb{X}) \rightarrow L^2(\Omega; \mathbb{Z})$ ,  $(t, x) \rightarrow F(t, x)$ , which is jointly continuous, is said to be quadratic mean almost periodic in  $t \in \mathbb{R}$  uniformly in  $x \in \mathbb{B}$  where  $\mathbb{B} \subset L^2(\Omega; \mathbb{X})$  is compact if for any  $\epsilon > 0$ , there exists  $l(\epsilon, \mathbb{B}) > 0$  such that any interval of length  $l(\epsilon, \mathbb{B})$  contains at least a number  $\tau$  for which

$$\sup_{t \in \mathbb{R}} (E \|N(t + \tau, y) - N(t, y)\|_{\mathbb{Z}}^2) < \epsilon.$$

for each stochastic process  $x : \mathbb{R} \rightarrow \mathbb{B}$ .

**Proposition 2.5.** Let  $N : \mathbb{R} \times L^2(\Omega; \mathbb{X}) \rightarrow L^2(\Omega; \mathbb{Z})$ ,  $(t, x) \rightarrow N(t, x)$ , be a quadratic mean almost periodic process in  $t \in \mathbb{R}$  uniformly in  $x \in \mathbb{B}$ , where  $\mathbb{B} \subset L^2(\Omega; \mathbb{X})$  is compact. Suppose that  $N$  is Lipschitz in the following sense:

$$E \|N(t, y) - N(t, x)\|_{\mathbb{Z}}^2 \leq \tilde{M} E \|y - x\|_{\mathbb{X}}^2$$

for all  $y, x \in L^2(\Omega; \mathbb{X})$  and for each  $t \in \mathbb{R}$ , where  $\tilde{A} > 0$ . Then for any quadratic mean almost periodic process  $\Psi : \mathbb{R} \rightarrow L^2(\Omega; \mathbb{X})$ , the stochastic process  $t \rightarrow F(t, \Psi(t))$  is quadratic mean almost periodic.

**Definition 2.4.** A  $N_t$ -progressively process  $\{y(t)\}_{t \in \mathbb{R}}$  is called a mild solution of

the problem (1.1) on  $\mathbb{R}$  if the function  $s \rightarrow MT(t-s)h(s, y(s))$  is integrable on  $(-\infty, t)$  for each  $t \in \mathbb{R}$ , and  $y(t)$  satisfies

$$y(t) = T(t-a)[y(a) - h(a, y(a))] + h(t, y(t)) \\ + \int_a^t MT(t-s)h(s, y(s))ds + \int_a^t T(t-s)S(s, y(s))d\omega(s)$$

for all  $t \geq a$  and for each  $a \in \mathbb{R}$ .

Let us list the following assumptions:

(H1) The operator  $M : D(M) \subset L^2(\Omega; \mathbb{H}) \rightarrow L^2(\Omega; \mathbb{H})$  is the infinitesimal generator of an analytic semigroup of linear operators  $\{T(t)\}_{t \geq 0}$  on  $L^2(\Omega; \mathbb{H})$  and  $A, \delta$  are positive numbers such that  $\|T(t)\| \leq Me^{-\delta t}$  for  $t \geq 0$ .

(H2) There exists a positive number  $\alpha \in (0, 1)$  such that  $g : \mathbb{R} \times L^2(\Omega; \mathbb{H}) \rightarrow L^2(\Omega; \mathbb{H}_\alpha)$  is quadratic mean almost periodic in  $t \in \mathbb{R}$  uniformly in  $y \in \mathbb{B}_1$  where  $\mathbb{B}_1 \subset L^2(\Omega; \mathbb{H})$  being a compact subspace. also,  $g$  is Lipschitz in the sense that: there exists  $L_g > 0$  such that

$$E\|(-M)^\alpha h(t, y) - (-M)^\alpha h(t, x)\|^2 \leq L_h E\|y - x\|^2,$$

for all  $t \in \mathbb{R}$  and for each stochastic processes  $x, y \in L^2(\Omega; \mathbb{H})$ .

(H3) The function  $G : \mathbb{R} \times L^2(\Omega; \mathbb{H}) \rightarrow L^2(\Omega; L^0_2)$  is quadratic mean almost periodic in  $t \in \mathbb{R}$  uniformly in  $y \in \mathbb{B}_2$  where  $\mathbb{B}_2 \subset L^2(\Omega; \mathbb{H})$  being a compact subspace. Moreover,  $S$  is Lipschitz in the sense that: there exists  $L_S > 0$  such that

$$E\|S(t, x) - S(t, y)\|_{L^0_2}^2 \leq L_S E\|y - x\|^2$$

for all  $t \in \mathbb{R}$  and for each stochastic processes  $x, y \in L^2(\Omega; \mathbb{H})$ .

### 3. Main Results

In this Section, we Prove our Main Theorem.

**Theorem 3.1.** suppose the conditions (H1)-(H3) are satisfied, then the problem (1.1) admits a unique quadratic mean almost periodic mild solution on  $\mathbb{R}$  provide that

$$L_0 = [3L_g\|(-M)^{-\alpha}\|^2 + 3A_{1-\alpha}^2 L_h \delta^{-2\alpha} [\Gamma(\alpha)]^2 + \frac{3T_r Q M^2 L_S}{2\delta}] < 1,$$

where  $\Gamma(\cdot)$  is the gamma function. *Proof.* Let  $\Lambda : AP(\mathbb{R}; L^2(\Omega; \mathbb{H})) \rightarrow C(\mathbb{R}; L^2(\Omega; \mathbb{H}))$  be the operator defined by

$$\Lambda y(t) = h(t, y(t)) + \int_{-\infty}^t MT(t-s)h(s, y(s))ds \\ + \int_{-\infty}^t T(t-s)S(s, y(s))d\omega(s), t \in \mathbb{R}.$$

First It proves that  $\Lambda y$  is well defined. From Proposition 2.5, we infer that  $s \rightarrow h(s, y(s))$  is in  $MP(\mathbb{R}; L^2(\Omega; \mathbb{H}))$ . therefore using Proposition 2.2 (ii) it follows that

there exists a constant  $N_g > 0$  such that  $E\|(-M)^\alpha h(t, y(t))\|^2 N_g$ , for all  $t \in \mathbb{R}$ . Moreover, from the continuity of  $s \rightarrow MT(t-s)$  and  $s \rightarrow T(t-s)$  in the uniform operator topology on  $(-\infty, t)$  for each  $t \in \mathbb{R}$  and the estimate

$$\begin{aligned} & E\left\|\int_{-\infty}^t MT(t-s)h(s, y(s))ds\right\|^2 \\ &= E\left\|\int_{-\infty}^t (-M)^{1-\alpha}T(t-s)(-M)^\alpha h(s, y(s))ds\right\|^2 \\ &\leq A_{1-\alpha}^2 E\left(\int_{-\infty}^t e^{-\delta(t-s)}(t-s)^{\alpha-1}\|(-M)^\alpha h(s, y(s))\|ds\right)^2 \\ &\leq A_{1-\alpha}^2 \left(\int_{-\infty}^t e^{-\delta(t-s)}(t-s)^{\alpha-1}ds\right)^2 \\ &\times \left(\int_{-\infty}^t e^{-\delta(t-s)}(t-s)^{\alpha-1}E\|(-M)^\alpha h(s, y(s))\|^2 ds\right) \\ &\leq F_h M_{1-\alpha}^2 \left(\int_{-\infty}^t e^{-\delta(t-s)}(t-s)^{\alpha-1}ds\right)^2 = F_h A_{1-\alpha}^2 \delta^{-2\alpha} [\Gamma(\alpha)]^2, \end{aligned}$$

Hence  $s \rightarrow MT(t-s)h(s, y(s))$  and  $s \rightarrow T(t-s)S(s, y(s))$  are integrable on  $(-\infty, t)$  for every  $t \in \mathbb{R}$ , then  $\Lambda y$  is well defined and continuous.

Next, we show that  $\Lambda y(t) \in MP(\mathbb{R}; L^2(\Omega; \mathbb{H}))$ . we can define

$$\Lambda_1 y(t) = \int_{-\infty}^t MT(t-s)h(s, y(s))ds$$

and

$$\Lambda_2 y(t) = \int_{-\infty}^t T(t-s)S(s, y(s))d\omega(s)$$

we show that  $\Lambda_1 y(t)$  is quadratic mean almost periodic. since  $h(\cdot, y(\cdot)) \in MP(\mathbb{R}; L^2(\Omega; \mathbb{H}_\alpha))$ , by Definition 2.2, it follows that for any  $\epsilon > 0$ , there exists  $l(\epsilon) > 0$  such that every interval of length  $l(\epsilon)$  contains at least a number  $\tau$  with the property that

$$E\|(-M)^\alpha h(t+\tau, y(t+\tau)) - (-M)^\alpha h(t, y(t))\|^2 < \frac{\epsilon}{A_{1-\alpha}^2 \delta^{-2\alpha} [\Gamma(\alpha)]^2},$$

for each  $t \in \mathbb{R}$ .

using Cauchy-Schwarz inequality, we get

$$\begin{aligned}
 & E\|\Lambda_1 y(t + \tau) - \Lambda_1 y(t)\|^2 \\
 & E\left\| \int_{-\infty}^t MT(t-s)[h(s + \tau, y(s + \tau)) - h(s, y(s))]ds \right\|^2 \\
 & = E\left\| \int_{-\infty}^t (-M)^{1-\alpha} T(t-s)[(-M)^\alpha h(t + \tau, y(t + \tau)) - (-M)^\alpha h(t, y(t))]ds \right\|^2 \\
 & \leq A_{1-\alpha}^2 E\left( \int_{-\infty}^t e^{-\delta(t-s)}(t-s)^{\alpha-1} \times \|(-M)^\alpha h(s + \tau, x(s + \tau)) - (-M)^\alpha h(s, y(s))\| ds \right)^2 \\
 & \leq A_{1-\alpha}^2 E\left[ \left( \int_{-\infty}^t e^{-\delta(t-s)}(t-s)^{\alpha-1} ds \right) \right. \\
 & \quad \times \left. \left( \int_{-\infty}^t e^{-\delta(t-s)}(t-s)^{\alpha-1} \|(-M)^\alpha h(s + \tau, y(s + \tau)) - (-M)^\alpha h(s, y(s))\|^2 ds \right) \right] \\
 & \leq A_{1-\alpha}^2 \left( \int_{-\infty}^t e^{-\delta(t-s)}(t-s)^{\alpha-1} \right) \\
 & \quad \times \left( \int_{-\infty}^t e^{-\delta(t-s)}(t-s)^{\alpha-1} E\|(-M)^\alpha h(s + \tau, y(s + \tau)) - (-M)^\alpha h(s, y(s))\|^2 ds \right) \\
 & \leq A_{1-\alpha}^2 \left( \int_{-\infty}^t e^{-\delta(t-s)}(t-s)^{\alpha-1} ds \right)^2 \times \sup_{t \in \mathbb{R}} E\|(-M)^\alpha h(t + \tau, y(t + \tau)) - (-M)^\alpha h(t, y(t))\|^2 \\
 & \leq \frac{\epsilon}{\delta^{-2\alpha}[\Gamma(\alpha)]^2} \left( \int_{-\infty}^t e^{-\delta(t-s)}(t-s)^{\alpha-1} ds \right)^2 = \epsilon
 \end{aligned}$$

Therefore,  $\Lambda_1 y(\cdot)$  is quadratic mean almost periodic.

Similarly, by using Proposition 2.5, hence  $s \rightarrow S(s, y(s))$  is quadratic mean almost periodic. Therefore, it follows from Definition 2.2 that for any  $\epsilon > 0$ , there exists  $l(\epsilon) > 0$  such that every interval of length  $l(\epsilon)$  contains at least a number  $\tau$  with the property that

$$E\|S(t + \tau, y(t + \tau)) - S(t, y(t))\|_{L_2^0}^2 < \frac{2\delta\epsilon}{TrQM^2},$$

for each  $t \in \mathbb{R}$ . Now, let us prove that  $\Lambda_2 y(t)$  is quadratic mean almost periodic. We adopt the techniques developed in [2]. Let  $\tilde{\omega}(t) := \omega(t + \tau) - \omega(\tau)$  for each  $t \in \mathbb{R}$ , note that  $\tilde{\omega}$  is also a Brownian motion and has the same distribution as  $\omega$ .

Now, we consider

$$\begin{aligned}
 & E\|\Lambda_2 y(t + \tau) - \Lambda_2 y(t)\|^2 \\
 & = E\left\| \int_{-\infty}^{t+\tau} T(t + \tau - s)S(s, y(s))d\omega(s) - \int_{-\infty}^t T(t - s)S(s, y(s))d\omega(s) \right\|^2 \\
 & = E\left\| \int_{-\infty}^t T(t - s)[S(s + \tau, y(s + \tau)) - S(s, y(s))]d\tilde{\omega}(s) \right\|^2
 \end{aligned}$$

Thus using estimate on Ito integral established in Ichikawa [11], we obtain that

$$\begin{aligned}
& E\|\Lambda_2 y(t + \tau) - \Lambda_2 y(t)\|^2 \\
&= E\left\| \int_{-\infty}^t T(t-s)[S(s + \tau, y(s + \tau)) - S(s, y(s))]d\tilde{\omega}(s) \right\|^2 \\
&\leq TrQE\left[ \int_{-\infty}^t \|T(t-s)[S(s + \tau, y(s + \tau)) - S(s, y(s))]\|^2 ds \right] \\
&\leq TrQE\left[ \int_{-\infty}^t \|T(t-s)\|^2 \|S(s + \tau, y(s + \tau)) - S(s, y(s))\|_{L_2^0}^2 ds \right] \\
&\leq TrQM^2 \int_{-\infty}^t e^{-2\delta(t-s)} E\|S(s + \tau, y(s + \tau)) - S(s, y(s))\|_{L_2^0}^2 ds \\
&\leq TrQM^2 \left( \int_{-\infty}^t e^{-2\delta(t-s)} ds \right) \sup_{t \in \mathbb{R}} E\|S(t + \tau, y(t + \tau)) - S(t, y(t))\|_{L_2^0}^2 \\
&< 2\delta\epsilon \int_{-\infty}^t e^{-2\delta(t-s)} ds = \epsilon.
\end{aligned}$$

Thus,  $\Lambda_2 y(\cdot)$  is quadratic mean almost periodic. And in view of the above, it is clear that  $\Lambda$  maps  $AP(\mathbb{R}; L^2(\Omega; \mathbb{H}))$  into itself.

Now we should prove that  $\Lambda$  is a strict contraction on  $AP(\mathbb{R}; L^2(\Omega; \mathbb{H}))$ . Indeed, for each  $t \in \mathbb{R}, x, y \in AP(\mathbb{R}; L^2(\Omega; \mathbb{H}))$ , we have

$$\begin{aligned}
& E\|\Lambda y(t) - \Lambda x(t)\|^2 \leq 3E\|h(t, y(t)) - h(t, x(t))\|_\alpha^2 \\
&+ 3E\left(\left\| \int_{-\infty}^t AT(t-s)[h(s, y(s)) - h(s, x(s))]ds \right\|\right)^2 \\
&+ 3E\left(\left\| \int_{-\infty}^t T(t-s)[S(s, y(s)) - S(s, x(s))]d\omega(s) \right\|\right)^2 \\
&\leq 3\|(-M)^{-\alpha}\|^2 E\|(-M)^\alpha h(t, y(t)) - (-M)^\alpha h(t, x(t))\|^2 \\
&+ 3E\left(\left\| \int_{-\infty}^t (-M)^{1-\alpha} T(t-s)[(-M)^\alpha h(s, y(s)) - (-M)^\alpha h(s, x(s))]ds \right\|\right)^2 \\
&+ 3TrQE\left(\int_{-\infty}^t \|(T(t-s)[S(s, y(s)) - S(s, x(s))]\|^2 ds\right) \\
&\leq 3\|(-M)^{-\alpha}\|^2 \sup_{t \in \mathbb{R}} E\|(-M)^\alpha h(t, y(t)) - (-M)^\alpha h(t, x(t))\|^2 \\
&+ 3A_{1-\alpha}^2 E\left(\int_{-\infty}^t e^{-\delta(t-s)}(t-s)^{\alpha-1} \|(-M)^\alpha h(s, y(s)) - (-M)^\alpha h(s, x(s))\| ds\right)^2 \\
&+ 3TrQE\left(\int_{-\infty}^t \|T(t-s)\|^2 \|S(s, y(s)) - S(s, x(s))\|_{L_2^0}^2 ds\right) \\
&\leq 3L_h\|(-M)^{-\alpha}\|^2 \sup_{t \in \mathbb{R}} E\|y(t) - x(t)\|^2 + 3A_{1-\alpha}^2 E\left[\left(\int_{-\infty}^t e^{-\delta(t-s)}(t-s)^{\alpha-1} ds\right) \right. \\
&\times \left. \left(\int_{-\infty}^t e^{-\delta(t-s)}(t-s)^{\alpha-1} \|(-M)^\alpha h(s, y(s)) - (-M)^\alpha h(s, x(s))\|^2 ds\right) \right]
\end{aligned}$$



$$\begin{aligned}
& + 3TrQA^2 \int_{-\infty}^t e^{-2\delta(t-s)} E \| [S(s, y(s)) - S(s, x(s))] \|_{L_2^0}^2 ds \\
& \leq 3L_h \| (-M)^{-\alpha} \|^2 \sup_{t \in \mathbb{R}} E \| y(t) - x(t) \|^2 + 3A_{1-\alpha}^2 E \left( \int_{-\infty}^t e^{-\delta(t-s)} (t-s)^{\alpha-1} ds \right) \\
& \times \left( \int_{-\infty}^t e^{-\delta(t-s)} (t-s)^{\alpha-1} E \| (-A)^\alpha g(s, x(s)) - (-A)^\alpha g(s, y(s)) \|^2 \right) \\
& + 3TrQA^2 L_h \int_{-\infty}^t e^{-2\delta(t-s)} ds \sup_{t \in \mathbb{R}} E \| y(t) - x(t) \|^2 \\
& \leq 3L_h \| (-M)^{-\alpha} \|^2 \sup_{t \in \mathbb{R}} E \| y(t) - x(t) \|^2 + 3A_{1-\alpha}^2 \left( \int_{-\infty}^t e^{-\delta(t-s)} (t-s)^{\alpha-1} ds \right)^2 \sup_{t \in \mathbb{R}} E \| y(t) - x(t) \|^2 \\
& + 3TrQA^2 L_h \frac{1}{2\delta} \sup_{t \in \mathbb{R}} E \| y(t) - x(t) \|^2 \\
& \leq 3L_h \| (-M)^{-\alpha} \|^2 \sup_{t \in \mathbb{R}} E \| y(t) - x(t) \|^2 \\
& + 3A_{1-\alpha}^2 L_h \delta^{-2\alpha} [\Gamma(\alpha)]^2 \times \sup_{t \in \mathbb{R}} E \| y(t) - x(t) \|^2 + \frac{3TrQM^2 L_h}{2\delta} \sup_{t \in \mathbb{R}} E \| y(t) - x(t) \|^2 \\
& = [3L_h \| (-M)^{-\alpha} \|^2 + 3A_{1-\alpha}^2 L_h \delta^{-2\alpha} [\Gamma(\alpha)]^2 + \frac{3TrQM^2 L_h}{2\delta}] \times \sup_{t \in \mathbb{R}} E \| y(t) - x(t) \|^2
\end{aligned}$$

by using the arithmetic geometric inequality, Cauchy-Schwarz inequality and Ito isometry identity.

Note that

$$\sup_{t \in \mathbb{R}} E \| y(t) - x(t) \|^2 \leq [\sup_{t \in \mathbb{R}} (E \| y(t) - x(t) \|^2)^{\frac{1}{2}}]^2$$

Thus, it follows that, for each  $t \in \mathbb{R}$ ,

$$(E \| \Lambda y(t) - \Lambda x(t) \|^2)^{\frac{1}{2}} \leq \sqrt{L_0} \| y - x \|_{MP(\mathbb{R}; L^2(\Omega; \mathbb{H}))}.$$

Hence

$$\| \Lambda y - \Lambda x \|_{MP(\mathbb{R}; L^2(\Omega; \mathbb{H}))} = \sup_{t \in \mathbb{R}} (E \| y(t) - x(t) \|^2)^{\frac{1}{2}} \leq \sqrt{L_0} \| y - x \|_{MP(\mathbb{R}; L^2(\Omega; \mathbb{H}))},$$

which implies that  $\Lambda$  is a contraction by (3.1).

#### 4. Conclusion

So by the contraction principle, we conclude that there exists a unique fixed point  $y(\cdot)$  for  $\Lambda$  in  $MP(\mathbb{R}; L^2(\Omega; \mathbb{H}))$ , such that  $\Lambda y = y$ , that is

$$y(t) = h(t, y(t)) + \int_{-\infty}^t MT(t-s)h(s, x(s))ds + \int_{-\infty}^t T(t-s)S(s, y(s))d\omega(s)$$

for all  $t \in \mathbb{R}$ . If we let  $y(a) = h(a, x(a)) + \int_{-\infty}^t MT(a-s)h(s, y(s))ds + \int_{-\infty}^t T(a-s)S(s, y(s))d\omega(s)$ , then

$$T(t-a)y(a) = T(t-a)h(a, y(a)) + \int_{-\infty}^t MT(t-s)h(s, y(s))ds + \int_{-\infty}^t T(t-s)S(s, y(s))d\omega(s).$$

But for  $t \geq a$ ,

$$\begin{aligned} & \int_a^t T(t-s)S(s, y(s))d\omega(s) \\ &= \int_{-\infty}^t T(t-s)S(s, y(s))d\omega(s) - \int_{-\infty}^a T(t-s)S(s, y(s))d\omega(s) \\ &= y(t) - h(t, y(t)) - \int_{-\infty}^t MT(t-s)h(s, y(s))ds - T(t-a)[y(a) - h(a, y(a))] \\ &+ \int_{-\infty}^a MT(t-s)h(s, y(s))ds = y(t) - h(t, y(t)) \\ &- \int_a^t MT(t-s)h(s, y(s))ds - T(t-a)[y(a) - h(a, y(a))]. \end{aligned}$$

In conclusion,

$y(t) = T(t-a)[y(a) - h(a, y(a))] + h(t, y(t)) + \int_a^t MT(t-s)h(s, y(s))ds + \int_a^t T(t-s)S(s, y(s))d\omega(s)$  is a mild solution of the problem (1.1) and  $y(\cdot) \in MP(\mathbb{R}; L^2(\Omega; \mathbb{H}))$ .

a simple example to illustrate main theorem. I examine the existence and uniqueness of quadratic mean almost periodic solutions for the following stochastic partial differential equation

$$\frac{\partial}{\partial t}[y(t, \xi) - h(t, y(t, \xi))] = \frac{\partial^2}{\partial \xi^2}y(t, \xi) + S(t, y(t, \xi))d\omega(t), t \in \mathbb{R}, \xi \in \mathcal{D}, \quad (2)$$

where  $\mathcal{D} \subset \mathbb{R}^n (n \geq 1)$  is a bounded subset with  $C^2$  boundary  $\partial\mathcal{D}$ .

Let  $H := L^2(\mathcal{D})$  be equipped with its natural topology and define an operator  $A$  on  $L^2(\mathbb{R}; H)$  by

$$Ax(t, \cdot) = \frac{\partial^2}{\partial \xi^2}x(t, \cdot), x \in H^2(\mathcal{D} \cap H_0^1(\mathcal{D})).$$

It is well known that (see [15-18])  $A$  is the infinitesimal generator of an analytic semigroup  $\{T(t)\}_{t \geq 0}$  on  $L^2(\mathbb{R}; H)$  satisfying (H1). Therefore, under assumptions (H2)-(H3), if we assume that (3.1) holds, by Theorem 3.1 we can say (4.1) has a unique quadratic mean almost periodic mild solution.

## References

- [1] Acquistapace. P and Terreni. B., A unified approach to abstract linear nonautonomous parabolic equations. Rend. Sem. Mat. Univ. Padova, **78** (1987) 47-107.
- [2] Bezandry. P and Diagana. T., Existence of almost periodic solutions to some stochastic differential equations. Appl. Anal, **86** (2007) 819-827.
- [3] Bezandr. P and Diagana. T., Square-mean almost periodic solutions nonautonomous stochastic differential equations. Electron. J. Differ. Equ, (2007) 1-10.

- [4] Bezandry. P., Existence of almost periodic solutions to some functional integro-differential stochastic evolution equations. *Stat. Probab. Lett.*, **78** (2008) 2844-2849.
- [5] Cao. J., Yang. Q., Huang. Z and Liu. Q., Asymptotically almost periodic solutions of stochastic functional differential equations. *Appl. Math. Comput.*, **218** (2011) 1499-1511.
- [6] Corduneanu. C., *Almost Periodic Functions*, 2nd edn. Chelsea, New York, (1989).
- [7] Da Prato. G and Zabczyk. J., *Stochastic Equations in Infinite Dimensions*. Cambridge University Press, Cambridge, (1992).
- [8] Dorogovtsev. A. Ya and Ortega. O. A., On the existence of periodic solutions of a stochastic equation in a Hilbert space. *Visnik Kiiv. Univ. Ser. Mat. Mekh.*, **115** (1988) 21-30.
- [9] Hernandez. E. M., Pelicer, M. L and dos Santos. J. P. C., Asymptotically almost periodic and almost periodic solutions for a class of evolution equations. *Electron. J. Differ. Equ.*, (2004) 1-15.
- [10] Hernandez. E and Pelicer, H. L., Asymptotically almost periodic and almost periodic solutions for partial neutral differential equations. *Appl. Math. Lett.*, **18** (2005) 1265-1272.
- [11] Lunardi. A., *Analytic Semigroups and Optimal Regularity in Parabolic Problems*, PNLDE, Birkhauser, Basel, **16** (1995).
- [12] F. Mirzaee and A. Hamzeh., A Sixth Order Method for Solving Nonlinear Equations, *ijm2c*, **4** (1) (2014).
- [13] N'Guerekata. G. M., *Almost Automorphic Functions and Almost Periodic Functions in Abstract Spaces*. Kluwer Academic Plenum Publishers, New York, (2001).
- [14] Pazy. A., *Semigroups of Linear Operators and Applications to Partial Equations*, in: *Applied Mathematical Sciences*, Springer, New York, **44** (1983).
- [15] P. K. Pandey., Non-standard Finite Difference Method for Numerical Solution of Second Order Linear Fredholm Integro-Differential Equations, *ijm2c*, **5** (3) (2015).
- [16] Tudor. C., Almost periodic solutions of affine stochastic evolutions equations. *Stoch. Rep.*, **38** (1992) 251-266.
- [17] Xie. B., Stochastic differential equations with non-Lipschitz coefficients in Hilbert spaces. *Stoch. Anal. Appl.*, **26** (2008) 408-433.
- [18] Zhao. Z. H., Chang. Y. K and Li. W. S., Asymptotically almost periodic, almost periodic and pseudo almost periodic mild solutions for neutral differential equations. *Nonlinear Anal. RWA*, **11** (2010) 3037-3044.