

Effects of synthetic pyrethroids and neonicotinoids insecticides and synergists on population of *Spodoptera littoralis* (Lepidoptera: Noctuidae)

A. Dini pour^{*1}, M. Oktay Gurkan²

1- Ph.D. Student of Agricultural Entomology, Agricultural Faculty, Ankara University, Ankara, Turkey

2- Plant Protection Department, Agricultural Faculty, Ankara University, Ankara, Turkey

Abstract

Spodoptera littoralis (Lepidoptera: Noctuidae) known as the African Cotton Leaf worm or Egyptian Cotton Leaf worm is one of the most common pests of cotton in the region of Adana, Turkey. The aim of this study was to determine the effects of a synthetic pyrethroid (Gamma-cyhalothrin) and neonicotinoids (Imidacloprid) on the pest to find out the resistance inducing process during several generations. Induced resistance to gamma-cyhalothrin up to 12 generations and to Imidacloprid for 7 generations was studied by assessing the resistance ratio (RR) of LC₅₀ and LC₉₀ with confidence limit of %95. The resistance ratios (RR) for LC₅₀ and LC₉₀ of gamma-cyhalothrin were calculated 16.85 and 18.82 and for Imidacloprid were 9.76 and 8.68, respectively. In order to study the effect of synergists on the insecticides, three different synergists including S, S, S-tributylfosforotritioat (DEF), piperonyl butoxide (PBO) and diethyl maleate (DEM) were used and compared the Synergism Ratio (SR) with control. The highest SR for both insecticides belonged to application of DEM synergist. It is concluded that DEM could be recommended to be applied with Gamma-cyhalothrin and Imidacloprid to prevent or postpone the resistance induction into *S. littoralis*.

Key words : *Spodoptera littoralis*, bioassay, resistance, insecticide, synergist

* Corresponding Author, E-mail: dinipourm@gmail.com

Received: 11 May 2012 -Accepted: 20 Dec. 2012

Introduction

Spodoptera littoralis (Lepidoptera: Noctuidae) (Boisduval) attacks to a wide range host of crops (at least 87 different hosts) in tropical and subtropical regions. It causes economic damages on plants throughout the year. In cotton plant it induces serious damages to leaves, seeds, buds, and bolls (Salama *et al* .1970).

Despite their efficiencies and being a definitive solution for pest control, pesticides, especially insecticides have endangered human health or have had negative impacts on non-target organisms. They have hazarded the environment, especially when improperly used by farmers during the past fifty years (Topuz, 2005). Moreover they have induced resistance in target organisms. Besides, overusing the insecticides against pests with several generations a year make them highly resistant in consequence of natural selection (Tiryaki *et al* .,2010 .)

Extensive investigations on some chemicals have been conducted to inhibit or at least lower the insecticides detoxification by insects. S, S, S-tributylfosforotritioat (DEF) as an inhibitor of esterase enzymes, piperonyl butoxide (PBO) as an inhibitor of Cytochrome P-450 monooxygenase enzyme, and Diethyl maleate (DEM) (as an inhibitor of Glutation S-transferase enzyme have been used with chemical to enhance the efficiency of insecticides by inhibiting their insecticides detoxification (Van Leeuwen *et al*) .2004. According to the published reports, changes in content of Cytochrome P-450 monooxygenase enzyme and esterase in the genus *Spodoptera* usually cause an increase the metabolic activities and resistance to insecticide (Armes *et al* .,1997 ;Zhou and Huang, 2002; Huang and Han, 2007; Ahmad, 2008).

Gamma-cyhalothrin is a contact insecticide which can also be effective through ingestion. It acts as a sodium channel modulator and causes neurotic disorder (Metcalf, 2000). Imidacloprid belongs to neo-nicotinoid insecticides class and is known as an agonist of nicotinic acetylcholine receptor (nAChR). Acetylcholine is an endogenous agonist as well as a cholinergic signaling of the nervous system which is known as a neurotransmitter (Pittendrigh *et al* .,2008.)

Imidacloprid is a systemic insecticide which can also be effective through ingestion or contact. It penetrates swiftly into the plant tissues after applying on plants. It has a variety of functions on the nervous system and especially on post-synaptic nicotinic acetylcholine receptor. In class of insects, these receptors are found solely on the central nervous system (CNS). Imidacloprid molecules after attachment to the nicotinic receptor, firstly neurotic signal discharging happens, and then, by any neurotic signal, nerve failure occurs. Reason of receptors non-stop activity is the adhering the insecticide molecules to acetylcholinesterase enzyme irreversibly (Gervais, et al., 2010).

Comparison of resistant strains of *S. litura* to organophosphates and pyrethroids collected from five different districts of China with a susceptible strain reared in laboratory, the values of LC_{50} increased 14-229-fold for organophosphates and 12-227-fold for pyrethroids respectively (Tong *et al.*, 2013). Application of DDT on *S. littoralis* resulted 43 times more resistant individuals after seventeenth generations; whereas, simultaneous use it with Du-ter (a type of synergists) has induced 11.5 times resistance (Abdallah *et al.*, 1979). Ahmed and his colleague (2011) using endosulfan on *S. litura* found that the resistance ratio in the 6th generation increased 3.97 fold compare to the 1st generation. Efficiency of pyrethroids and organophosphates on populations of *Spodoptera* significantly increased by using DEF and PBO enzyme inhibitors. It was shown that the resistance to insecticides was associated with esterase and monooxygenase detoxification respectively (Ahmad *et al.*, 2008). Carboxylesterase is one of the most commonly used enzyme which associated with insecticide resistance. Metabolic resistance to organophosphates increases the esterase activity in insects resistant to carbamates. Some evidence suggest that cytochrome P-450 monooxygenase enzyme involves in synthesis of resistance to carbamates in cases of frequent use of insecticides (Farnsworth *et al.*, 2010).

Until now, investigations have indicated that improper use of insecticides with various modes of actions or broad spectrum insecticides cause resistance or reduce susceptibility to a special class of insecticide or cross resistance. The results of these investigations show that esterase, Glutathion S-transferase, Cytochrome P-450 monooxygenase induce resistance by increasing metabolism or reducing penetration to cuticle (Riskalla, 1983; Ishaq, 1990).

This study aims to find out the effects of two insecticides, Gamma-cyhalothrin (VANTEX) (and Imidacloprid (Confidor) on *S. littoralis* using bioassay under laboratory conditions.

Materials and Methods

S. littoralis (Boisd.) was collected from cotton fields of Adana region, Turkey, in 2008, and since then reared in conditions of $25 \pm 2^{\circ}C$, 65 ± 5 relative humidity and 16:8 photoperiod (light: dark) at the Department of Plant Protection, Faculty of Agriculture, University of Ankara. Third instars larvae (6 days after hatching) were used in all experiments.

Synthetic pyrethroid with the active ingredient of gamma-cyhalothrin (VANTEX, 60 SC, Dow AgroSciences) and Neo-nicotinoid with the active ingredient of imidacloprid (CONFIDOR: 350 SC: Bayer CropScience) were used in this study. We also used the S, S, S-tributylfosforotritioat (DEF) as an inhibitor of esterase enzymes, piperonyl butoxide (PBO) as an inhibitor of the of cytochrome P-450 monooxygenase and carboxylesterase enzymes, and diethyl maleate (DEM) as an inhibitor of Glutathion S-transferase enzyme diluted in acetone (1000 g / ml solution (Van Leeuwen *et al.*, 2004).

Bioassays and selections

A susceptible strain of *S. littoralis* (Boisd.) was collected in 2008 from cotton fields of Adana province in Turkey and reared until now in laboratory without exposure to insecticides. Bioassays were conducted in growth chamber and mortality data were used for susceptibility to insecticides. The newly third-instar larvae of *S. littoralis* (Boisd.) used in bioassay experiments. Serial dilutions of active ingredient were done using distilled water. Lettuce leaves were cut into 9 cm pieces and dipped into the insecticide solution for 10s. These leaves were air dried on tissue paper, under lumina hood. The leaves were dipped in distilled water to use as controls. Leaves treated with insecticides were then transferred to each petri dish lined with moistened filter paper. At least six concentrations and three replications (12 larvae per replication) were used to estimate mortality compare to previous generation. Mortality data were scored 48 h after exposure to insecticides. Data were analyzed using probit analysis through POLO-PLUS software to estimate LC_{50} , LC_{75} and LC_{90} values and 95 % confidence limits. The LC_{50} values for each generation was calculated and used for selection the survived individuals for the next step. All these steps were repeated for 13 and 7 times for gamma-cyhalothrin and imidacloprid respectively.

Synergistic tests

Synergistic bioassay were performed on sensitive population (non-exposed to insecticides (as control and two selected populations of *S. littoralis*) Boisd. (which were resistant to Gamma-cyhalothrin and Imidacloprid. PBO, DEM and DEF were used as synergists to determine the type and amount of the insecticide detoxification. These substances were diluted using acetone solution, and brought to a concentration of 1,000 micrograms per milliliter (Van Leeuwen *et al* .2004). In each synergistic test, the third instar larvae exposed to different synergists by topical application to the prothorax at the concentration of 1000 $\mu\text{g/ml}$ before performing the bioassay tests. Acetone was applied on individuals instead the synergist in controls. After two hours, the larvae exposed to different doses of insecticide by the leaf dipping method. Finally, the mortality rate was analyzed using Polo-Plus software at 95 % confidence level.

Results

In bioassay tests, the value of LC_{50} in F0 for gamma-cyhalothrin was 8.4.1ppm. This value increased gently until the seventh generation) F₇. (However it increased sharply from F₈ to F₁₂ and reached to 9.13-fold of its initial value, i.e. 11194. Similarly, after a sudden rise from the eighth generation, the LC_{90} value reached to 15.7-fold compare to initial value, i.e. 24340 ppm in F₁₂. In Figure 1 shows LC_{50} and LC_{90} values of gamma-cyhalothrin on selected and sensitive populations of *S.lithuralis* larvae. Accordingly, the resistance ratio for LC_{50} and LC_{90} values reached to 16.85 and 18.82-fold in 13th generation, respectively. Consequently, reduced sensitivity or increasing resistance to this insecticide appeared gradually. The heterogeneity ratio (h) in population exposed to gamma-cyhalothrin reduced from 2.7 to 0.24 in 13th generation. This means that homogeneity to gamma-cyhalothrin increased gradually in selected populations (Table 1). The resistance ratios (RR) which is considered as a resistance factor was calculated using LC_{50} or LC_{90} values of resistance population divided by LC_{50} or LC_{90} values of sensitive population increased 16.85 and 18.82 fold compared to sensitive populations (table 1)

Table 1- The bioassay parameters of 13 generations of third instar larvae of *S. littoralis* exposed to Gamma-cyhalothrin

RR LC ₉₀	RR LC ₅₀	LC ₉₀ ppm (CL 0.95)	LC ₅₀ ppm (CL 0.95)	Slope ±SE	h	χ ² (df)	n	Sel.
1.00	1.00	1551.5 (1315.2 – 2010.7)	804.1 (441.1 – 734.6)	4.49 ±0.61	2.70	14.82 (21)	216	F0
1.10	0.86	1313.0 (1063.6 – 1917.7)	604.6 (441.1 – 734.6)	3.80 ±0.59	1.32	14.52 (11)	216	F1
1.01	1.01	1629.2 (1443.4 – 3415.4)	651.1 (486.6 -799.5)	2.71 ±0.45	1.07	12.95 (12)	216	F2
1.17	1.03	1510.6 (1059.0 – 4017.6)	517.8 (269.2 – 703.3)	2.75 ±0.47	2.25	27 (12)	216	F3
1.08	1.27	1083.9 (944.5 – 1312.6)	597.9 (515.8 – 675.3)	4.96 ±0.63	1.90	10.83 (12)	216	F4
2.80	2.40	2549.0 (1640.4 – 10571.0)	957.6 (685.5 – 1395.2)	3.01 ±0.47	2.70	32.48 (12)	216	F5
1.56	2.34	1830.0 (1581.1 – 2443.,1)	1264.4 (1096.3 – 1438.4)	7.98 ±0.98	1.43	29.27 (12)	216	F6
1.35	2.28	1501.7 (1397.9 – 1684.5)	1169.8 (1118.9 – 1229.2)	11.81 ±1.60	0.71	8.71 (13)	216	F7
4.22	5.58	4389.1 (3603.4 – 6115.6)	2410.9 (2179.8 – 2768.2)	4.925 ±0.68	0.44	7.49 (18)	216	F8
4.36	6.82	4826.6 (4038.1 – 7263.1)	2746.1 (2451.7 – 3037.9)	5.23 ±0.12	0.41	5.6 (12)	216	F9
11.75	10.10	15318.0 (11796.0 – 23892.0)	6278.1 (4789.7 – 8230.0)	3.24 ±1	0.26	2.41 (15)	216	F10
12.53	11.01	15598.0 (8534.9 – 19803.4)	6483.4 (5669.8 – 7673.5)	3.43 ±0.49	0.26	3.94 (15)	216	F11
18.82	16.85	24340.0 (16427.0 - 47492.0)	11194.0 (9341.2 – 16705.0)	3.79 ±0.88	0.24	3.31 (13)	216	F12

n =Number of larvae used

h =Heterogeneous value

RR =resistance ratio (LC₉₀. or LC₉₅. values of resistance population / LC₉₀. or LC₉₅. values of sensitive population)

CL=confidence limit




Fig. 1- The trend of changes in LC₅₀ and LC₉₀ values of imidacloprid on selected and sensitive populations of *S.littoralis* larvae

The values of LC₅ and LC₉ in F0 for imidacloprid were 369.1ppm and 955.2ppm which respectively increased to 3772.1 and 7548.8 in seventh selection (F) (Fig.2). This means that the LC₅ increased tenfold and LC₉ increased eightfold in the last selection. The RR values reached to 9.76 and 8.68 for LC50 and LC90 respectively (table 2). The heterogeneity ratio (h) in population exposed to imidacloprid decreased from 1.34 to 0.13 in 6th generation. This means that homogeneity to this insecticide increased gradually in selected populations (Table 2).

Table 2- The bioassay parameters of 7 generations of third instar larvae of *S. littoralis* exposed to Imidacloprid

RR LC ₉₀	RR LC ₅₀	LC ₉₀ ppm (Confidence limit 0.95)	LC ₅₀ ppm (Confidence limit 0.95)	Slope ±SE	h	χ ² (df)	n	Sel.
1.00	1.00	955.2 (722.5 – 1494.0)	369.1 (306.7 - 451.0)	3.10 ±0.35	1.34	29.46 (22)	216	F0
3.23	2.25	3386.9 (2388.9 – 5889.1)	921.1 (766.5 – 1142.5)	2.26 ±0.28	0.66	14.03 (21)	216	F1
5.91	5.65	5416.6 (4466.6 – 7016.7)	2215.8 (1928.7 – 2550.8)	3.30 ±0.31	0.96	20.18 (21)	216	F2
6.38	7.27	7056.4 (5881.5 – 9701.7)	3050.6 (2626.0 – 3708.4)	3.34 ±0.54	0.65	5.71 (12)	216	F3
6.68	7.75	7166.1 (5951.6 – 9653.2)	3624.5 (3250.9 – 4088.4)	4.32 ±0.57	0.47	9.81 (15)	216	F4
8.11	9.64	7371.0 (5504.1 – 12570.0)	3715.1 (2996.5 – 3925.3)	4.06 ±0.64	0.18	2.42 (13)	216	F5
8.68	9.76	7548.8 (5885.4 – 11536.0)	3772.3 (3199.7 – 4418.1)	4.09 ±0.59	0.13	2.2 (16)	216	F6

n =Number of larvae used

h =Heterogeneous value

RR =resistance ratio (LC₅₀ or LC₉₀ values of resistance population / LC₅₀ or LC₉₀ values of sensitive population)

CL=confidence limit

synergistic tests

According to the results, all three synergists increased the effect of the insecticides. The greatest impact was observed in DEM synergist, so that in the case of the simultaneous use of gamma-cyhalothrin in the LC₉₀ level, the impact of pesticide increased by 3.9 times (Table 3). At the same time, the simultaneous use of synergist and insecticide in sensitive population has no significant impact on the effectiveness of insecticides.

Table 3- Comparison of bioassay parameters Synergism of gamma-cyhalothrin by DEM, DEF and PBO

SR LC ₉₀	SR LC ₅₀	RR LC ₉₀	RR LC ₅₀	LC ₉₀ ppm (Confidence limit 0.95)	LC ₅₀ ppm (Confidence limit 0.95)	Slope ±SE	h	χ ² (df)	n	
		18.82	16.85	24340.0 (16427.0 - 47492.0)	11194.0 (9341.2 – 16705.0)	3.79 ±0.88	0.25	3,31 (13)	216	F12
		1.00	1.00	1105.0 (912.0 – 1458.1)	664.1 (558.1 – 791.9)	4.93 ±0.86	0.92	4,52 (11)	216	control
1.7	2.1	13.92	11.5	14062.0 (10848.0 – 22080.0)	5248.6 (4444.9 – 6150.8)	2.99 ±0.46	0.62	8,13 (13)	216	F12 +PBO
1.1	1.4	1.00	1.00	1010.2 (897.7 – 1318.8)	456.1 (382.6 – 535.7)	3.32 ±0.42	0.44	5,37 (12)	216	control +PBO
3.9	3.6	6.00	6.58	6177.8 (4966.0 – 8733.5)	3092.1 (2649.4 – 3663.0)	4.26 ±0.59	0.46	8,34 (18)	216	F12 +DEM
1.1	1.4	1.00	1.00	1028.9 (851.3 – 1350.6)	469.9 (400.2 – 545.2)	3.76 ±0.46	0.75	8,98 (12)	216	control +DEM
2.0	3.1	11.51	7.19	12029 (9061.5 – 20279.0)	3553.9 (2677.5- 4322.9)	2.42 ±0.42	0.21	2,84 (13)	216	F12 +DEF
1.1	1.3	1.00	1.00	1045.1 (994.9 – 1442.9)	494.1 (414.2 – 582.6)	3.19 ±0.41	0.48	5,74 (12)	216	control +DEF

n =Number of larvae used

h =Heterogeneous value

RR =resistance ratio (LC₉₀ or LC₅₀ values of resistance population / LC₉₀ or LC₅₀ values of sensitive population)

SR =Synergistic ratio (LC50 or LC90 of the insecticide / LC50 or LC90 of the insecticide plus synergist)

On the other hand, the homogeneity of the sensitive population is deductible. The use of gamma-cyhalothrin plus three synergists of DEM, DEF and PBO ,synergistic ratios (SR) for LC₅ & .LC₉ .3.6 ,3.9 ;2 ,3.1 and 2.1, 2.7 values were obtained respectively. In the case of use Imidacloprid plus DEM, DEF and PBO, SR were calculated 2.03 ,2.07 ;1.6 ,1.7 and 1.04, 1.25 at the values of LC₅ & .LC₉ .respectively (Table 4).

Table 4- Comparison of the bioassay parameters of 6th generation with the sensitive (control) population of *S. littoralis* exposed to Imidacloprid after application of the synergists DEM, DEF and PBO

SR LC ₇ 5	SR LC ₅ 0	RR LC ₇ 5	RR LC ₅ 0	LC ₉₀ ppm (Confidence limit 0.95)	LC ₅₀ ppm (Confidence limit 0.95)	Slope±SE	h	χ ² (df)	n	
		9.1 0	9.2 1	8359.2 (6323.8 - 13640)	3854.4 (3318.8 - 4757.9)	3.81 ±0.5 7	0.1 5	2.42 (16)	21 6	F6
		1.0 0	1.0 0	918.1 (716.3 - 1374.6)	418.3 (354.2 - 504.1)	3.76 ±0.5 4	0.3 8	4.63 (12)	21 6	Sensitive
1.0	1.2	9.0 6	8.4 9	8056.3 (6199.2 - 12131)	3078.7 (2561.4 - 4016.4)	2.73 ±0.4 3	0.3 4	5.1 (15)	21 6	F6 +PBO
1.1	1.1	1.0 0	1.0 0	809 (634 - 1192.2)	362.7 (306.3 - 434.1)	3.67 ±0.5 1	0.7 2	8.63 (12)	21 6	Sensitive +PBO
2.0	2.0	4.9 9	6.1 6	4121.9 (3361.3 - 5630)	1862.8 (1634.1 - 2138.2)	3.71 ±0.4 6	0.7 4	11.1 9 (15)	21 6	F6 +DEM
1.1	1.4	1.0 0	1.0 0	825.8 (620.5 - 1283.7)	302.4 (258.3 - 358.6)	2.93 ±0.3 8	0.7 7	9.3 (12)	21 6	Sensitive +DEM
1.6	1.7	6.9 9	7.7 3	5114.2 (3490.4 - 13202)	2287.7 (1874.3 - 3268.5)	3.66 ±0.6 4	1.3 8	8.5 (12)	21 6	F6 +DEF
1.3	1.4	1.0 0	1.0 0	730.9 (565.7 - 1082)	296.1 (248.8 - 358.9)	3.27 ±0.4 2	0.7 3	8.77 (12)	21 6	Sensitive +DEF

n =Number of larvae used

h =Heterogeneous value

RR =resistance ratio (LC₅ or LC₉₀ values of resistance population / LC₅ or LC₉₀ values of sensitive population(

RS =Synergistic ratio (LC50 or LC90 of the insecticide / LC50 or LC90 of the insecticide plus synergist(




Fig. 2- The trend of changes in LC₅₀ and LC₉₀ values of gamma-cyhalothrin on selected and sensitive populations of *S. littoralis* larvae

Discussion

In this study, all three synergists were effective on *S. littoralis* exposed to frequent application of gamma-cyhalothrin. Therefore, all the detoxification enzymes have efficiency to raise the resistance to both insecticide. However, DEM had the highest value of SR that it probably is an agent to inhibit the glutation *S*-transferase which has the major role in inducing resistance into *S. littoralis* (Van Leeuwen *et al.* 2004).

Pyrethroids are axonic excitoxins which the toxic effects of them are mediate through preventing the closure of the voltage-gated sodium channels in the axonal membranes. Gamma-cyhalothrin acts as a voltage-dependent sodium channel blocker belonging to the oxadiazine insecticide group. Resistance to this insecticide probably is a major cause of changes in activity or structure of this part. The resistance to synthetic pyrethroids is also due to an increase in the amount or activity of cytochrome P-450 monooxygenase enzyme. However, sometimes carboxylesterase enzyme is also effective on reducing the sensitivity of individuals to the insecticide (Delorme *et al.* 1988, Ahmad *et al.* 2007 .)

To reveal the effect of enzymatic factor(s), the synergistic tests were conducted in this study. The most commonly used insecticide synergists are metabolic inhibitors that block certain detoxification enzymes so that insecticide detoxification in target insect is abolished or significantly reduced (Capinera, 2008). PBO is a well-known inhibitor of cytochrome P-450 monooxygenases; but recently it has been shown to inhibit esterases as well (Young *et al.*, 2006) . DEF is an established esterase inhibitor, but it also acts as a substrate for cytochrome P-450 monooxygenases (Sanchez *et al.* ,2001) .There is no specific inhibitor available for glutation *S*-transferase) GST), which conjugates with and rapidly depletes glutation, has been commonly used as a synergist to suppress the GST activity (Capinera ,2008) .Yang *et al.* 2001) evaluated the susceptibility and possible detoxification mechanisms of *Oligonychus paratensis*) Banks), and two-spotted spider mite to three selected insecticides with three synergists (DEM, DEF and PBO). Their study suggests that esterases, GST and Cytochrome P-450 monooxygenase all play important roles in the detoxification of the pyrethroid, because DEF, DEM and PBO enhanced the toxicity of insecticide by 6.2, 4.1 and 4.5 fold respectively. Pacy *et al.* (2009) showed lower levels of esterase activity with DEF; lower levels of GST activity with DEM and lower levels of cytochrome monooxygenase activity with PBO by the enzyme assays .

Usually, AChE enzyme plays a major role in the resistance that is created as a result of repeated use of imidacloprid insecticide (Pittendrigh ,*et al.* , 2008) ,while, according to research, the increased activity of cytochrome P-450 monooxygenase enzyme has been involved in creating resistance mechanisms in some cases (Puinean *et al.* , 2010) .In this research, the PBO synergist had very low effect on increasing the imidacloprid efficiency. This has been approved by many researchers that cytochrome P-450 monooxygenases has no (or little) involvement in resistance ability of the pest .

References

- Abdallah, M. D., El Attal, Z. M., Moustafa Omayma, K. 1979.** Development of resistance of *Spodoptera littoralis* (Boisd.) to Du-Ter. Journal of Agricultural Science, 93: 257-261.
- Ahmad, I., Ahad, I., Gupta, R. K., Monobrullah, M., Bhagat, R. M. and Ahmad, H. 2011 .** Toxicity of conventional insecticides to fourth instar larvae of tobacco caterpillar (*Spodoptera litura*) Fab.) in different generations. Journal of Phytology, 3: 04-08 .
- Ahmad, M., Saleem, M. A., Sayyed, A. 2008.** Evidence for field evolved resistance to newer insecticides in *Spodoptera litura* (Lepidoptera: Noctuidae) (from Pakistan) Crop Prot., 27: 1367-1372 .
- Ahmad, M., Saleem, M. A., Sayyed, A. 2008.** Efficacy of insecticide mixtures against pyrethroid- and organophosphate-resistant populations of *Spodoptera litura* (Lepidoptera: Noctuidae). Pest Management Science, 65 :266–274.
- Ahmad, M., Sayyed, A. H., Crickmore, N. and Saleem, M. A. 2007.** Genetics and mechanism of resistance to deltamethrin in a field population of *Spodoptera litura* (Lepidoptera: Noctuidae). Pest Management Science, 63–1002:1010.
- Alin, M., Puinean, Ian Denholm, Neil S. Millar, Ralf Nauen, Martin S. Williamson, 2010 .** characterisation of imidacloprid resistance mechanisms in the brown planthopper *Nilaparvata lugens* Stål (Hemiptera :Delphacidae). Pesticide Biochemistry and Physiology, 97: 129–132.
- Armes, N.J., Wightman, J.A., Jadhav, D., Ranga Rao, G. 1997.** Status of insecticide resistance in *Spodoptera litra* in Andhara Pradesh, India. Pestic. Sci .,50 :240-248.
- Capinera, J. L., 2008 .**Encyclopedia of Entomology, 4: 3669-3671.
- Delorme, R., Fournier, D., Chaufaux, J., Cuany, A., Bride, J. M., Auge, D., Berge, J. B. 1988 .** Esterase metabolism and reduced penetration are causes of resistance to deltamethrin in *Spodoptera exigua* HUB. (Noctuidea ;lepidoptera). Pesticide Biochemistry and Physiology, 32: 240–246.
- Farnsworth, C. A., Teese, M. G., Yuan, G., Li Y., Scott, C., Zhang, X., Wu Y., Russell, R .J., Oakeshott, J. G. 2010.** Esterase-based metabolic resistance to insecticides in heliothine and spodopteran pests. Journal of Pesticide Science, 35: 275-289.
- Gervais, J. A., Luukinen, B., Buhl, K., Stone, D. 2010 .***Imidacloprid Technical Fact Sheet* ; National Pesticide Information Center, Oregon State University Extension Services, 5:1-7 .
- Haug, S., Han, Z. 2007 .**Mechanisms for multiple resistance in field populations of common cutworm (*Spodoptera Litura*) Fab.) in China. Pestic .Biochem. Physiol., 87: 14-22.
- Ishaa ya, I., Klein, M. 1990.** Response of susceptible laboratory and resistant field strains of *Spodoptera littoralis* (Lepidoptera: Noctuidae) to teflubenzuron. Journal of Economic Entomology, 83: 59-62.
- LeOra Software 2003 .**Poloplus, a user's guide to probit or logit analysis. LeOra Software, Berkeley, CA, USA.
- Metcalf, R. L. 2000 .**Insect Control. Ullmann's Encyclopedia of Industrial Chemistry ,Published Online: 15 JUN 2000 DOI. 14356007/10/1002.a14_263.
- Pasay, C., Arlian, L., Morgan, M., Gunning, R., Rossiter, L., Holt D., Walton, Sh., Beckham, S., McCarthy, J. 2009.** The Effect of Insecticide Synergists on the Response of Scabies Mites to Pyrethroid Acaricides .PLOS Neglected Tropical Diseases ,204-215 .DOI: 10.1371.
- Pittendrigh, B. R., Margam, V. M., Sun, L., and Huesing, J. E. 2008.** Resistance in the Post-Genomics Age, 45-47. Insect Resistance Management .David W. Onstad. Biology, Economics and Prediction, Illinois ,Urbana, USA, 318.
- Pittendrigh, B. R., Sun, L., Gaffney, P. and Huesing, J. 2008 .**Negative cross resistance”, in insect resistance management. Ed. David Onstad.1 108-124.
- Riskallah, M.R .1983 .**esterases and resistance to synthetic pyrethroids in the egyptian cotton leafworm. Pesticide Biochemistry and Physiology, 19 :184–189.

- Salama, H. S., Dimetry, N. Z.; Salem, S. A. 1970.** On the host preference and biology of the cotton leaf worm *Spodoptera littoralis* .zeitung für angewandte entomologie ,67 :261-266 .
- Sanchez-Arroyo, H., Koehler, P. G., Valles, S. M. 2001.** Effects of the synergists piperonyl butoxide and S,S,S-tributyl phosphorithioate on propoxur pharmacokinetics in *Blattella germanica* (Blattodea :Blatellidae). J. Econ. Entomol., 94: 209-216.
- TİRYAKİ, O., CANHİLAL, R., HORUZ, S. 2010.** tarım ilaçları kullanımı ve riskleri. Erciyes Üniversitesi Fen Bilimleri Enstitüsü Dergisi ,26 :154-169.
- Tong, H., Su, Q., Zhou, X., Bai, L. 2013.** Field resistance of *Spodoptera litura* (Lepidoptera: Noctuidae) to organophosphates, pyrethroids, carbamates and four newer chemistry insecticides in Hunan, China. J Pest Sci 86 :599–609.
- Topuz, E. 2005** .tarımsal zararlılarla mücadelede kimyasal pestisitlere alternatif bazı yöntemler, Derim, 22: 53-59.
- Van Leeuwen, T., Stillatus, V. and Tirry, L. 2004.** Genetic analysis and cross-resistance spectrum of a laboratory-selected chlorfenapyr resistant strain of two-spotted spider mite (Acari: Tetranychidae) .(Experimental and Applied Acarology, 32: 249-261.
- Yang, X., Zhu, K., Buschman, L., Margolies, D. C. 2001.** Comparative susceptibility and possible detoxification mechanisms for selected mitecides in banks grass mite and two-spotted spider mite (Acari: Tetranychidae). Exp .Appl. Acarol., 25: 239-299.
- Yonggyun, K., Choa, J .R., Lee, J., Kang, S., Han, S. C., Hongb, K. J., Kima, H. S., Yoob, J. K., Lee, J. O. 1998.** Insecticide resistance in the tobacco cutworm ,*Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae). Journal of Asia-Pacific Entomology, 1: 115–122.
- Young, S. J., Gunning, R. V., Moores, G. D. 2006.** Effect of pretreatment with piperonyl butoxide on pyrethroid efficacy against insecticide-resistance *Helicoverpa armigera* (Lep : Noctuidae). Pest management Sci ,62 :114-119.
- Zhou XM., Huang BO 2002** .Insecticides resistance of the common cutworm) *Spodoptera litura* (and its control strategies. Entomol. Knowl., 39: 98-102.

تأثیرات حشره کش‌های پیرتروئیدی مصنوعی و نیونیکوتینوئیدی و سینرژست‌ها روی جمعیت (*Spodoptera littoralis* (Lepidoptera: Noctuidae))

عبداله دینی پور*، اکتای گورکان^۲

۱- دانشجوی دکتری تخصصی حشره‌شناسی کشاورزی دانشگاه آنکارا

۲- گروه حشره‌شناسی دانشکده کشاورزی دانشگاه آنکارا

چکیده

Spodoptera littoralis (Lepidoptera: Noctuidae) تحت عنوان کرم برگ‌خوار پنبه مصری شناخته شده و یکی از متداولترین آفات پنبه در ناحیه آدانای ترکیه به حساب می‌آید. هدف از این تحقیق توصیف تأثیرات سموم پیرتروئیدی مصنوعی (Gamma-cyhalothrin) و نیونیکوتینوئیدی (Imidacloprid) روی این آفت به منظور یافتن روند القای مقاومت طی چندین نسل متوالی بوده است. مقاومت ایجاد شده به سم gamma-cyhalothrin تا ۱۲ نسل و به سم Imidacloprid تا ۷ نسل به وسیله محاسبه نرخ مقاومت (RR) مربوط به LC_{50} و LC_{90} در بازه خطای ۹۵٪ انجام شد. میزان مقاومت (RR) برای LC_{50} و LC_{90} سم gamma-cyhalothrin ۱۶/۸۲ و برای سم Imidacloprid به ترتیب ۹/۷۶ و ۸/۶۸ به دست آمد. به منظور بررسی تأثیر سینرژست‌ها روی سموم، سه سینرژست مختلف شامل (DEF) S, S, S-tributylfosforotritioat، piperonyl butoxide (PBO) و diethyl maleate (DEM) استفاده و به وسیله محاسبه نرخ سینرژستی (SR) نسبت به شاهد به دست آمد. در مورد هر دو سم، بیشترین مقدار SR به سینرژست DEM تعلق یافت که می‌توان نتیجه گرفت، استفاده و کاربرد این سینرژست همراه با سموم gamma-cyhalothrin و Imidacloprid می‌تواند به جلوگیری و یا به تاخیر انداختن مقاومت به این سموم مفید واقع شود.

واژه‌های کلیدی: *Spodoptera littoralis*، سنجش زیستی، مقاومت، حشره‌کش، سینرژست

* نویسنده رابط، پست الکترونیکی dinipourm@gmail.com

تاریخ دریافت مقاله (۹۱/۲/۲۲) - تاریخ پذیرش مقاله (۹۱/۹/۳۰)

