



Phyto-pharmacological review of *Calotropis procera* – A nature's drug house in tropical countries

Alok Maithani¹, Versha Parcha¹, Noor Fatma¹, Deepak Kumar¹, Santosh Kumar Karn^{2*}

¹Department of Applied Chemistry, Sardar Bhagwan Singh University, Balawala, Dehradun, India, 248100;

²Department of Biochemistry and Biotechnology, Sardar Bhagwan Singh University, Balawala, Dehradun, India, 248100;

*Email: santoshkarn@gmail.com

ARTICLE INFO

Type: Review

Topic: Medicinal Plants

Received June 06th 2020

Accepted September 21th 2020

Key words:

- ✓ *Calotropis procera*
- ✓ Phyto-constituents
- ✓ Pharmacological activity
- ✓ Latex

ABSTRACT

Background & Aim: The use of traditional medicine and medicinal plants in most developing countries, as a normative basis for the maintenance of good health, has been widely observed. In the last century, approximately 121 pharmaceutical products have been discovered based on the information obtained from the traditional healers. In the present study, we thoroughly investigated the phyto-pharmacological potential of *Calotropis procera* which is extremely effective as medicine and its phyto-constituents show strong antioxidant properties.

Results: The world health organization has estimated that world population of about 80 % from developing countries depends on herbal medicine for their basic health care needs. *Calotropis procera* is an important drug of Ayurvedic medicine and researchers are exploring the therapeutic potential of this plant. *Calotropis* is a plant with excellent medicinal properties which is used for curing different human ailments.

Recommended applications/industries: The present review article provides a viewpoint on phytopharmacological potential of *Calotropis procera* which could be beneficial for future research on development of better and economically superior therapeutic agents.

1. Introduction

Calotropis procera is a drought resistant, salt tolerant, shrub of India and an important drug of Ayurveda (Anonymous, 2010). It is native of Asia and widely found in China, Malaysia and Vietnam. It is a small, erect and compact shrub, which is used in several traditional medicines to cure various diseases. There are two common species of *Calotropis*, viz. *Calotropis gigantea* (Linn.) R.Br. and *Calotropis procera* (Ait.) R.Br. described by the Sanskrit writers (Parrotta, 2001). Both the species are used as substitutes for one another and are said to have similar effects. It grows in varieties of soils and environmental conditions and don't require

cultivation practices. Each part of the plant bears excellent medicinal properties and used for curing different human ailments. The decoction of the plant is used in Indian traditional medicine for the treatment of painful muscular spasm, dysentery, fever, rheumatism, asthma and as an expectorant and purgative. Taxonomic Classification of the plant is as follows (Grieve, 1994; Ali *et al.*, 1996):

Kingdom: Plantae

Subkingdom: Tracheobionta (Vascular Plants)

Superdivision: Spermatophyta (Seed Plants)

Division: Magnoliophyta (Flowering Plants)

Class: Magnoliopsida (Dicotyledones)

Sub class: Asteridae
Order: Gentianales
Family: Apocynaceae (Milkweed Family)
Genus: *Calotropis*

Species: *procera*
 The various common and popular names of *Calotropis procera* are discussed in Table 1 (Palejkar et al., 2012).

Table 1. Various common names of *Calotropis procera* in different place.

Countries	Synonyms	India	Synonyms
English	Crown flower, Giant Indian milkweed	Hindi	Madar, Aak
Malaysia	Remiga, Rembega	Sanskrit	Ravi
Indonesia	Bidhuri, Sidaguri	Assamies	Akand
Philippines	Kapal-kapal (Tagalog)	Bangoli	Akon
Laos	Kok may, Dok kap, Dok hak	Cannad	Ekka
Thailand	Po thuean, Paan thuean, Rak	Kashmiri	Acka
Vietnam	Bootng	Malyalm	Erikku
French	Faux arbre de soie	Marathi	Rui
German	Wahre Mudarpflamzer	Urea	Arkha
Italian	Calotropo	Panjabi	Ak
Spanish	Algodon extranjero	Urdu	Madar
Turkish	Ipekag	Telgu	Gilledu
Arabic	Oshar	Gujarati	Aakando

2. Botanical description

Calotropis procera (CP) is a perennial, evergreen and soft wooded shrub. It usually attends heights of 3 meter and rarely reaches to 5 meter. White coloured latex is obtained when leaves or stems are cut (Rastogi et al., 1997; Kokate et al., 1994; Evans et al., 2005; Alikhan et al., 2005).

2.1. Leaves

The leaves are opposite and externally shorts petioles belongs to hearts shaped base .The blades are broadly elliptic or nearly orbicular, short pointed to blunt at the apex. The leaves are light to dark green and contain nearly white veins. These are 7-18 cm long and 5-12 cm broad, slightly leathery and cover with fine coat of soft hairs that are easy to rub off (Figure 1a).

2.2. Flowers

The species having white flowers is a most common and superior variety and is referred to as *Calotropis procera*. The corolla is slightly succulent and consist of 5 small triangular dirty white sepals, 5 thick ovate petals (1 cm²) (Figure 1d).

2.3. Fruits

The fruits are green inflated, and spongy ovoid follicles that split open to release plumed, papery light brown seeds with white filaments. These are 3-4.5

inches long and 2-2.5 inches wide. The main flowering period is from March to October (Figure 1b).

2.4. Root and root bark

Giant milkweed has a very deep, stout taproot with few or no near-surface lateral roots. Giant milkweed roots were found to have few branches and reach depths of 1.7 to 3.0 m in Indian sandy desert soils. The roots are covered with root bark with rounded head and rest of the portion spirally curved. The bark of the older root is cracked at places. The bark is yellowish gray outside and yellowish white inside. The upper cork portion is spongy and rough while the inner portion of bark is smooth and mucilaginous (Figure 1e).




Figure 1. a) Leaf, b) Fruit, c) Stem, d) Flower, e) Root.

3. Phyto-pharmacological discussion

Each part of *Calotropis procera* (CP) contains several phytoconstituents belonging to different classes of natural products, which makes the genus *Calotropis* as the nature's drug house. About 70 chemicals have been isolated and characterized from different part of this plant. The availability of various compounds and important pharmacological activity associated to each part of *Calotropis procera* is discussed in this paper.

3.1. Flower

Flower of *Calotropis procera* is rich in cardiac glycosides, carbohydrates, enzymes and steroidal contents. The major cardiac glycosides are calotropin (1), calactin (2), calotoxin (3), calotropagenine (4), uscharin (5), voruscharin (6), uscharidine (7), uzarigenin (8), proceragenine (9) (Figure 2). The carbohydrates present in flowers are D-arabinose (10), glucose (11), glucosamine (12), and rhamnose (13). However, the most studied sterols in flower are lupeol (14), stigma sterols (15), cycloart-2,3-ene-3-B-25-diol (16), multiflorenol (17), clycosadol (18), 3-epimoratenol, prosterol (19) (Figure 3). Two enzymes namely 3-protinase, calotropain (20), have also been isolated from the flowers of *Calotropis procera*. The flower also contain one calotropenyl acetate (21)

(Figure 3). The flowers of the plant exhibit hepatoprotective activity, anti-inflammatory, antipyretic, analgesic and antimicrobial effect and larvicidal activity (Ansari *et al.*, 1999; Meena *et al.*, 2011). Structures of all the phytochemicals are shown in Figure 3.

Zafar *et al.* (2005) studied the effect of crude aqueous and methanolic extract of flowers of *Calotropis procera* on live *Haemonchus contortus* and evaluated the rate and extent of mortality and temporary paralysis. In their experiment, the extracts of *Calotropis procera* flowers showed good antihelmentic activity in sheep infected with nematodes.

Setty *et al.* (2007) reported hepatoprotective effect of ethanolic extract (70%) against paracetamol induced hepatitis in rats. The treatment with ethanolic extract (200mg and 400mg/kg) restored the altered levels of biochemical markers (SGPT, SGOT, ALP, Bilirubin and cholesterol, HDL, and tissue GSH) in a dose dependent manner.

Mukharjee *et al.* (2010) reported the CNS depressant activity of methanolic extracts of fresh flowers of *calotropis procera* in an *in-vivo* model. Administration of Diazepam (1.5mg/kg body weight ip) and flower extracts (100mg/kg oral) produces significant CNS depression to the treated mice.




Figure 2. Structures of phyto-constituents present in *Calotropis procera* flower. calotropin (1), calactin (2), calotoxin (3), calotropagenine (4), uscharin (5), voruscharin (6), uscharidine (7), uzarigenin (8), proceragenine (9).

Sharma and Sharma (2000) studied the antimarial effects of ethanolic extracts of different parts of *calotropis procera*. The IC₅₀ values were obtained from

0.11-0.47mg/ml against *P. falciperum*. It was observed that flower and bud extracts are most active antimicrobial component in *Calotropis procera*.




Figure 3. Structures of carbohydrates, sterols and enzymes present in flowers of *Calotropis procera*. D-arabinose (10), glucose (11), glucosamine (12), rhamnose (13), lupeol (14), stigma sterols (15), cycloart-2,3-ene-3-B-25-diol (16), multiflorenol (17), clycosadol (18), 3-epimoratenol, prosterol (19), 3-protinase, calotropain (20), calotropenyl acetate (21).

3.2. Latex

Latex is a complex mixture of more than 9 cardiac glycosides which are calotoxin, calactin, uscharine, voruscharine, uzarigenine, syriogenin (22), proceroside (23), cholin (24). One enzyme named as trypsine was also isolated from latex of *calotropis procera* (25) (Figure 4). The presence of these compounds enables

latex to use as anti-inflammatory, analgesic, antinociceptive and antidiabetic remedy in folklore medicines. The Latex of the plant is reported to possess antiasthmatic, antileprotic, antirheumatism, anticold activity (Murti *et al.*, 1943; Atef *et al.*, 1999; Kew *et al.*, 1985).




Figure 4. Glycosides and enzyme isolated from latex of *calotropis procera*: syriogenin (22), proceroside (23), cholin (24), trypsin (25).

Kumar *et al.* (2000) studied the analgesic effect of *calotropis procera* latex against acetic acid induced writhing drug. Administration of dry latex at a dose of 830 mg/kg produces marginal analgesia in a tail-flick rodent model. This dose of latex did not show any toxic effects in mice and LD₅₀ was found to be 3000 mg/kg.

Vasconcelos *et al.* (2005) administered fraction of *calotropis procera* latex protein to male mice and observed a dose-dependent antinociceptive effect compared with the respective controls in all assays. Different doses of latex protein reduced the nociception produced by formalin and the effect was not reversed by pretreatment with naloxone. They observed that protein fraction of latex possesses good antinociceptive activity.

Kumar *et al.* (2005) observed that daily oral administration of latex in diabetic rats at 100 and 400 mg/kg caused a dose-dependent decrease in blood glucose and an increase in hepatic glycogen. It also restores the body weight and daily water consumption capacity of the treated animals. Dry latex also increases hepatic levels of endogenous antioxidants such as superoxide dismutase (SOD), catalase, glutathione and reduces the levels of thiobarbituric acid reactive substance (TBARS) in alloxan-induced diabetic rats.

Al-Yahya *et al.* (1985) studied the latex of *Calotropis procera* for its inflammatory activity in rats using pedal oedema and air pouch models of inflammation. Subcutaneous injection of aqueous solution of dry latex into the plantar surface of paw induces inflammation. Maximum inflammation obtained one hour after the

injection and maintained for next one hour. The inflammatory activity (response) accompanied by an increase in vascular permeability that reaches its maximum within 15 minutes. The model characterized for the exudates volume and its protein concentration and wet and dry weights of granuloma. Their results indicated maximum exudates volume and weight of granuloma on day 5 after dry latex injection and the protein concentration peaked on the 3rd day. Further study of both the model for the anti-inflammatory effect of various drugs, they observed that in paw oedema model, phenylbutazone was more effective than prednisolone and almost complete inhibition was observed by mepiramine and cycloheptadine. In air pouch model, prednisolone was more effective than phenylbutazone in inhibiting the inflammation.

Kumar *et al.* (2001) studied the anti-diarrhoeal activity of *Calotropis procera* dry latex. They evaluated its effect on intestinal transit, castor oil-induced intestinal fluid accumulation and electrolyte concentration in intestinal fluid. Dry latex caused a decrease in intestinal transit (27%-37%) compared with normal and castor oil-treated animals. Dry latex significantly inhibited castor oil-induced enter pooling, but it didn't alter the electrolyte concentration in the intestinal fluid compared with castor oil-treated rats.

Uddin *et al.* (2012) studied the antibacterial effect of various extracts of milky latex of *Calotropis procera* using modified agar well diffusion method. Streptomycin was used as standard drug at the concentration of 2mg/ml. The crude methanolic extract

exhibited good antibacterial activity with zone of inhibition ranging from 12-20 mm diameter.

3.3. Roots and root bark

Root and root bark of *Calotropis procera* is rich in terpenoids, steroids, heterocyclics and naphthalene derivatives. Terpenoids were mainly Calotropaleanyl ester (26), proceroleanol A and B (27, 28), Calotropesquiterpenol (29), Calotropsesterpenol (30), Calotropursenyl acetate, quercetine -3- ratinoside (31). One naphthalene derivative namely Calotropnaphthalene (32) was isolated from the root bark. Few sterols like benzoyllineolone (33), benzoylisolineolane (34) and teraxasterol (35) have been isolated and characterized from the roots of *Calotropis procera*. One heterocyclic compound named as chlorobenzofuranone (36) has also reported from the roots. Compounds like procerursenyl acetate (37), proceranol -2 (38), n- tetradecanyl palmitoleate (39), proceraursenolide (40), pyrocatechuic acid (41) are also isolated from the root of *Calotropis procera* (Figure 5). The root and root bark of plant are reported to have antidiarrheal and anticholeral activity (Ansari et al., 2001; Akhtar et al., 1998; Murti et al., 1945; Gupta et al., 2000; Alam et al., 2009; Ansari et al., 2000; Mittal et al., 2011; Parrota et al. 2001).

Jalalpure et al. (2009) studied anticonvulsant activity of root extract of *Calotropis procera* in rats induced by seizures using maximal electroshock seizures (MES), pentylenetetrazol (PTZ), lithiumpilocarpine and kindling seizures. Chloroform extract of *Calotropis procera* roots showed the most significant ($P < 0.01$) anticonvulsant effect by decreasing the duration of hind limb extension in the MES test. In the PTZ test, chloroform extract exhibited highly significant effect ($P < 0.001$) and aqueous extract exhibited most significant ($P < 0.01$) effect. By the study, they observed

that the chloroform extracts and aqueous extract of roots are useful in the absence and tonic clonic (grand mal), (petitmal) types of seizures.

Mathura et al. (2009) studied the anti-tumor activity of various root extracts of *calotropis procera*. Treatment of step 2 cancer cells with these extracts at different dosage of 1,5,10 and 25 mg/ml revealed that all the extracts, except aqueous, possesses good cytotoxic effects. Among them, ethyl acetate extract showed strongest cytotoxic effect (96.3%) followed by methanolic (72.7%) and hexane extract (60.5%).

Danmalam et al. (2007) studied the anticonvulsant of 70% ethanolic extract of root bark of *Calotropis procera* using maximum electroshock induced seizures in chicks. The extracts proved to be effecting in inhibiting electroshock induced seizures in chicks. The observed inhibition effect in hind limb tonic extension chicks was comparable to the effect of phenitoin (100% protection) in this model. Thus, they suggested the usefulness of ethanolic extracts of *Calotropis procera* root bark in management of grandma epilepsy.

3.4. Leaves

The leaves of *Calotropis procera* have been reported to possess cardiac glycosides, terpenoides and sterols. Two major cardiac glycosides isolated from the leaves are calotropin and calotropagenin, where as terpenoids were mainly alpha-amyrin (42), beta-amyrin and amyrin acetate (43). Beta sitosterol (44) was the only steroid present in the leaves of *Calotropis procera*. Ursolic acid (45), ratinoside (46) and 3-thiozoline (47) was also isolated from the leaves of *Calotropis procera* (Figure 6). The leaves of plant exhibit antirheumatism and antiasthmatic activity (Palejkar, et al., 2012; Jan, et al., 2008). Phytopharmacological details of *Calotropis procera* are summarized in Table 2.




Figure 5. Terpenoids, steroids, heterocyclics and naphthalene derivatives in root and root bark of *Calotropis procera*: Calotropaleanyl ester (26), proceroleanol A and B (27, 28), Calotropsesquiterpenol (29), Calotropsesterpenol (30), Calotropursenyl acetate, quercetin -3- ratinoside (31), Calotropnaphthalene (32), benzoyllineolone (33), benzoylisolineolane (34), teraxasterol (35), chlorobenzofuranone (36), procerursenyl acetate (37), proceranol -2 (38), n- tetradecanyl palmitoleate (39), procaursenolide (40), Pyrocatechuic acid (41).




Figure 6. Cardiac glycosides, terpenoides and sterols of *Calotropis procera* leaves: alpha-amyrin (42), beta-amyrin, amyrin acetate (43), beta sitosterol (44), ursolic acid (45), ratinoside (46) and 3-thiozoline (47).

Yashmin *et al.* (2008) evaluated antioxidant potential of methanolic and aqueous extract of leaves of *Calotropis procera* using scavenging activity of the

stable DPPH free radical. IC₅₀ of the methanolic extract was 110.25 mg/ml indicating strong antioxidant effect of aqueous extract.

Table 2. Phyto-pharmacological detail of various parts of *Calotropis procera*

Parts	Chemistry	Pharmacological Activity	Reference
Flower	Cardiac glycoside	Heptoprotective,	(Setty <i>et al.</i> , 2007)
	Calotropagenin	Anti- inflammatory	(Markouk <i>et al.</i> , 2000; Mascolo <i>et al.</i> , 1989)
	Calotropin	Antimicrobial,	(Markouk <i>et al.</i> , 2000; Mascolo <i>et al.</i> , 1989)
	Usharin	Larvicidal	(Jain <i>et al.</i> , 2008)
	Calotoxins	Antimalarial	(Kumar <i>et al.</i> , 2009)
	Calactin	Antiasthmatic	(Anis <i>et al.</i> , 2000)
	Voruscharin	Antimoebic activity	
	Usharidin		
	Uzarigenin		
	Proceroside		
	Proceragenin		
	Syriogenin		
	Carbohydrates		
	D- Arabinose		
	Glucose		
	L- Rhamnose		
	Glucosamine		
	Sterols		
	Proesterol		
	Multiflorenol		
Cyclosadol			
Cycloart-23-Ene-3b-25 Diol			
Lupeol			

	Stigmasterol		
	Calotropain		
	Calotropenyl acetate		
Latex	Cardiacglycosides		
	Calotropin	Antirabies	(Kumar, <i>et al.</i> , 1994)
	Calotoxin	Arthritis	(Negi, <i>et al.</i> , 2002)
	Calactin	Analgesic, Anti-	
	Uscharin	inflammatory, antimicrobial,	
	Voruscharin	Antiliprotic	
	Uzariogenin		
	Syriogenin		
	Proceroside		
	Trypsin		
	Choline		
Leaves	Cardiac glycosides		
	Calotropagenin	Antigout	(Reddy, <i>et al.</i> , 2008)
	Calotropin	Antileprotic	(Reddy, <i>et al.</i> , 2008)
	α -Amyrin	Antimalarial	(Bhogaonkar, <i>et al.</i> , 2007)
	β -Amyrin	Antirhumatism	(Reddy, <i>et al.</i> , 2008)
	Amyrin Acetate	Analgesic	(Sha, <i>et al.</i> , 2006)
	β -Sitosterol		
	Urosolic Acid		
	Ratinoside		
	3-thiozolin		
Root and root bark	Sterols	Anticanceractivity	(Dhar, <i>et al.</i> , 1968)
	Benzoyllineolone	Antimalarial	(Joshua, <i>et al.</i> , 2006)
	Benzoylisolineolane	Antirheumatism, Antiasthmatic,	(Sen, <i>et al.</i> , 2007)
	Proceranol-2	Antileprotic	
	Teraxasterol		
	Triterpenoids		
	Calotropsesquiterpenol		
	Calotropsesterpenol		
	Calotropleanyl Ester		
	Proceraleanol A		
	Proceraleanol B		
	Others		
	Calotropnaphthalene		
	Calotropbenzofuranone		
	Procerursenyl Acetate		
	N- Tetradecanyl Palmitoleate		
	Proceraursenolide		
	Pyrocatechuric Acid		
	Queretin-3-retinoside		

From the above vast literature search it was observed that *Calotropis procera* itself contains enormous potential to cure various chronic ailment and is being recommended by various popular systems of medicines. Keeping this aim in mind, *Calotropis procera* latex was taken and preliminary phytochemical screening and antimicrobial activity assay was performed with various extracts of *Calotropis procera* latex. Significant effect was found in petroleum ether extract and the study further elaborated for investigation of antimicrobial principal (if any), which is under progress.

4. Conclusion

Treatment of a number of chronic diseases is still a big challenge before modern medicinal system and World Health Organization has recognised that there are strict requirements of alternative medicines in treatment and cure of such complex diseases. *Calotropis procera* and other species of this plant are well known for a number of phyto-pharmaceutical agents which are been used for their excellent therapeutic properties by Ayurvedic and other traditional practitioners. The present detailed review article thus provides a viewpoint on

phytopharmacological potential of *Calotropis procera* which could be beneficial for future research on development of better and economically superior therapeutic agents.

5. Acknowledgment

The authors are thankful to Ms Monika Khatri, MSc Pharmaceutical Chemistry, SBS PGI, Balawala, Dehradun for drawing beautiful sketch of plant parts shown in Figure 1. Authors are also thankful to Ms Neelam Kandari. Further authors are thankful to Dr. Gaurav Deep Singh, Managing Secretary, Sardar Bhagwan Singh University, Balawala, Dehradun, Uttarakhand, India, for providing facility, space and resources to conduct this work.

6. References

- Akhtar, N. and Malik, A. 1998. Proceragenin, an antibacterial cardenolide from *Calotropis procera*. *Phytochemistry*, 31(8): 2821-2824.
- Alam, P. and Ali, M. 2009. Phytochemical investigation of *Calotropis procera* (Ait) roots. *Indian Journal of chemistry*, 48B: 443-446.
- Ali, A.M. Mackeen M.M, and El- Sharkawy S.H. 1996. Antiviral and cytotoxic activities of some plants used in Malaysian indigenous medicines. *Pertanika Journal of Tropical Agricultural Science*, 19: 129.
- Alikhan, I. and Khanum, A. 2005. Medicinal and Aromatic Plants of India, Ukaaz Publication, 133-134.
- Al-Yahya, M.A., Al-Meshal I.A., Mossa, J.S., and Tariq, M. 1985. Phytochemical and pharmacological studies on *Calotropis procera*. *Proceeding of the 3rd International Conference of Traditional and Folk Medicine*. Lecatecas, Mexico.
- Anis, M., Sharma, M.P. and Muhammad, I. 2000. Herbal ethno-medicine of the Gwalior Forest Division in Madhya Pradesh, India. *Pharmaceutical Biology*, 38(4): 241-253.
- Anonymous, 2010. The Ayurvedic Pharmacopoeia of India. Government of India Ministry of Health and Family Welfare Department of Ayush, New Dehli, India, 1(1): 11-14.
- Ansari, S.H. and Ali, M. 2000. New oleanene triterpenes from root bark of *Calotropis procera* (Ait) R.Br. *Indian journal of chemistry*, 39B: 287-290.
- Ansari, S.H. and Ali, M. 1999. New oleanene triterpenes from root bark of *Calotropis procera*. *Medicinal and Aromatic Plant Science*, 21(4): 978-981.
- Ansari, S.H. and Ali, M. 2001. Norditerpenic ester and pentacyclic triterpenoids from root bark of *Calotropis procera* (Ait) R. Br. *Pharmazie*, 56(2):175-177.
- Atef G.H., Elgamal, M.H.A., Morsy, N.A.M., Duddeck, H., Kovacs, J. and Toth, G. 1999. Two cardenolides from *Calotropis procera*. *Journal of Magnetic Resonance Chemistry*, 17: 754-757.
- Bhogaonkar, P.Y. and Kadam, V.N. 2007. Herbal antidotes used for snakebite by banjara people of Umardhed region in Maharashtra, India. *Ethnobotany*, 19: 71-78.
- Danmalan, U.H., Abdul, R.A., Yaro, A.H. and Magaji, M.G. 2007. Phytochemical and anticonvulsant activity of the ethanol extracts of the root bark of *calotropis procera*. *Chemical Class Journal*, 4: 113-116.
- Dhar, M.L., Dhar, M.M., Dhawan, B.N., Mehrotra, B.N. and Ray C. 1968. Screening of Indian medicinal plants for biological activity, Part I. *Indian Journal of Experimental Biology*, 6: 232-247.
- Evans, W.C. 2005. Trease and Evans Pharmacognosy. Saunders an Imprint of Elsevier, 41-47.
- Grieve, M.M. 1994. A Modern Herbal. London: Tigers Book International, 10: 154.
- Gupta, J. and Ali, M. 2000. Rare chemical constituents from *Calotropis gigantea* roots. *Indian Journal of pharmaceutical sciences*, 62(1),29-32.
- Iqbal, Z., Lateef, M. and Jabbar, A. 2005. Anthelmintic activity of *Calotropis procera* (Ait.), flowers in sheep. *Journal of Ethnopharmacology*, 102 (2): 256-261.
- Jain, A., Katewa, S.S., Galav, P. and Nag, A. 2008. Some therapeutic uses of biodiversity among the tribals of Rajasthan, India. *Journal of Traditional Knowledge*, 7 (2): 256-262.
- Jalalpure, S.S. 2009. Anticonvulsant effects of *Calotropis procera* root in rats. *Pharmaceutical biology*, 47 (2): 162-167.
- Jan, S., Ajabkhan, M., Uddin, S., Murad, W., Hussain, M. and Ghani, A. 2008. Herbal remedies used for gastrointestinal disorders Kaghan Valley, Nwep, Pakistan. *Pakistan Journal of Weed Science Research*, 14(3-4):169-200.

- Joshua, K. 2006. Conservation of indigenous medicinal botanicals in Ekiti State, Nigeria. *Journal of Zhejiang University Science B*, 7(9): 713-718.
- Kew, F. 1985. The useful plants of west tropical Africa, 2nd ed. Vol. 1. In Burkill, H.M. *Royal Botanical Gardens*, pp. 219-222.
- Kokate, C.K. 1994. Practical pharmacognosy. Vallabh Prakashan; pp. 115-121.
- Kumar, A. 2009. Some important medicinal plants from different regions of Rajasthan, Scientific Biology, Science.
- Kumar, V.I. and Basu, N. 1994. Anti-inflammatory activity of the latex of *Calotropis procera*. *Journal of Ethnopharmacology*, 44: 123-125.
- Kumar, V.L., Padhy, B.M., Sehgal, R. and Roy, S. 2005. Antioxidant and protective effect of latex of *Calotropis procera* against alloxan-induced diabetes in rats. *Journal of Ethnopharmacology*, 102 (3): 470-473.
- Kumar, V.L., Sangraula, H., Dewan, S. and Kumar, S. 2001. Antidiarrheal activity of the latex of *Calotropis procera*. *Journal of Ethnopharmacology*, 76 (1):115-118.
- Kumar, V.L., Sangraula, H. and Dewan, S. 2000. Preliminary studies on the analgesic activity of latex of *Calotropis procera*. *Journal of Ethnopharmacology*, 73(1-2): 307-311.
- Markouk, M., Bekkouche, K., Larhsini, M., Bousaid, M., Lazrek, H.B. and Jana, M. 2000. Evaluation of some Moroccan medicinal plant extracts for larvicidal activity. *Journal of Ethnopharmacology*, 73 (1-2): 293-297.
- Mascolo, N., Jain, R., Jain, S.C. and Capasso, F. 1989. Ethnopharmacologic investigation of ginger (*Zingiber officinale*). *Journal of Ethnopharmacology*, 27 (1-2): 129-140.
- Mathura, R. and Gupta, S.K. 2009. Anti-tumor studies with extracts of *Calotropis procera* (Ait.) R.Br. root employing Hep2 cells and their possible mechanism of action. *Indian Journal of Experimental Biology*, 47 (5): 343-348.
- Meena, A.K., Yadav, A. and Raw, M.M. 2011. Ayurvedic uses and pharmacological activities of *Calotropis procera* Linn. *Asian journal of traditional medicines*, 6(2): 45-53.
- Mittal, A. and Ali, M. 2011. Phytochemical investigation of the roots of *Calotropis procera* (Ait) R.Br. *international journal of pharmacy*, 2(9):52-54.
- Mukherjee, B., Bose, S. and Dutta, S.K. 2010. Phytochemical and pharmacological investigation of fresh flower of extract of *Calotropis procera* linn. *International Journal of Pharmaceutical Sciences and Research*, 1(12): 182-187.
- Murti, P.B.R. and Seshadri, T.R. 1943. Chemical composition of *Calotropis gigantea*: Part1. Wax and Resin components of the latex. *Proceeding of the Indian Academy Science*, 18: 145-159.
- Murti, P.B.R. and Seshadri, T.R. 1945. Chemical composition of *Calotropis gigantea* : Part V. Further examination of the latex and root bark. *Proceeding of the Indian Academy Science*, 21: 143-147.
- Negi, C.S., Navtiyal, S., Dasila, L., Rao, K.S. and Maikhuri, R.K. 2002. Ethnomedicinal plant uses in a small tribal community in a part of central Himalaya, India, *Journal of Human Ecology*, 14 (1): 23-31.
- Palejkar, C.J., Palejkar, J.H., Patel, M.A. and Patel, A.J. 2012. A Comprehensive review on plant *Calotropis gigantea*. *International Journal of Institutional Pharmacy and Life Sciences*, 2(2): 463-470.
- Parrota, J.A. 2001. Healing of plants of Penninsular India. AB International Wallingford, U.K., pp. 944
- Rastogi, R.P. and Mehrotra, B.N. 1997. Compendium of Indian Medicinal Plant, Central Drug Research Institute, Lucknow, pp. 147.
- Reddy, V.B. 2008. Use of various bio fencing plants in the control of human diseases by Lambda Tribe Inhabiting Nagaland District, A.P, India. *Ethnobotanical Leaflets*, 12: 520-523.
- Sen, S.K. and Behra, L.M. 2007. Ethnomedicinal plants used in touch therapy at Bargarh district of Orissa. *Ethnobotany*, 19: 100-104.
- Setty, R.S., Quereshi, A.A., Swamy, V., Patil, T., Prakash, T., Prabhu, K. and Gouda, V. 2007. Hepatoprotective activity of *Calotropis procera* flowers against paracetamol-induced hepatic injury in rats. *Fitoterapia*, 78 (1): 451-454.
- Sha, S.R.U., Hassan, G., Rehman, A. and Ahmed, I. 2006. Ethnobotanical studies of the flora of the district Musakhel and Barkhan in Balochistan. Pakistan. *Pakistan Journal of Weed Science Research*, 12(3): 199-211.
- Sharma, P. and Sharma, J.D. 2000. In-vitro Schizontocidal screening of *Calotropis procera*. *Fitoterapia*, 71: 77-79.

- Uddin, G., Rauf, A., Muhammad, N. and Mulik, N. 2012. Phytochemical and pharmacological studied of the whole plant of *Calotropis procera*. *Middle-East Journal of medicinal Plants Research*, 1(4): 71-74.
- Vasconcelos, S.M.M. 2005. Antinociceptive activity of *Calotropis procera* latex in mice. *Journal of Ethnopharmacology*, 99(1): 125-129.
- Yasmin, N., Uddin, S.N., Mubassara, S. and Akond, M.A. 2008. Antioxidant and antibacterial activity of *calotropis procera* linn. *Ammerican –Eurasian Journal of agricultural. and environmental science*, 4(5): 550-553.