

Fixed Point Theorems in Orthogonal Intuitionistic Fuzzy b-metric Spaces with an Application to Fredholm Integral Equation

Fahim Uddin , Muhammad Saeed , Khaleel Ahmed , Umar Ishtiaq , Salvatore Sessa* 

Abstract. In this manuscript, the concept of an orthogonal intuitionistic fuzzy b-metric space is initiated as a generalization of an intuitionistic fuzzy b-metric space. We presented some fixed point results in this setting. For the validity of the obtained results, some non-trivial examples are given. In the last part, we established an application on the existence of a unique solution of a Fredholm-type integral equation.

AMS Subject Classification 2020: 47H10; 54H25

Keywords and Phrases: Orthogonal set, Intuitionistic fuzzy metric space, Unique solution, Integral equation.

1 Introduction

A publication showing there are solutions to differential equations established fixed-point theory in the second quarter of the eighteenth century (Joseph Liouville, 1837). This approach was further improved as a sequential approximation technique (Charles Emile Picard, 1890), and in the setting of complete normed space, it was generalized as a fixed-point theorem (Stefan Banach, 1922). It presents the a priori and a posteriori approximations for the convergence rate as well as a general way to actually determine the fixed point. Additionally, it ensures that a fixed point exists and is distinct. This information is helpful for studying metric spaces. Stefan Banach is acknowledged for developing fixed-point theory after that. Fixed-point theorems allow us to guarantee that the main problem has been resolved, as has the existence of a fixed point for a given function. In a large variety of scientific problems that are derive from many different branches of mathematics, the existence of a solution is equivalent to the existence of a fixed point for a suitable mapping.

In 1989, Bakhtin [2] established the notion of quasi-metric spaces and established some results for contraction mappings. In 1993, Czerwik [4] established the concept of b-metric spaces and discussed several fixed-point results. Eshaghi et al. [8] introduced the notion of orthogonal metric spaces and derived well-known Banach fixed point theorem. Uddin et al. [27] established orthogonal m-metric spaces and solve the integral equation. Eshaghi and Habibia [7] derived several fixed point results in the context of generalized orthogonal metric space. Senapati et al. [23] established some new fixed point theorems in the context of orthogonal metric spaces. In 1965, Zadeh [28] established the notion of fuzzy sets (FSs) to deal with those problems that do have not any clear boundaries.

*Corresponding Author: Salvatore Sessa, Email: salvatore.sessa2@unina.it, ORCID: 0000-0002-4303-2884

Received: 21 March 2024; **Revised:** 18 June 2024; **Accepted:** 21 June 2024; **Available Online:** 21 June 2024; **Published Online:** dd mmmm yyyy.

How to cite: Uddin F, Saeed M, Ahmed K, Ishtiaq U, Sessa S. Fixed point theorems in orthogonal intuitionistic fuzzy b-metric spaces with an application to fredholm integral equation. *Transactions on Fuzzy Sets and Systems*. 2024; 3(2): pp-pp. DOI: <http://doi.org/10.xxxxx/xxxx.yyyyy.zzzz>

In 1960, Schweizer [21] introduced the notion of continuous t-norm and worked on statistical metric spaces. In 1975, the combination of metric spaces and FSs, named fuzzy metric spaces (FMSs), have been introduced by Kramosil and Michlek [14]. In 1994, George and Veeramani [9] modified the notion of FMSs and gave an interesting analysis of FMSs in 1997 in a research paper [10]. Deng [5] established the notion of fuzzy pseudo-metric spaces and proved neumours results in the existence and uniqueness of a solution. Shukla and Abbas [24] established the notion of fuzzy metric-like spaces as a generalization of FMSs. Hezarjaribi [11] established the notion of orthogonal FMSs as a generalization of FMSs. Ndban [17] established the concept of fuzzy b-metric spaces (FBMSs) and Javed et al. [12] introduced fuzzy b-metric like spaces as a generalization of FBMSs. The authors [22, 6, 20, 15] derived several fixed points results under some circumstances in the context of FBMSs. In 2004, Park [18] introduced the notion of intuitionistic fuzzy metric spaces (IFMSs), in which he combined the notions of continuous t-norm, continuous t-conorm, FSs and metric space.

Rafi and Noorani [19], Sintunavarat and Kumam [25], Alaca et al. [1] and Mohamad [16] derived some fixed point results for contraction mappings in the context of IFMSs. Konwar [13] introduced the notion of intuitionistic fuzzy b-metric spaces (IFBMSs) as a generalization of IFMSs and derived fixed point results. Baleanu and Rezapour [3] and Sudsutad and Tariboon [26] worked on fractional differential equations. In this manuscript, we aim to toss the notion of orthogonal Intuitionistic fuzzy b-metric spaces (OIFBMSs) as a generalization of IFBMSs. We provide some related fixed point theorems, including non-trivial examples and an application. Some of the following notions are used throughout this paper, as CTN for a continuous t-norm, CTCN for a continuous t-conorm and FP for fixed point.

2 preliminaries

In this section, we will discuss some important definitions that support our main result.

Definition 2.1. [2] Suppose $\Xi \neq \phi$. Given a five tuple $(\Xi, G, H, *, \Delta)$ where $*$ is a CTN, Δ is a CTCN, $\theta \geq 1$ and G, H are FSs on $\Xi \times \Xi \times (0, \infty)$. If $(\Xi, G, H, *, \Delta)$ meets the below conditions for all $w, k \in \Xi$ and $\pi, \tau > 0$:

$$(B1) \quad G(w, k, \tau) + H(w, k, \tau) \leq 1;$$

$$(B2) \quad G(w, k, \tau) > 0;$$

$$(B3) \quad G(w, k, \tau) = 1 \Leftrightarrow w = k;$$

$$(B4) \quad G(w, k, \tau) = G(k, w, \tau);$$

$$(B5) \quad G(w, e, \theta(\tau + \pi)) \geq G(w, k, \tau) * G(k, e, \Pi);$$

$$(B6) \quad G(w, k, \cdot) \text{ is a non decreasing function of } R^+ \text{ and } \lim_{\tau \rightarrow \infty} G(w, k, \tau) = 1;$$

$$(B7) \quad H(w, k, \tau) > 0;$$

$$(B8) \quad H(w, k, \tau) = 0 \Leftrightarrow w = k;$$

$$(B9) \quad H(w, k, \tau) = H(k, w, \tau);$$

$$(B10) \quad H(w, e, \theta(\tau + \pi)) \leq H(w, k, \tau) \Delta H(k, e, \Pi);$$

$$(B11) \quad H(w, k, \cdot) \text{ is a non increasing function of } R^+ \text{ and } \lim_{\tau \rightarrow \infty} H(w, k, \tau) = 1;$$

Then $(\Xi, G, H, *, \Delta)$ is an IFBMS.

Definition 2.2. Assume $\Xi \neq \phi$. Let $\perp \in \Xi \times \Xi$ be a binary relation. Suppose there exists $w_0 \in \Xi$ such that $w_0 \perp w$ or $w \perp w_0$ for all $w \in \Xi$. Thus, Ξ is known as orthogonal set (OS) and denoted by (Ξ, \perp)

Definition 2.3. Assume that (Ξ, \perp) is an OS. A sequence $\{w_n\}$ for $n \in \mathbf{N}$ is known to be an O-sequence if $(\forall n, w_n \perp w_{n+1})$ or $(\forall n, w_{n+1} \perp w_n)$

3 Orthogonal Intuitionistic Fuzzy b-metric Spaces

Now, we establish the notion of OIFBMSs and derive several FP results with non-trivial examples.

Definition 3.1. $(\Xi, G, H, *, \Delta)$ is known to be an OIFBMS if Ξ is a (non empty) OS, $*$ is a CTN, Δ is a CTCN, and G, H are FSS on $\Xi \times \Xi \times (0, \infty)$ verifying the below conditions for a given real number $\theta \geq 1$:

- $(B_{\perp 1})$ $G(w, k, \tau) + H(w, k, \tau) \leq 1$ for all $w, k \in \Xi, \tau > 0$ such that $w \perp k$ and $k \perp w$;
- $(B_{\perp 2})$ $G(w, k, \tau) > 0$ for all $w, k \in \Xi, \tau > 0$ such that $w \perp k$ and $k \perp w$;
- $(B_{\perp 3})$ $G(w, k, \tau) = 1 \Leftrightarrow w = k$; for all $w, k \in \Xi, \tau > 0$ such that $w \perp k$ and $k \perp w$;
- $(B_{\perp 4})$ $G(w, k, \tau) = G(k, w, \tau)$ for all $w, k \in \Xi, \tau > 0$ such that $w \perp k$ and $k \perp w$;
- $(B_{\perp 5})$ $G(w, e, \theta(\tau + \pi)) \geq G(w, k, \tau) * G(k, e, \Pi)$ for all $w, k \in \Xi, \tau > 0$ such that $w \perp k$ and $k \perp w$;
- $(B_{\perp 6})$ $G(w, k, \cdot)$ is a non decreasing function of R^+ and $\lim_{\tau \rightarrow \infty} G(w, k, \tau) = 1$ for all $w, k \in \Xi, \tau > 0$ such that $w \perp k$ and $k \perp w$;
- $(B_{\perp 7})$ $H(w, k, \tau) > 0$ for all $w, k \in \Xi, \tau > 0$ such that $w \perp k$ and $k \perp w$;
- $(B_{\perp 8})$ $H(w, k, \tau) = 0 \Leftrightarrow w = k$ for all $w, k \in \Xi, \tau > 0$ such that $w \perp k$ and $k \perp w$;
- $(B_{\perp 9})$ $H(w, k, \tau) = H(k, w, \tau)$ for all $w, k \in \Xi, \tau > 0$ such that $w \perp k$ and $k \perp w$;
- $(B_{\perp 10})$ $H(w, e, \theta(\tau + \pi)) \leq H(w, k, \tau) \Delta H(k, e, \Pi)$ for all $w, k \in \Xi, \tau > 0$ such that $w \perp k$ and $k \perp w$;
- $(B_{\perp 11})$ $H(w, k, \cdot)$ is a non increasing function of R^+ and $\lim_{\tau \rightarrow \infty} H(w, k, \tau) = 1$ for all $w, k \in \Xi, \tau > 0$ such that $w \perp k$ and $k \perp w$;

Then $(\Xi, G, H, *, \Delta)$ is an IFBMS.

Example 3.2. Let $\Xi = R$ and define $\sigma * \theta = \sigma\theta, \sigma\Delta\theta = \min\{\sigma, \theta\}$ and \perp by $w \perp k$ iff $w + k \geq 0$. Let

$$G(w, k, \tau) = \begin{cases} 1 & \text{if } w = k, \\ \frac{\tau}{\tau + \max\{w, k\}^\alpha} & \text{otherwise.} \end{cases} \quad (1)$$

and

$$H(w, k, \tau) = \begin{cases} 0 & \text{if } w = k, \\ \frac{\max\{w, k\}^\alpha}{\tau + \max\{w, k\}^\alpha} & \text{otherwise.} \end{cases} \quad (2)$$

for all $w, k \in \Xi, \tau > 0$ with α belong to odd natural numbers.

Proof. $(B_{\perp 1}) - (B_{\perp 3}), (B_{\perp 5}) - (B_{\perp 9})$ and $(B_{\perp 11})$ are obvious. Here, we prove $(B_{\perp 4})$ and $(B_{\perp 10})$. $(B_{\perp 4})$: for a random number $\theta \geq 1$, one writes

$$\max\{w, e\}^\alpha \leq \theta[\max\{w, k\}^\alpha + \max\{k, e\}^\alpha]$$

Thus,

$$\tau\pi \max\{w, e\}^\alpha \leq \theta(\tau + \pi)\pi \max\{w, k\}^\alpha + \theta(\tau + \pi)\tau \max\{k, e\}^\alpha.$$

Consequently,

$$\tau\pi \max\{w, e\}^\alpha \leq \theta(\tau + \pi)\pi \max\{w, k\}^\alpha + \theta(\tau + \pi)\tau \max\{k, e\}^\alpha + \theta(\tau + \pi) \max\{k, e\}^\alpha.$$

Thus,

$$\tau\pi \max\{w, e\}^\alpha \leq \theta(\tau + \pi)[\pi \max\{w, k\}^\alpha + \tau \max\{k, e\}^\alpha + \max\{w, k\}^\alpha \max\{k, e\}^\alpha].$$

one write

$$\theta(\tau + \pi)\tau\pi + \tau\pi \max\{w, e\}^\alpha \leq \theta(\tau + \pi)\tau\pi + \theta(\tau + \pi)[\pi \max\{w, k\}^\alpha + \tau \max\{k, e\}^\alpha + \max\{w, k\}^\alpha \max\{k, e\}^\alpha].$$

Therefore,

$$\theta(\tau + \pi)\tau\pi + \tau\pi \max\{w, e\}^\alpha \leq \theta(\tau + \pi)[\tau\pi + \pi \max\{w, k\}^\alpha + \tau \max\{k, e\}^\alpha + \max\{w, k\}^\alpha \max\{k, e\}^\alpha].$$

That is,

$$\tau\pi[\theta(\tau + \pi) + \max\{w, e\}^\alpha] \leq \theta(\tau + \pi)[\tau + \max\{w, k\}^\alpha][\pi + \max\{k, e\}^\alpha]$$

Hence,

$$\frac{\theta(\tau + \pi)}{\theta(\tau + \pi) + \max\{w, e\}^\alpha} \geq \frac{\tau\pi}{[\tau + \max\{w, k\}^\alpha][\pi + \max\{k, e\}^\alpha]}.$$

$$\frac{\theta(\tau + \pi)}{\theta(\tau + \pi) + \max\{w, e\}^\alpha} \geq \frac{\tau}{\tau + \max\{w, k\}^\alpha}.$$

That is,

$$G(w, e, \theta(\tau + \pi)) \geq G(w, k, \tau) * G(k, e, \pi).$$

($B_{\perp 10}$): One writes

$$\max\{w, e\}^\alpha = \max\{w, e\}^\alpha \max \left\{ \frac{\max\{w, k\}^\alpha}{\max\{w, k\}^\alpha}, \frac{\max\{k, e\}^\alpha}{\max\{k, e\}^\alpha} \right\}.$$

Then

$$\max\{w, e\}^\alpha \leq [\theta(\tau + \pi) + \max\{w, e\}^\alpha] \max \left\{ \frac{\max\{w, k\}^\alpha}{\max\{w, k\}^\alpha}, \frac{\max\{k, e\}^\alpha}{\max\{k, e\}^\alpha} \right\}.$$

That is,

$$\frac{\max\{w, e\}^\alpha}{\theta(\tau + \pi) + \max\{w, e\}^\alpha} \leq \max \left\{ \frac{\max\{w, k\}^\alpha}{\tau + \max\{w, k\}^\alpha}, \frac{\max\{k, e\}^\alpha}{\pi + \max\{k, e\}^\alpha} \right\}.$$

Hence,

$$H(w, e, \theta(\tau + \pi)) \leq H(w, k, \tau)\Delta H(k, e, \pi).$$

Now, we show it's not an IFBM. Indeed, for $\pi = \tau = 1$, $w = -1$, $k = -\frac{1}{2}$ and $\alpha = 3$, (B4) and (B10) fail. \square

Example 3.3. Let $\Xi = \mathbb{R}$ and define $\sigma * \theta = \sigma\theta$, $\sigma\Delta\theta = \min\{\sigma, \theta\}$ and \perp by $w \perp k$ iff $w + k \geq 0$. Let

$$G(w, k, \tau) = \begin{cases} 1 & \text{if } w = k, \\ \left[e^{\frac{\max\{w, k\}^\alpha}{\tau}} \right]^{-1} & \text{otherwise.} \end{cases} \quad (3)$$

and

$$H(w, k, \tau) = \begin{cases} 0 & \text{if } w = k, \\ 1 - \left[e^{\frac{\max\{w, k\}^\alpha}{\tau}} \right]^{-1} & \text{otherwise.} \end{cases} \quad (4)$$

for all $w, k \in \Xi, \tau > 0$ with α belong to odd natural numbers.

Proof. $(B_{\perp 1}) - (B_{\perp 3}), (B_{\perp 5}) - (B_{\perp 9})$ and $(B_{\perp 11})$ are obvious. Here, we prove $(B_{\perp 4})$ and $(B_{\perp 10})$.
 $(B_{\perp 4})$: for a random number $\theta \geq 1$, one writes

$$\max\{w, e\}^\alpha \leq \theta [\max\{w, k\}^\alpha + \max\{k, e\}^\alpha].$$

Therefore,

$$\max\{w, e\}^\alpha \leq \theta \left[\frac{\tau + \pi}{\tau} \max\{w, k\}^\alpha + \frac{\tau + \pi}{\pi} \max\{k, e\}^\alpha \right]$$

Then

$$\frac{\max\{w, e\}^\alpha}{\theta(\tau + \pi)} \leq \frac{\max\{w, k\}^\alpha}{\tau} + \frac{\max\{k, e\}^\alpha}{\pi}$$

Since, e^w is an increasing function, one gets

$$e^{\frac{\max\{w, e\}^\alpha}{\theta(\tau + \pi)}} \leq e^{\frac{\max\{w, k\}^\alpha}{\tau}} \cdot e^{\frac{\max\{k, e\}^\alpha}{\pi}}.$$

That is

$$\left[e^{\frac{\max\{w, e\}^\alpha}{\theta(\tau + \pi)}} \right]^{-1} \geq \left[e^{\frac{\max\{w, k\}^\alpha}{\tau}} \right]^{-1} \cdot \left[e^{\frac{\max\{k, e\}^\alpha}{\pi}} \right]^{-1}.$$

Hence,

$$G(w, e, \theta(\tau + \pi)) \geq G(w, k, \tau) * G(k, e, \pi).$$

$(B_{\perp 10})$: For a random $\theta \geq 1$, we write

$$\frac{\max\{w, e\}^\alpha}{\theta(\tau + \pi)} \leq \max \left\{ \frac{\max\{w, k\}^\alpha}{\tau}, \frac{\max\{k, e\}^\alpha}{\pi} \right\}.$$

That is,

$$e^{\frac{\max\{w, e\}^\alpha}{\theta(\tau + \pi)}} \leq \max \left\{ e^{\frac{\max\{w, k\}^\alpha}{\tau}}, e^{\frac{\max\{k, e\}^\alpha}{\pi}} \right\}.$$

Then,

$$\left[e^{\frac{\max\{w, e\}^\alpha}{\theta(\tau + \pi)}} \right]^{-1} \geq \max \left\{ \left[e^{\frac{\max\{w, k\}^\alpha}{\tau}} \right]^{-1}, \left[e^{\frac{\max\{k, e\}^\alpha}{\pi}} \right]^{-1} \right\}.$$

That is,

$$1 - \left[e^{\frac{\max\{w, e\}^\alpha}{\theta(\tau + \pi)}} \right]^{-1} \leq \max \left\{ 1 - \left[e^{\frac{\max\{w, k\}^\alpha}{\tau}} \right]^{-1}, 1 - \left[e^{\frac{\max\{k, e\}^\alpha}{\pi}} \right]^{-1} \right\}.$$

Hence,

$$H(w, e, \theta(\tau + \pi)) \leq H(w, k, \tau) \Delta H(k, e, \pi). \forall w, k, e \in \Xi, \forall \tau, \pi > 0.$$

Now, we show it's not an IFBM. Indeed, for $\pi = \tau = 1, w = -1, k = -\frac{1}{2}, e = -2$ and $\alpha = 3$, (B_4) and (B_{10}) is not satisfy. \square

Example 3.4. Let $\Xi = \mathbb{R}$ and define $\sigma * \theta = \sigma\theta$, $\sigma\Delta\theta = \max\{\sigma, \theta\}$ and \perp by $w \perp k$ iff $w + k \geq 0$. Suppose

$$G(w, k, \tau) = \frac{\tau + \min\{w, k\}^\alpha}{\tau + \max\{w, k\}^\alpha} \quad (5)$$

and

$$H(w, k, \tau) = 1 - \frac{\tau + \min\{w, k\}^\alpha}{\tau + \max\{w, k\}^\alpha} \quad (6)$$

for all $w, k \in \Xi, \tau > 0$ with α belong to odd natural numbers. Here, $(\Xi, G, H, *, \Delta, \perp)$ is an OIFBMS. It is not an IFBMS. Indeed, if it is the case, from (B4),

$$\frac{\theta(\tau + \pi) + \min\{w, k\}^\alpha}{\theta(\tau + \pi) + \max\{w, k\}^\alpha} \geq \frac{\tau + \min\{w, k\}^\alpha}{\tau + \max\{w, k\}^\alpha} \cdot \frac{\pi + \min\{w, k\}^\alpha}{\pi + \max\{w, k\}^\alpha}$$

and from case (B10)

$$1 - \frac{\theta(\tau + \pi) + \min\{w, k\}^\alpha}{\theta(\tau + \pi) + \max\{w, k\}^\alpha} \leq \max \left[1 - \frac{\tau + \min\{w, k\}^\alpha}{\tau + \max\{w, k\}^\alpha} \cdot 1 - \frac{\pi + \min\{w, k\}^\alpha}{\pi + \max\{w, k\}^\alpha} \right].$$

Then by taking $w = k, e = -2$ and $\alpha = \frac{1}{2}$, the above inequalities are not satisfied.

Remark 3.5. Every IFBMS is an OIFBMS, but the converse is not true. The above examples confirm this reverse statement.

Definition 3.6. An O-sequence $\{w_n\}$ is an OIFBMS $(\Xi, G, H, *, \Delta, \perp)$ is called an orthogonal convergent (O-convergent) to $w \in \Xi$, if

$$\lim_{n \rightarrow \infty} G(w_n, w, \tau) = 1, \forall \tau > 0,$$

and

$$\lim_{n \rightarrow \infty} H(w_n, w, \tau) = 0, \forall \tau > 0,$$

Definition 3.7. An O-sequence $\{w_n\}$ is an OIFBMS $(\Xi, G, H, *, \Delta, \perp)$ is known to be an orthogonal Cauchy (O-Cauchy) if

$$\lim_{n \rightarrow \infty} G(w_n, w, \tau) = 1,$$

and

$$\lim_{n \rightarrow \infty} H(w_n, w, \tau) = 0,$$

for all $\tau > 0, p \geq 1$.

Definition 3.8. Let $\xi : \Xi \rightarrow \Xi$ is \perp -continuous at $w \in \Xi$ is an OIFBMS $(\Xi, G, H, *, \Delta, \perp)$, whenever for each O-sequence w_n for all $n \in \mathbb{N}$ in Ξ if $\lim_{n \rightarrow \infty} G(w_n, w, \tau) = 1$ and $\lim_{n \rightarrow \infty} H(w_n, w, \tau) = 0$ for all $\tau > 0$, then $\lim_{n \rightarrow \infty} G(\xi w_n, \xi w, \tau) = 1$ and $\lim_{n \rightarrow \infty} H(\xi w_n, \xi w, \tau) = 0$ for all $\tau > 0$. Furthermore, ξ is \perp -continuous on Ξ if $\xi \perp$ -continuous at each $w \in \Xi$. Also, ξ is \perp -preserving if $\xi w \perp \xi k$, whence $w \perp k$.

Definition 3.9. An OIFBMS $(\Xi, G, H, *, \Delta, \perp)$ is known to be orthogonally complete (O-complete) if every O-Cauchy O-sequence is O-convergent.

Remark 3.10. It is necessary that the limit of an O-convergent O-sequence is unique in an OIFBMS.

Remark 3.11. It is necessary that the limit of an O-convergent O-sequence is O-Cauchy in an OIFBMS.

Lemma 3.12. *If for some $v \in (0, 1)$ and $w, k \in \Xi$,*

$$G(w, k, \tau) \geq G\left(w, k, \frac{\tau}{v}\right), \tau > 0,$$

and

$$H(w, k, \tau) \leq H\left(w, k, \frac{\tau}{v}\right), \tau > 0,$$

then $w = k$.

Proof. *The proof is follows from [8].* \square

Definition 3.13. Suppose $(\Xi, G, H, *, \Delta, \perp)$ be an OIFBMS. A mapping $\xi : \Xi \rightarrow \Xi$ is an orthogonal contraction (\perp -contraction) if there exists $\varrho \in (0, 1)$ such that for every $\tau > 0$ and $w, k \in \Xi$ with $w \perp k$, we have

$$G(\xi w, \xi k, \varrho\tau) \geq G(w, k, \tau), \tag{7}$$

$$H(\xi w, \xi k, \varrho\tau) \leq H(w, k, \tau). \tag{8}$$

Theorem 3.14. *Let $(\Xi, G, H, *, \Delta, \perp)$ be an O-complete IFBMS such that*

$$\lim_{\tau \rightarrow \infty} G(w, k, \tau) = 1,$$

and

$$\lim_{\tau \rightarrow \infty} H(w, k, \tau) = 0.$$

for all $w, k \in \Xi$. Suppose $\xi : \Xi \rightarrow \Xi$ be an \perp -continuous and \perp -preserving mapping. Thus, ξ has a unique FP, say $w_* \in \Xi$. Furthermore,

$$\lim_{\tau \rightarrow \infty} G(\xi^n w, k, \tau) = 1,$$

and

$$\lim_{\tau \rightarrow \infty} H(\xi^n w, k, \tau) = 0.$$

for all $w, k \in \Xi$.

Proof. Let $(\Xi, G, H, *, \Delta, \perp)$ be an O-complete IFBMS, there exists $w_0 \in \Xi$ such that $w_0 \perp k$ for all $k \in \Xi$, that is, $w_0 \perp \xi w_0$. Take $w_n = \xi^n w_0 = \xi w_{n-1}$ for all $n \in \mathbb{N}$. Since ξ is \perp -preserving, $\{w_n\}$ is an O-sequence. From assumption that ξ is an \perp -contraction, we have

$$G(w_{n+1}, w_n, \varrho\tau) = G(\xi w_n, \xi w_{n-1}, \varrho\tau) \geq G(w_n, w_{n-1}, \tau)$$

for all $n \in \mathbb{N}$ and $\tau > 0$. Note that G is non-decreasing on $(0, \infty)$. By utilizing above inequality, we have

$$\begin{aligned} G(w_{n+1}, w_n, \tau) &\geq G(w_{n+1}, w_n, \varrho\tau) = G(\xi w_{n+1}, \xi w_n, \varrho\tau) \geq G(w_n, w_{n-1}, \tau) \\ &= G(\xi w_{n-1}, \xi w_{n-2}, \tau) \geq G\left(w_{n-1}, w_{n-2}, \frac{\tau}{\varrho}\right) \geq \dots \geq G\left(w_1, w_0, \frac{\tau}{\varrho^n}\right) \end{aligned} \tag{9}$$

for all $n \in \mathbb{N}$ and $\tau > 0$. Thus, from (9) and (B4), we deduce

$$\begin{aligned} G(w_n, w_{n+m}, \tau) &\geq G\left(w_n, w_{n+1}, \frac{\tau}{\theta}\right) * G\left(w_{n+1}, w_{n+m}, \frac{\tau}{\theta}\right) \\ &\geq G\left(w_n, w_{n+1}, \frac{\tau}{\theta}\right) * G\left(w_{n+1}, w_{n+2}, \frac{\tau}{\theta^2}\right) * G\left(w_{n+2}, w_{n+3}, \frac{\tau}{\theta^3}\right) * \dots * G\left(w_{n+m-1}, w_{n+m}, \frac{\tau}{\theta^{n+m}}\right) \end{aligned}$$

$$\geq G\left(w_1, w_0, \frac{\tau}{\theta \rho^n}\right) * G\left(w_{n+1}, w_{n+2}, \frac{\tau}{\theta^2 \rho^n}\right) * G\left(w_{n+2}, w_{n+3}, \frac{\tau}{\theta^3 \rho^n}\right) * \cdots * G\left(w_{n+m-1}, w_{n+m}, \frac{\tau}{\theta^{n+m} \rho^n}\right) \quad (10)$$

We know that $\lim_{\tau \rightarrow \infty} G(w, k, \tau) = 1$, for all $w, k \in \Xi$ and $\tau > 0$. So, from (10), we have

$$\lim_{\tau \rightarrow \infty} G(w_n, w_{n+m}, \tau) \geq 1 * 1 * \cdots * 1 = 1. \quad (11)$$

Similarly,

$$H(w_{n+1}, w_n, \rho\tau) = H(\xi w_n, \xi w_{n-1}, \rho\tau) \leq H(w_n, w_{n-1}, \tau)$$

for all $n \in \mathbb{N}$ and $\tau > 0$. By utilizing above inequality, we have

$$\begin{aligned} H(w_{n+1}, w_n, \tau) &\leq H(w_{n+1}, w_n, \rho\tau) = H(\xi w_{n+1}, \xi w_n, \rho\tau) \leq H(w_n, w_{n-1}, \tau) \\ &= H(\xi w_{n-1}, \xi w_{n-2}, \tau) \leq H\left(w_{n-1}, w_{n-2}, \frac{\tau}{\rho}\right) \leq \cdots \leq H\left(w_1, w_0, \frac{\tau}{\rho^n}\right) \end{aligned} \quad (12)$$

for all $n \in \mathbb{N}$ and $\tau > 0$. Thus, from (12) and (B10), we deduce

$$\begin{aligned} H(w_n, w_{n+m}, \tau) &\leq H\left(w_n, w_{n+1}, \frac{\tau}{\theta}\right) \Delta H\left(w_{n+1}, w_{n+m}, \frac{\tau}{\theta}\right) \\ &\leq H\left(w_n, w_{n+1}, \frac{\tau}{\theta}\right) \Delta H\left(w_{n+1}, w_{n+2}, \frac{\tau}{\theta^2}\right) \Delta H\left(w_{n+2}, w_{n+3}, \frac{\tau}{\theta^3}\right) \Delta \cdots \Delta H\left(w_{n+m-1}, w_{n+m}, \frac{\tau}{\theta^{n+m}}\right) \\ &\leq H\left(w_1, w_0, \frac{\tau}{\theta \rho^n}\right) \Delta H\left(w_{n+1}, w_{n+2}, \frac{\tau}{\theta^2 \rho^n}\right) \Delta H\left(w_{n+2}, w_{n+3}, \frac{\tau}{\theta^3 \rho^n}\right) \Delta \cdots \Delta H\left(w_{n+m-1}, w_{n+m}, \frac{\tau}{\theta^{n+m} \rho^n}\right) \end{aligned} \quad (13)$$

We know that $\lim_{\tau \rightarrow \infty} H(w, k, \tau) = 0$, for all $w, k \in \Xi$ and $\tau > 0$. So, from (13), we have

$$\lim_{\tau \rightarrow \infty} H(w_n, w_{n+m}, \tau) \leq 0 \Delta 0 \Delta \cdots \Delta 0 = 0. \quad (14)$$

So, $\{w_n\}$ is an O -sequence. The O -sequence. The O -completeness of the IFBMS $(\Xi, w, k, *, \Delta, \perp)$ ensure that there exists $w_* \in \Xi$ such that $G(w_n, w_*, \tau) \rightarrow 1$, and $H(w_n, w_*, \tau) \rightarrow 0$, as $n \rightarrow +\infty$ for all $\tau > 0$. Now, since ξ is an \perp -continuous mapping, $G(w_{n+1}, \xi w_*, \tau) = G(\xi w_{n+1}, \xi w_*, \tau) \rightarrow 1$ and $H(w_{n+1}, \xi w_*, \tau) = H(\xi w_{n+1}, \xi w_*, \tau) \rightarrow 0$ as $n \rightarrow +\infty$. Now, we have

$$\begin{aligned} G(w_*, \xi w_*, \tau) &\geq G\left(w_*, w_{n+1}, \frac{\tau}{2\theta}\right) * G\left(w_{n+1}, \xi w_*, \frac{\tau}{2\theta}\right), \\ H(w_*, \xi w_*, \tau) &\leq H\left(w_*, w_{n+1}, \frac{\tau}{2\theta}\right) \Delta H\left(w_{n+1}, \xi w_*, \frac{\tau}{2\theta}\right). \end{aligned}$$

Taking limit as $n \rightarrow \infty$, we get $G(w_*, \xi w_*, \tau) = 1 * 1 = 1$ and $H(w_*, \xi w_*, \tau) = 0 \Delta 0 = 0$ and hence $\xi w_* = w_*$.

Uniqueness:

Let w_* and k_* be two FPs of ξ such that $w_* \neq k_*$. We have $w_0 \perp w_*$ and $w_0 \perp k_*$. Since T is \perp -preserving, we have $\xi w_0 \perp \xi^n w_*$ and $\xi^n w_0 \perp k_*$ for all $n \in \mathbb{N}$. So from (7), we can drive

$$G(\xi^n w_0, \xi^n w_*, \tau) \geq G(\xi^n w_0, \xi^n w_*, \rho\tau) \geq G\left(w_0, w_*, \frac{\tau}{\rho^n}\right)$$

and

$$G(\xi^n w_0, \xi^n k_*, \tau) \geq G(\xi^n w_0, \xi^n k_*, \rho\tau) \geq G\left(w_0, k_*, \frac{\tau}{\rho^n}\right)$$

Therefore,

$$\begin{aligned} G(w_*, k_*, \tau) &= G(\xi^n w_*, \xi^n k_*, \tau) \geq G\left(\xi^n w_0, \xi^n w_*, \frac{\tau}{2\theta}\right) * G\left(\xi^n w_0, \xi^n k_*, \frac{\tau}{2\theta}\right) \\ &\geq G\left(w_0, w_*, \frac{\tau}{2\theta \rho^n}\right) * G\left(w_0, k_*, \frac{\tau}{2\theta \rho^n}\right) \rightarrow 1 \end{aligned}$$

as $n \rightarrow \infty$ So from (8), we can derive

$$H(\xi^n w_0, \xi^n w_*, \tau) \leq H(\xi^n w_0, \xi^n w_*, \rho\tau) \leq H\left(w_0, w_*, \frac{\tau}{\rho^n}\right)$$

and

$$H(\xi^n w_0, \xi^n k_*, \tau) \leq H(\xi^n w_0, \xi^n k_*, \rho\tau) \leq H\left(w_0, k_*, \frac{\tau}{\rho^n}\right)$$

Therefore,

$$\begin{aligned} H(w_*, k_*, \tau) &= H(\xi^n w_*, \xi^n k_*, \tau) \leq H\left(\xi^n w_0, \xi^n w_*, \frac{\tau}{2\theta}\right) * H\left(\xi^n w_0, \xi^n k_*, \frac{\tau}{2\theta}\right) \\ &\leq H\left(w_0, w_*, \frac{\tau}{2\theta \rho^n}\right) \Delta H\left(w_0, k_*, \frac{\tau}{2\theta \rho^n}\right) \rightarrow 0 \end{aligned}$$

as $n \rightarrow \infty$ So, $w_* = k_*$, hence w_* is the unique FP. \square

Corollary 3.15. Suppose $(\Xi, G, H, *, \Delta, \perp)$ be an O -complete IFBMS. Assume $\xi : \Xi \rightarrow \Xi$ be \perp -contraction and \perp -preserving. Assume that if $\{w_n\}$ is an O -sequence with $w_n \rightarrow w \in \Xi$, Then $w \perp w_n$ for all $n \in \mathbb{N}$. Then ξ has a unique FP, say $w_* \in \Xi$, Moreover, $\lim_{n \rightarrow \infty} G(\xi^n w, w_*, \tau) = 1$ and $\lim_{n \rightarrow \infty} H(\xi^n w, w_*, \tau) = 0$, for all $w \in \Xi$ and $\tau > 0$.

Proof. Follows from Theorem 2.1 that w_n is a O -Cauchy O -sequence and so it O -converges to $w_* \in \Xi$. Hence $w_* \perp w_n$ for all $n \in \mathbb{N}$ from (7), we have

$$G(\xi w_*, w_{n+1}, \tau) = G(\xi w_*, \xi w_n, \tau) \geq G(\xi w_*, \xi w_n, \tau \rho) \geq G(w_*, w_n, \tau)$$

and

$$\lim_{n \rightarrow \infty} G(\xi w_*, w_{n+1}, \tau) = 1.$$

Then, we can write

$$G(w_*, \xi w_*, \tau) \geq G\left(w_*, \xi w_{n+1}, \frac{\tau}{2\theta}\right) * G\left(w_{n+1}, \xi w_*, \frac{\tau}{2\theta}\right)$$

Taking limit as $n \rightarrow +\infty$, We get $G(w_*, \xi w_*, \tau) = 1 * 1 = 1$

and from (8)

$$H(\xi w_*, w_{n+1}, \tau) = H(\xi w_*, \xi w_n, \tau) \leq H(\xi w_*, \xi w_n, \tau \rho) \leq H(w_*, w_n, \tau)$$

and

$$\lim_{n \rightarrow \infty} H(\xi w_*, w_{n+1}, \tau) = 0.$$

Then, we can write

$$H(w_*, \xi w_*, \tau) \leq H\left(w_*, \xi w_{n+1}, \frac{\tau}{2\theta}\right) \Delta H\left(w_{n+1}, \xi w_*, \frac{\tau}{2\theta}\right)$$

Taking limit as $n \rightarrow +\infty$, We get $H(w_*, \xi w_*, \tau) = 0 \Delta 0 = 0$, So $\xi w_* = w_*$. Next follows from Theorem 3.13.

\square

Example 3.16. Let $\Xi = [-2, 2]$. We define \perp by

$$w \perp k \Leftrightarrow w + k \in \{|w|, |k|\} \quad (15)$$

$$G(w, k, \tau) = \begin{cases} 1 & \text{if } w = k, \\ \left[e^{\frac{\max\{w, k\}^\alpha}{\tau}} \right]^{-1} & \text{otherwise.} \end{cases} \quad (16)$$

and

$$H(w, k, \tau) = \begin{cases} 0 & \text{if } w = k, \\ 1 - \left[e^{\frac{\max\{w, k\}^\alpha}{\tau}} \right]^{-1} & \text{otherwise.} \end{cases} \quad (17)$$

for all $w, k \in \Xi, \tau > 0$ with $\sigma \times \theta = \sigma \cdot \theta$ and $\sigma \Delta \theta = \max\{\sigma, \theta\}$. Then $(\Xi, G, *, \Delta, \perp)$ is an O-complete IFBMS. Define $\xi : \Xi \rightarrow \Xi$ by

$$\xi(w) = \begin{cases} \frac{w}{4}, & \text{if } w \in [-2, 0] \\ 0, & \text{if } w \in (0, 2]. \end{cases} \quad (18)$$

Then the below cases fulfilled:

1. if $w \in [-2, 0]$ and $k \in (0, 2]$, then $\xi(w) = \frac{w}{4}$ and $\xi(k) = 0$,
2. if $w, k \in [-2, 0]$, then $\xi(w) = \frac{w}{4}$ and $\xi(k) = \frac{k}{4}$,
3. if $w, k \in (0, 2]$, then $\xi(w) = 0$ and $\xi(k) = 0$,
4. if $w \in (0, 2]$ and $k \in [-2, 0]$, then $\xi(w) = 0$ and $\xi(k) = \frac{k}{4}$,

This is easy to see that $\xi((w)) + \xi(k) \in \{|\xi(w)|, |\xi(k)|\}$. Hence, ξ is \perp -preserving. Let $\{w_n\}$ be an arbitrary O-sequence in Ξ that $\{w_n\}$ O-converges to $w \in \Xi$. That is

$$\lim_{n \rightarrow \infty} G(w_n, w, \tau) = \lim_{n \rightarrow \infty} \left[e^{\frac{\max\{w_n, w\}^\alpha}{\tau}} \right]^{-1} = 1,$$

$$\lim_{n \rightarrow \infty} H(w_n, w, \tau) = 1 - \lim_{n \rightarrow \infty} \left[e^{\frac{\max\{w_n, w\}^\alpha}{\tau}} \right]^{-1} = 0.$$

We can easily see that if $\lim_{n \rightarrow \infty} G(w_n, w, \tau) = 1$, then $\lim_{n \rightarrow \infty} G(\xi w_n, \xi w, \tau) = 1$, and if $\lim_{n \rightarrow \infty} H(w_n, w, \tau) = 0$, then $\lim_{n \rightarrow \infty} H(\xi w_n, \xi w, \tau) = 0$, for all $w \in \Xi$ and $\tau > 0$. That is, ξ is \perp -continuous. if $w = k$, then it is obvious. Suppose $w \neq k$, then there are following four cases for $\varrho \in [\frac{1}{2}, 1)$:

Case 1) if $w \in [-2, 0]$ and $k \in (0, 2]$, then $\xi w = \frac{w}{4}$ and $\xi k = 0$. Here,

$$G(\xi w_n, \xi w, \varrho \tau) = G\left(\frac{w}{4}, 0, \varrho \tau\right) = \left[e^{\frac{[\frac{w}{4}]^\alpha}{\varrho \tau}} \right]^{-1} \geq \left[e^{\frac{\max\{w, k\}^\alpha}{\tau}} \right]^{-1} = G(w, k, \tau),$$

$$H(\xi w_n, \xi w, \varrho \tau) = H\left(\frac{w}{4}, 0, \varrho \tau\right) = 1 - \left[e^{\frac{[\frac{w}{4}]^\alpha}{\varrho \tau}} \right]^{-1} \leq 1 - \left[e^{\frac{\max\{w, k\}^\alpha}{\tau}} \right]^{-1} = H(w, k, \tau),$$

Case 2) If $w, k \in [-2, 0]$, then $\xi w = \frac{w}{4}$ and $\xi k = \frac{k}{4}$. We have

$$G(\xi w_n, \xi w, \varrho \tau) = G\left(\frac{w}{4}, \frac{k}{4}, \varrho \tau\right) = \left[e^{\frac{\max\{\frac{w}{4}, \frac{k}{4}\}^\alpha}{\varrho \tau}} \right]^{-1} \geq \left[e^{\frac{\max\{w, k\}^\alpha}{\tau}} \right]^{-1} = G(w, k, \tau),$$

$$H(\xi w_n, \xi w, \varrho\tau) = H\left(\frac{w}{4}, \frac{k}{4}, \varrho\tau\right) = 1 - \left[e^{\frac{\max\{\frac{w}{4}, \frac{k}{4}\}^\alpha}{\varrho\tau}} \right]^{-1} \leq 1 - \left[e^{\frac{\max\{w, k\}^\alpha}{\tau}} \right]^{-1} = H(w, k, \tau),$$

Case 3) If $w, k \in (0, 2]$, then $\xi w = 0$ and $\xi k = 0$. Here,

$$G(\xi w, \xi k, \varrho\tau) = G(0, 0, \varrho\tau) = e^0 \geq \left[e^{\frac{\max\{w, k\}^\alpha}{\tau}} \right]^{-1} = G(w, k, \tau),$$

$$H(\xi w, \xi k, \varrho\tau) = H(0, 0, \varrho\tau) = 1 - e^0 \leq 1 - \left[e^{\frac{\max\{w, k\}^\alpha}{\tau}} \right]^{-1} = H(w, k, \tau),$$

Case 4) If $w \in (0, 2]$ and $k \in [-2, 0]$, then $\xi w = 0$ and $\xi k = \frac{k}{4}$. We have

$$G(\xi w, \xi k, \varrho\tau) = G\left(0, \frac{k}{4}, \varrho\tau\right) = \left[e^{\frac{\max\{0, \frac{k}{4}\}^\alpha}{\varrho\tau}} \right]^{-1} \geq \left[e^{\frac{\max\{w, k\}^\alpha}{\tau}} \right]^{-1} = G(w, k, \tau),$$

$$H(\xi w, \xi k, \varrho\tau) = H\left(0, \frac{k}{4}, \varrho\tau\right) = 1 - \left[e^{\frac{\max\{0, \frac{k}{4}\}^\alpha}{\varrho\tau}} \right]^{-1} \leq 1 - \left[e^{\frac{\max\{w, k\}^\alpha}{\tau}} \right]^{-1} = H(w, k, \tau),$$

From all the above cases, We obtain that

$$G(\xi w, \xi k, \varrho\tau) \geq G(w, k, \tau), \tag{19}$$

$$H(\xi w, \xi k, \varrho\tau) \geq H(w, k, \tau), \tag{20}$$

Hence, ξ is an orthogonal contraction. But, ξ is not a contraction. In fact, let $w = -1$ and $k = -2$ and $\alpha = 3$, then

$$G(\xi w, \xi k, \varrho\tau) = \left[e^{\frac{\max\{\frac{w}{4}, \frac{k}{4}\}^\alpha}{\varrho\tau}} \right]^{-1} \geq 1,$$

$$H(\xi w, \xi k, \varrho\tau) = 1 - \left[e^{\frac{\max\{\frac{w}{4}, \frac{k}{4}\}^\alpha}{\varrho\tau}} \right]^{-1} \leq 0.$$

Which is not true. Hence, all assumptions of Theorem 3.13 are fulfilled and 0 is the unique FP of ξ . Also,

$$G(w, w, \tau) = G(0, 0, \tau) = e^0 = 1, \forall \tau > 0 \tag{21}$$

and

$$H(w, w, \tau) = H(0, 0, \tau) = 1 - e^0 = 0, \forall \tau > 0 \tag{22}$$

Theorem 3.17. Suppose $(\Xi, G, H, *, \Delta, \perp)$ be an O-complete IFBMS such that $\lim_{t \rightarrow \infty} G(w, k, \tau) = 1$, and $\lim_{t \rightarrow \infty} H(w, k, \tau) = 0, \forall w, k \in \Xi$ and $\tau > 0$. Suppose $\xi : \Xi \rightarrow \Xi$ be \perp -continuous, \perp -contraction, and \perp -preserving. Suppose $\varrho \in (0, \frac{1}{\varrho})$ and $\tau > 0$, such that

$$G(\xi w, \xi k, \varrho\tau) \geq \min\{G(\xi w, w, \tau), G(\xi k, k, \tau)\} \tag{23}$$

$$H(\xi w, \xi k, \varrho\tau) \leq \min\{H(\xi w, w, \tau), H(\xi k, k, \tau)\} \tag{24}$$

for all $w, k \in \Xi, \tau > 0$. Then ξ has a unique FP, say $w_* \in \Xi$. Moreover, $\lim_{n \rightarrow \infty} G(\xi^n w, w_*, \tau) = 1$ and $\lim_{n \rightarrow \infty} H(\xi^n w, w_*, \tau) = 0$ for all $w \in \Xi$ and $\tau > 0$.

Proof. Let $(\Xi, G, H, *, \Delta, \perp)$ be an O-complete IFBMS, There exists $w_0 \in \Xi$ such that

$$w_0 \perp k \forall k \in \Xi \tag{25}$$

Therefore, ξ is \perp -preserving, and $\{w_n\}$ is an O -sequence. We have

$$G(w_{n+1}, n, \tau) \geq G(w_{n+1}, n, \varrho\tau) = G(\xi w_n, \xi w_{n-1}, \varrho\tau) \geq \min\{G(\xi w_n, n, \tau), G(\xi w_{n-1}, w_{n-1}, \tau)\}$$

$$H(w_{n+1}, n, \tau) \leq H(w_{n+1}, n, \varrho\tau) = H(\xi w_n, \xi w_{n-1}, \varrho\tau) \leq \min\{H(\xi w_n, n, \tau), H(\xi w_{n-1}, w_{n-1}, \tau)\}$$

Two cases arise.

Case 1: If $G(w_{n+1}, n, \tau) \geq G(\xi w_n, w_n, \tau)$, then

$$G(w_{n+1}, w_n, \tau) \geq G(w_{n+1}, w_n, \varrho\tau) \geq G(\xi w_n, w_n, \tau) = G(w_{n+1}, w_n, \tau)$$

and

$$H(w_{n+1}, w_n, \tau) \leq H(w_{n+1}, w_n, \varrho\tau) \leq H(\xi w_n, w_n, \tau) = H(w_{n+1}, w_n, \tau)$$

Then, by Lemma 3.12, $w_n = w_{n+1}$ for all $n \in \mathbb{N}$

Case 2): If $G(w_{n+1}, n, \tau) \geq G(\xi w_{n-1}, w_{n-1}, \tau)$, then

$$G(w_{n+1}, w_n, \tau) \geq G(w_{n+1}, w_n, \varrho\tau) \geq G(\xi w_{n-1}, w_{n-1}, \tau) \geq G(w_n, w_{n-1}, \tau)$$

and $H(w_{n+1}, n, \tau) \leq H(\xi w_{n-1}, w_{n-1}, \tau)$, then

$$H(w_{n+1}, w_n, \tau) \leq H(w_{n+1}, w_n, \varrho\tau) \leq H(\xi w_{n-1}, w_{n-1}, \tau) \leq H(w_n, w_{n-1}, \tau)$$

for all $n \in \mathbb{N}$ and $\tau > 0$. By utilizing Theorem 3.13, we have an O -Cauchy sequence. Since $(\Xi, G, H, *, \Delta, \perp)$ is complete, there exists $w_* \in \Xi$, such that

$$\lim_{n \rightarrow \infty} G(w_n, w_*, \tau) = 1, \quad (26)$$

and

$$\lim_{n \rightarrow \infty} H(w_n, w_*, \tau) = 0, \quad (27)$$

for all $\tau > 0$. Science, ξ is an \perp -continuous, We have

$$\lim_{n \rightarrow \infty} G(w_{n+1}, w_*, \tau) = G(\xi w_n, \xi w_*, \tau) = 1,$$

and

$$\lim_{n \rightarrow \infty} H(w_{n+1}, w_*, \tau) = H(\xi w_n, \xi w_*, \tau) = 0,$$

Next, we examine that w_* is a FP of ξ . Let $\tau_1 \in (\varrho\theta, 1)$ and $\tau_2 = 1 - \tau_1$. then

$$\begin{aligned} G(\xi w_*, w_*, \tau) &\geq G\left(\xi w_*, w_{n+1}, \frac{\tau\tau_1}{\theta}\right) * G\left(\xi w_{n+1}, w_*, \frac{\tau\tau_2}{\theta}\right), \\ &= G\left(\xi w_*, \xi w_n, \frac{\tau\tau_1}{\theta}\right) * G\left(\xi w_{n+1}, w_*, \frac{\tau\tau_2}{\theta}\right), \\ &\geq \min\left\{G\left(\xi w_*, w_*, \frac{\tau\tau_1}{\theta}\right) * G\left(\xi w_n, w_n, \frac{\tau\tau_2}{\theta}\right)\right\} * G\left(\xi w_{n+1}, w_*, \frac{\tau\tau_2}{\theta}\right) \\ &= \min\left\{G\left(\xi w_*, w_*, \frac{\tau\tau_1}{\theta}\right) * G\left(\xi w_{n+1}, w_n, \frac{\tau\tau_2}{\theta}\right)\right\} * G\left(\xi w_{n+1}, w_*, \frac{\tau\tau_2}{\theta}\right) \end{aligned}$$

Taking $n \rightarrow \infty$, We get

$$G(\xi w_*, w_*, \tau) \geq \min\left\{G\left(\xi w_*, w_*, \frac{\tau\tau_1}{\theta}\right), 1\right\} * 1,$$

$$G(\xi w_*, w_*, \tau) \geq G\left(\xi w_*, w_*, \frac{\tau}{\nu}\right) \tau > 0, ,$$

and

$$\begin{aligned} H(\xi w_*, w_*, \tau) &\leq H\left(\xi w_*, w_{n+1}, \frac{\tau\tau_1}{\theta}\right) \Delta H\left(\xi w_{n+1}, w_*, \frac{\tau\tau_2}{\theta}\right), \\ &= H\left(\xi w_*, \xi w_n, \frac{\tau\tau_1}{\theta}\right) \Delta H\left(\xi w_{n+1}, w_*, \frac{\tau\tau_2}{\theta}\right), \\ &\leq \min\left\{H\left(\xi w_*, w_*, \frac{\tau\tau_1}{\theta}\right) \Delta H\left(\xi w_n, w_n, \frac{\tau\tau_2}{\theta}\right)\right\} \Delta H\left(\xi w_{n+1}, w_*, \frac{\tau\tau_2}{\theta}\right) \\ &= \min\left\{H\left(\xi w_*, w_*, \frac{\tau\tau_1}{\theta}\right) \Delta H\left(\xi w_{n+1}, w_n, \frac{\tau\tau_2}{\theta}\right)\right\} \Delta H\left(\xi w_{n+1}, w_*, \frac{\tau\tau_2}{\theta}\right) \end{aligned}$$

Taking $n \rightarrow \infty$, We get

$$\begin{aligned} H(\xi w_*, w_*, \tau) &\leq \min\left\{H\left(\xi w_*, w_*, \frac{\tau\tau_1}{\theta}\right), 0\right\} * 0, \\ H(\xi w_*, w_*, \tau) &\leq H\left(\xi w_*, w_*, \frac{\tau}{\nu}\right) \tau > 0, , \end{aligned}$$

There is $\nu = \frac{\theta\rho}{\tau_1} \in (0, 1)$, and by utilizing Lemma 3.12, we get $\xi w_* = w_*$.

Uniqueness : Suppose $w_* \neq k_*$ are two FPs of ξ . We get $w_0 \perp w_*$ and $w_0 \perp k_*$. Therefore, since ξ is an \perp -preserving, we have $\xi^n w_0 \perp \xi^n w_*$ and $\xi^n w_0 \xi^n \perp k_*$. for all $n \in \mathbb{N}$. we can write

$$G(\xi^n w_0, \xi^n w_*, \tau) \geq G(\xi^n w_0, \xi^n w_*, \rho\tau) \geq \min\{G(\xi^n w_0, w_0, \tau), G(\xi^n w_*, w_*, \tau)\},$$

and

$$G(\xi^n w_0, \xi^n k_*, \tau) \geq G(\xi^n w_0, \xi^n k_*, \rho\tau) \geq \min\{G(\xi^n w_0, w_0, \tau), G(\xi^n k_*, k_*, \tau)\},$$

Hence, we write that

$$G(w_0, k_*, \tau) = G(\xi^n w_*, \xi^n k_*, \tau) \geq \min\left\{G\left(\xi^n w_*, w_*, \frac{\tau}{\rho}\right), G\left(\xi^n k_*, k_*, \frac{\tau}{\rho}\right)\right\},$$

and

$$H(\xi^n w_0, \xi^n w_*, \tau) \leq G(\xi^n w_0, \xi^n w_*, \rho\tau) \leq \min\{H(\xi^n w_0, w_0, \tau), H(\xi^n w_*, w_*, \tau)\},$$

and

$$H(\xi^n w_0, \xi^n k_*, \tau) \leq H(\xi^n w_0, \xi^n k_*, \rho\tau) \leq \min\{H(\xi^n w_0, w_0, \tau), H(\xi^n k_*, k_*, \tau)\},$$

Hence, we write that

$$H(w_0, k_*, \tau) = H(\xi^n w_*, \xi^n k_*, \tau) \leq \min\left\{H\left(\xi^n w_*, w_*, \frac{\tau}{\rho}\right), H\left(\xi^n k_*, k_*, \frac{\tau}{\rho}\right)\right\},$$

for all $\tau > 0$. Thus, $w_* = k_*$. \square

Corollary 3.18. Suppose $(\Xi, G, H, *, \Delta, \perp)$ be a complete OIFBMS and $\xi : \Xi \rightarrow \Xi$ be an \perp -continuous and \perp -preserving. Let $\rho \in (0, \frac{1}{\theta})$ for all $\tau > 0$, with

$$G(\xi w, \xi k, \rho\tau) \geq \min\{G(\xi w, w, \tau), G(\xi k, k, \tau)\},$$

$$H(\xi w, \xi k, \rho\tau) \leq \min\{H(\xi w, w, \tau), H(\xi k, k, \tau)\}.$$

Then, ξ has a unique FP. Furthermore, $\lim_{n \rightarrow \infty} G(\xi^n w, w_*, \tau) = 1$ and $\lim_{n \rightarrow \infty} H(\xi^n w, w_*, \tau) = 0$, for all $w \in \Xi$ and $\tau > 0$.

Proof. It is obvious from Theorem 3.14 and 3.17 \square

Example 3.19. Suppose $\Xi = [-2, 2]$ and by \perp by $w \perp k \Leftrightarrow w + k \geq 0$. Define G and H by

$$G(w, k, \tau) = \begin{cases} 1 & \text{if } w = k, \\ \frac{\tau}{\tau + \max\{w, k\}^\alpha} & \text{otherwise} \end{cases} \quad (28)$$

$$H(w, k, \tau) = \begin{cases} 0 & \text{if } w = k, \\ \frac{\max\{w, k\}^\alpha}{\tau + \max\{w, k\}^\alpha} & \text{otherwise} \end{cases} \quad (29)$$

for all $w, k \in \Xi$ and $\tau > 0$, with $\sigma * \theta = \sigma \cdot \theta$ and $\sigma \Delta \theta = \max\{\sigma, \theta\}$, Then $(\Xi, G, H, *, \Delta, \perp)$ is an O-complete IFBMS. Note that $\lim_{n \rightarrow \infty} Gw, k, \tau = 1$ and $\lim_{n \rightarrow \infty} Hw, k, \tau = 0$. Define $\xi : \Xi \rightarrow \Xi$ by

$$\xi(w) = \begin{cases} \frac{w}{4}, & w \in [-2, \frac{2}{3}], \\ 1 - w, & w \in (\frac{2}{3}, 1], \\ w - \frac{1}{2}, & w \in (1, 2]. \end{cases} \quad (30)$$

There are following four cases:

1. If $w, k \in [-2, \frac{2}{3}]$ then $\xi(w) = \frac{w}{4}$ and $\xi(k) = \frac{k}{4}$.
2. If $w, k \in (\frac{2}{3}, 1]$ then $\xi(w) = 1 - w$ and $\xi(k) = 1 - k$.
3. If $w, k \in (1, 2]$ then $\xi(w) = w - \frac{1}{2}$ and $\xi(k) = k - \frac{1}{2}$.
4. If $w \in [-2, \frac{2}{3}]$ and $k \in (\frac{2}{3}, 1]$ then $\xi(w) = \frac{w}{4}$ and $\xi(k) = k - \frac{1}{2}$.
5. If $w \in [-2, \frac{2}{3}]$ and $k \in (1, 2]$ then $\xi(w) = \frac{w}{4}$ and $\xi(k) = k - \frac{1}{2}$.
6. If $w \in (\frac{2}{3}, 1]$ and $k \in (\frac{2}{3}, 1]$ then $\xi(w) = 1 - w$ and $\xi(k) = k - \frac{1}{2}$.
7. If $w \in (1, 2]$ and $k \in (\frac{2}{3}, 1]$ then $\xi(w) = w - \frac{1}{2}$ and $\xi(k) = 1 - k$.
8. If $w \in (1, 2]$ and $k \in [-2, \frac{2}{3}]$ then $\xi(w) = w - \frac{1}{2}$ and $\xi(k) = \frac{k}{4}$.
9. If $w \in (\frac{2}{3}, 1]$ and $k \in [-2, \frac{2}{3}]$ then $\xi(w) = 1 - w$ and $\xi(k) = \frac{k}{4}$.

Because $w \perp k \Leftrightarrow w + k \geq 0$, it is clearly implies that $\xi w + k \geq 0$. that is, ξ is \perp -preserving. Suppose $\{w_n\}$ be any O-sequence in Ξ that O-converges to $w \in \Xi$. We get

$$\lim_{n \rightarrow \infty} G(w_n, w, \tau) = \lim_{n \rightarrow \infty} \frac{\tau}{\tau + \max\{w_n, w\}^3} = 1, \quad (31)$$

$$\lim_{n \rightarrow \infty} H(w_n, w, \tau) = \lim_{n \rightarrow \infty} \frac{\max\{w_n, w\}^3}{\tau + \max\{w_n, w\}^3} = 0, \quad (32)$$

Note that if $G(w_n, w, \tau) = 1$ and $H(w_n, w, \tau) = 0$, then $G(\xi w_n, \xi w, \tau) = 1$ and $H(\xi w_n, \xi w, \tau) = 0$ for all $\tau > 0$. that is, ξ is orthogonal continuous. For $w = k$, it is obvious. Assume $w \neq k$. We get

$$G(\xi w, \xi k, \rho \tau) \geq \min\{G(\xi w, w, \tau), G(\xi k, k, \tau)\}$$

$$H(\xi w, \xi k, \rho \tau) \leq \min\{H(\xi w, w, \tau), H(\xi k, k, \tau)\}.$$

It fulfilled above all cases. Now, we show that ξ is not a contraction. Suppose

$$\min\{G(\xi w, w, \tau), G(\xi k, k, \tau)\} = G(\xi w, w, \tau)$$

$$\min\{H(\xi w, w, \tau), H(\xi k, k, \tau)\} = H(\xi w, k, \tau).$$

then for $w = -1$ and $k = -2$, we have

$$G(\xi w, \xi k, \varrho\tau) = \frac{\varrho\tau}{\varrho\tau + \max\{\frac{w}{4}, \frac{k}{4}\}^3} = \frac{64\varrho\tau}{64\varrho\tau - 1} \geq 1,$$

$$H(\xi w, \xi k, \varrho\tau) = \frac{\max\{\frac{w}{4}, \frac{k}{4}\}^3}{\varrho\tau + \max\{\frac{w}{4}, \frac{k}{4}\}^3} = \frac{-1}{64\varrho\tau - 1} \leq 0.$$

Which is not true. That is, all assumptions of Theorem 2.2 are fulfilled, and 0 is a unique FP of ξ .

Definition 3.20. Suppose $(\Xi, G, H, *, \Delta, \perp)$ be an OIFBMS. A mapping $\xi : \Xi \rightarrow \Xi$ is called a fuzzy θ - \perp -contraction if there exists $\varrho \in (0, 1)$ such that

$$\frac{1}{G(\xi w, \xi k, \tau)} - 1 \leq \varrho \left[\frac{1}{G(w, k, \tau)} - 1 \right] \quad (33)$$

$$H(\xi w, \xi k, \tau) \leq \varrho H(w, k, \tau) \quad (34)$$

for all $w, k \in \Xi$ and $\tau > 0$. Where ϱ is said to be an IFB- \perp -contractive constant of ξ .

Theorem 3.21. Suppose $(\Xi, G, H, *, \Delta, \perp)$ be an OIFBMS. Such that

$$\lim_{\tau \rightarrow \infty} G(w, k, \tau) = 1, \quad (35)$$

$$\lim_{\tau \rightarrow \infty} H(w, k, \tau) = 0, \forall w, k \in \Xi. \quad (36)$$

Assume a mapping $\xi : \Xi \rightarrow \Xi$ be a \perp -continuous, IFB- \perp -contraction and \perp -preserving mapping. Thus, ξ has a FP, call $\nu \in \Xi$. Moreover, $G(\nu, \nu, \alpha) = 1$ and $H(\nu, \nu, \alpha) = 0$ for all $\alpha > 0$.

Proof. Suppose $(\Xi, G, H, *, \Delta, \perp)$ be an O-complete IFBMS. For any point $w_0 \in \Xi$, $w_0 \perp k$, for all $k \in \Xi$. That is, $w_0 \perp \xi w_0$. Consider $w_n = \xi^n w_0 = \xi w_{n-1}$ for all $n \in \mathbb{N}$. Therefore, ξ is \perp -preserving and $\{w_n\}$ is an O-sequence. If $w_n = w_{n-1}$ for some $n \in \mathbb{N}$ then w_n is a FP of ξ . We suppose that $w_n \neq w_{n-1}$ for all $n \in \mathbb{N}$. For all $\tau > 0$, $n \in \mathbb{N}$ and utilizing (9), we have

$$\frac{1}{G(w_n, w_{n+1}, \tau)} - 1 = \frac{1}{G(\xi w_{n-1}, \xi w_n, \tau)} - 1 \leq \varrho \left[\frac{1}{G(w_{n-1}, w_n, \tau)} - 1 \right]$$

$$H(w_n, w_{n+1}, \tau) = H(\xi w_{n-1}, \xi w_n, \tau) \leq \varrho H(w_{n-1}, w_n, \tau).$$

We have

$$\frac{1}{G(w_n, w_{n+1}, \tau)} - 1 = \frac{\varrho}{G(w_{n-1}, w_n, \tau)} + (1 - \varrho), \forall \tau > 0$$

$$\frac{\varrho}{G(\xi w_{n-2}, \xi w_{n-1}, \tau)} + (1 - \varrho) \leq \frac{\varrho^2}{G(w_{n-2}, w_{n-1}, \tau)} + \varrho(1 - \varrho) + (1 - \varrho).$$

Continuing in this way, we get

$$\begin{aligned} \frac{\varrho}{G(w_n, w_{n+1}, \tau)} &\leq \frac{\varrho^n}{G(w_0, w_1, \tau)} + \varrho^{n-1}(1 - \varrho) + \varrho^{n-2}(1 - \varrho) + \dots + \varrho(1 - \varrho) + (1 - \varrho). \\ &\leq \frac{\varrho^n}{G(w_0, w_1, \tau)} + (\varrho^{n-1} + \varrho^{n-2} + \dots + 1)(1 - \varrho) \end{aligned}$$

$$\leq \frac{\varrho^n}{G(w_0, w_1, \tau)} + (1 - \varrho^n)$$

We have

$$\frac{1}{\frac{\varrho^n}{G(w_0, w_1, \tau)} + (1 - \varrho^n)} \leq G(w_n, w_{n+1}, \tau), \forall \tau > 0, n \in \mathbb{N} \quad (37)$$

and

$$\begin{aligned} H(w_n, w_{n+1}, \tau) &= H(\xi w_{n-1}, \xi w_n, \tau) \leq \varrho H(w_{n-1}, w_n, \tau) = \varrho H(\xi w_{n-2}, \xi w_{n-1}, \tau) \\ &\leq \varrho^2 H(w_{n-2}, w_{n-1}, \tau) \leq \cdots \leq \varrho^n H(w_0, w_1, \tau) \forall \tau > 0, n \in \mathbb{N} \end{aligned} \quad (38)$$

Now, for $m \geq 1$ and $n \in \mathbb{N}$, we have

$$\begin{aligned} G(w_n, w_{n+m}, \tau) &\geq G\left(w_n, w_{n+1}, \frac{\tau}{\theta}\right) * G\left(w_{n+1}, w_{n+2}, \frac{\tau}{\theta^2}\right) \\ &\geq G\left(w_n, w_{n+1}, \frac{\tau}{\theta}\right) * G\left(w_{n+1}, w_{n+2}, \frac{\tau}{\theta^2}\right) * G\left(w_{n+2}, w_{n+3}, \frac{\tau}{\theta^3}\right) \end{aligned}$$

Again, continuing in this way, we get

$$G(w_n, w_{n+m}, \tau) \geq G\left(w_n, w_{n+1}, \frac{\tau}{\theta}\right) * G\left(w_{n+1}, w_{n+2}, \frac{\tau}{\theta^2}\right) * \cdots * G\left(w_{n+m-1}, w_{n+m}, \frac{\tau}{\theta^{m-1}}\right)$$

and

$$\begin{aligned} H(w_n, w_{n+m}, \tau) &\leq H\left(w_n, w_{n+1}, \frac{\tau}{\theta}\right) \Delta H\left(w_{n+1}, w_{n+2}, \frac{\tau}{\theta^2}\right) \\ &\leq H\left(w_n, w_{n+1}, \frac{\tau}{\theta}\right) \Delta H\left(w_{n+1}, w_{n+2}, \frac{\tau}{\theta^2}\right) \Delta H\left(w_{n+2}, w_{n+3}, \frac{\tau}{\theta^3}\right) \end{aligned}$$

Continuing in this way, we get

$$H(w_n, w_{n+m}, \tau) \leq H\left(w_n, w_{n+1}, \frac{\tau}{\theta}\right) \Delta H\left(w_{n+1}, w_{n+2}, \frac{\tau}{\theta^2}\right) \Delta \cdots \Delta H\left(w_{n+m-1}, w_{n+m}, \frac{\tau}{\theta^{m-1}}\right)$$

By utilizing (37) in the above inequality, we get

$$\begin{aligned} G(w_n, w_{n+m}, \tau) &\geq \frac{1}{\frac{\varrho^n}{G(w_0, w_1, \frac{\tau}{\theta})} + (1 - \varrho^n)} * \frac{1}{\frac{\varrho^{n+1}}{G(w_0, w_1, \frac{\tau}{\theta^2})} + (1 - \varrho^{n+1})} * \cdots \\ &\quad * \frac{1}{\frac{\varrho^{n+m-1}}{G(w_0, w_1, \frac{\tau}{\theta^{m-1}})} + (1 - \varrho^{n+m-1})} \\ &\geq \frac{1}{\frac{\varrho^n}{G(w_0, w_1, \frac{\tau}{\theta})} + 1} * \frac{1}{\frac{\varrho^{n+1}}{G(w_0, w_1, \frac{\tau}{\theta^2})} + 1} * \cdots * \frac{1}{\frac{\varrho^{n+m-1}}{G(w_0, w_1, \frac{\tau}{\theta^{m-1}})} + 1} \end{aligned}$$

Also, using (38), we have

$$H(w_n, w_{n+p}, \tau) \leq H\left(w_n, w_{n+1}, \frac{\tau}{\theta}\right) \Delta H\left(w_{n+1}, w_{n+2}, \frac{\tau}{\theta^2}\right) \Delta \cdots \Delta H\left(w_{n+m-1}, w_{n+m}, \frac{\tau}{\theta^{m-1}}\right)$$

As $\varrho \in (0, 1)$, we have $\lim_{n \rightarrow \infty} G(w_n, w_{n+m}, \tau) = 1$ and $\lim_{n \rightarrow \infty} H(w_n, w_{n+m}, \tau) = 0$ for all $\tau > 0$, $m \geq 1$. Therefore, a sequence $\{w\}$ is an O -Cauchy in $(\Xi, G, H, *, \Delta, \perp)$ is complete, and we have ξ is an \perp -continuous, there exist $\nu \in \Xi$ such that

$$\lim_{n \rightarrow \infty} G(w_{n+1}, \nu, \tau) = \lim_{n \rightarrow \infty} G(\xi w_n, \xi \nu, \tau) = 1, \forall \tau > 0, \quad (39)$$

$$\lim_{n \rightarrow \infty} H(w_{n+1}, \nu, \tau) = \lim_{n \rightarrow \infty} H(\xi w_n, \xi \nu, \tau) = 0, \forall \tau > 0, \quad (40)$$

Now, we show that ν is a FP of ξ . By utilizing (33), we have

$$\frac{1}{G(\xi w, \xi \nu, \tau)} - 1 \leq \rho \left[\frac{1}{G(w_n, \xi \nu, \tau)} - 1 \right] = \frac{\rho}{G(w, \xi \nu, \tau)} - \rho.$$

That is,

$$\frac{1}{G(\xi w, \xi \nu, \tau) + 1 - \rho} \leq G(\xi w_n, \xi \nu, \tau).$$

Using the above inequality, we obtain

$$\begin{aligned} G(\nu, \xi \nu, \tau) &\geq G\left(\nu, w_{n+1}, \frac{\tau}{2\theta}\right) * G\left(w_{n+1}, \xi \nu, \frac{\tau}{2\theta}\right) \\ &= G\left(\nu, w_{n+1}, \frac{\tau}{2\theta}\right) * G\left(\xi w_n, \xi \nu, \frac{\tau}{2\theta}\right) \\ &\geq G\left(\nu, w_{n+1}, \frac{\tau}{2\theta}\right) * \frac{\rho}{G(w_n, \nu, \frac{\tau}{2\theta}) + 1 - \rho} \end{aligned}$$

and

$$\begin{aligned} H(w, \nu, \tau) &= H(\xi w, \xi \nu, \tau) \leq \rho H(w, \nu, \tau) < H(w, \nu, \tau) \\ &= H\left(w, w_{n+1}, \frac{\tau}{2\theta}\right) \Delta H\left(\xi w_n, \xi \nu, \frac{\tau}{2\theta}\right) \\ &\leq H\left(w, w_{n+1}, \frac{\tau}{2\theta}\right) \Delta \rho H\left(w_n, w, \frac{\tau}{2\theta}\right) \end{aligned}$$

Taking limit as $n \rightarrow \infty$ and using (39) and (40) in the above expression, we get $G(\nu, \xi \nu, \tau) = 1$, that is, $\xi \nu = \nu$. Therefore, ν is a FP of ξ , and $G(\nu, \nu, \tau) = 1$ and $H(\nu, \nu, \tau) = 0$ for all $\tau > 0$. \square

Corollary 3.22. Suppose $(\Xi, G, H, *, \Delta, \perp)$ be an O-complete IFBMS such that $\lim_{n \rightarrow \infty} G(w, k, \tau) = 1$ and $\lim_{n \rightarrow \infty} H(w, k, \tau) = 0$, for all $w, k \in \Xi$ and $\xi : \Xi \rightarrow \Xi$ satisfy

$$\frac{1}{G(\xi^n w, \xi^n k, \tau)} - 1 \leq \rho \left[\frac{1}{G(w, k, \tau)} - 1 \right] \quad (41)$$

$$H(\xi^n w, \xi^n k, \tau) \leq \rho H(w, k, \tau) \quad (42)$$

for all $n \in \mathbb{N}$, $w, k \in \Xi$, $\tau > 0$, where $0 < \rho < 1$. Then ξ has a FP, say $\nu \in \Xi$ and $G(\nu, \nu, \tau) = 1$, for all $\tau > 0$.

Proof. $\nu \in \Xi$ is a unique FP of ξ^n by utilizing Theorem 3.22, and $G(\nu, \nu, \tau) = 1$, for all $\tau > 0$. $\xi \nu$ is also a FP of $\xi^n(\xi \nu) = \xi \nu$ from Theorem 3.22, $\xi \nu = \nu$. Hence, the FP of ξ is also a FP of ξ^n . \square

Example 3.23. Suppose $\Xi = [-1, 2]$ and define \perp by $w \perp k \Leftrightarrow w + k \geq 0$. Define G, H as in Example 3.4 with $\alpha = 3$,

$$G(w, k, \tau) = \frac{\tau + \min\{w, k\}^3}{\tau + \min\{w, k\}^3} \forall w, k \in \Xi, \tau > 0, \quad (43)$$

and

$$H(w, k, \tau) = 1 - \frac{\tau + \min\{w, k\}^3}{\tau + \min\{w, k\}^3} \forall w, k \in \Xi, \tau > 0, \quad (44)$$

with $\sigma * \theta = \sigma \cdot \theta$ and $\sigma \Delta \theta = \max\{\sigma, \theta\}$, then $(\Xi, G, H, *, \Delta, \perp)$ is an O-complete IFBMS. see that $\lim_{\tau \rightarrow \infty} G(w, k, \tau) = 1$ and $\lim_{\tau \rightarrow \infty} H(w, k, \tau) = 0$ for all $w, k \in \Xi$. Define $\xi : \Xi \rightarrow \Xi$ by

$$G(w, k, \tau) = \begin{cases} 2 - w & w \in [-1, 1), \\ 1 & w \in [1, 2), \end{cases} \quad (45)$$

We have the following four cases:

1. if $w, k \in [-1, 1)$ then $\xi w = 2 - w$ and $\xi k = 2 - k$,
2. if $w, k \in [1, 2]$ then $\xi w = \xi k = 1$,
3. if $w \in [-1, 1)$ and $k \in [1, 2]$ then $\xi w = 2 - w$ and $\xi k = 1$,
3. if $w \in [1, 2]$ and $k \in [-1, 1)$ then $\xi w = 1$ and $\xi k = 2 - k$,

Because $w \perp k \Leftrightarrow w + k \geq 0$, it is clearly implies that $\xi(w) + \xi(k) \geq 0$. That is, ξ is \perp -preserving. Suppose $\{w_n\}$ be any O-sequence in Ξ that O-converges to $w \in \Xi$. we get

$$\lim_{n \rightarrow \infty} G(w, k, \tau) = \lim_{n \rightarrow \infty} \frac{\tau + \min\{w, k\}^3}{\tau + \min\{w, k\}^3} = 1 \forall w, k \in \Xi, \tau > 0,$$

and

$$\lim_{n \rightarrow \infty} H(w, k, \tau) = 1 - \lim_{n \rightarrow \infty} \frac{\tau + \min\{w, k\}^3}{\tau + \min\{w, k\}^3} = 0 \forall w, k \in \Xi, \tau > 0,$$

we can easily see that if $\lim_{n \rightarrow \infty} G(w_n, w, \tau) = 1$, and $\lim_{n \rightarrow \infty} H(w_n, w, \tau) = 0$, then $\lim_{n \rightarrow \infty} G(\xi w_n, \xi w, \tau) = 1$ and $\lim_{n \rightarrow \infty} H(\xi w_n, \xi w, \tau) = 0$ for all $\tau > 0$. That is, ξ is orthogonal continuous. For $w = k$, it is obvious.

$$\frac{1}{G(\xi w, \xi k, \tau)} - 1 \leq \varrho \left[\frac{1}{G(w, k, \tau)} - 1 \right]$$

$$H(\xi w, \xi k, \tau) \leq \varrho H(w, k, \tau).$$

All conditions of Theorem 3.21 are satisfied and 1 is a FP of ξ

4 An Application to an Integreal Equation

Let $\Xi = C([\sigma, \theta], \mathbb{R})$ be the set of all continuous real valued functions defined on $[\sigma, \theta]$. Now, we consider the Fredholm type integral equation of first kind:

$$w(\eta) = \int_{\sigma}^{\theta} F(\eta, j)w(\eta)kj, \text{ for } \eta, j \in [\sigma, \theta] \quad (46)$$

Where, $F \in \Xi$. Define G as in Example 3.2, That is

$$G(w(\eta), k(\eta), \tau) = \sup_{\eta \in [\sigma, \theta]} \begin{cases} 1 & \text{if } w = k, \\ \left[e^{\frac{\max\{w(\eta), k(\eta)\}^{\alpha}}{\tau}} \right]^{-1} & \text{otherwise,} \end{cases} \quad (47)$$

and

$$H(w(\eta), k(\eta), \tau) = \sup_{\eta \in [\sigma, \theta]} \begin{cases} 0 & \text{if } w = k, \\ 1 - \left[e^{\frac{\max\{w(\eta), k(\eta)\}^{\alpha}}{\tau}} \right]^{-1} & \text{otherwise,} \end{cases} \quad (48)$$

for all $w, k \in \Xi$ and $\tau > 0$. Then $(\Xi, G, H, *, \Delta, \perp)$ is an O-complete IFBMS.

Theorem 4.1. Assume that $\max\{F(\eta, j)w(\eta), F(\eta, j)k(\eta)\} \leq \varrho \max\{w(\eta), k(\eta)\}$ for $w, k \in \Xi$, $\varrho \in (0, 1)$ and $\eta, j \in [\sigma, \theta]$. Also, consider $\int_{\sigma}^{\theta} kj = 1$. Then the Fredholm type integral equation of first kind in equation (46) has a unique solution.

Proof. Define $\xi : \Xi \rightarrow \Xi$ by $w(\eta) = \int_{\sigma}^{\theta} F(\eta, j)w(\eta)kj, \text{ for } \eta, j \in [\sigma, \theta]$. Define Orthogonality as: $w(\eta) \perp k(\eta) \Leftrightarrow w(\eta)k(\eta) \in \{|w(\eta)|, |k(\eta)|\}$. We see that $w(\eta)$ and $\xi w(\eta)$ belong to Ξ . So, observe that if $w(\eta) \perp k(\eta)$,

then must be $\xi w(\eta) \perp \xi k(\eta)$. Observe that the existence of a FP of the operator ξ is equivalent to the existence of a solution of the Fredholm type integral equation (46). Now, for $w(\eta) = k(\eta)$, the contraction condition holds. While for $w \neq k$, We have

$$\begin{aligned} G(\xi w(\eta), \xi k(\eta), \varrho\tau) &= \left[e^{\frac{\max\{w(\eta), k(\eta)\}^\alpha}{\varrho\tau}} \right]^{-1} \\ &= \left[e^{\frac{\max\{ \int_\sigma^\theta F(\eta, j)w(\eta)k_j, \int_\sigma^\theta F(\eta, j)k(\eta)k_j \}^\alpha}{\varrho\tau}} \right]^{-1} \\ &= \left[e^{\frac{(\int_\sigma^\theta \max\{F(\eta, j)w(\eta)k_j, F(\eta, j)k(\eta)k_j\})^\alpha}{\varrho\tau}} \right]^{-1} \\ &\geq \left[e^{\frac{(\int_\sigma^\theta \max\{w(\eta)k_j, k(\eta)k_j\})^\alpha}{\varrho\tau}} \right]^{-1} \\ &\geq \sup_{\eta \in [\sigma, \theta]} \left[e^{\frac{(\varrho \max\{w(\eta)k_j, k(\eta)\})^\alpha (\int_\sigma^\theta k_j)^\alpha}{\varrho\tau}} \right]^{-1} \\ &= \sup_{\eta \in [\sigma, \theta]} \left[e^{\frac{(\max\{w(\eta)k_j, k(\eta)\})^\alpha}{\tau}} \right]^{-1} \\ &= G(w(\eta), k(\eta), \tau), \end{aligned}$$

and

$$\begin{aligned} H(\xi w(\eta), \xi k(\eta), \varrho\tau) &= 1 - \left[e^{\frac{\max\{w(\eta), k(\eta)\}^\alpha}{\varrho\tau}} \right]^{-1} \\ &= 1 - \left[e^{\frac{\max\{ \int_\sigma^\theta F(\eta, j)w(\eta)k_j, \int_\sigma^\theta F(\eta, j)k(\eta)k_j \}^\alpha}{\varrho\tau}} \right]^{-1} \\ &= 1 - \left[e^{\frac{(\int_\sigma^\theta \max\{F(\eta, j)w(\eta)k_j, F(\eta, j)k(\eta)k_j\})^\alpha}{\varrho\tau}} \right]^{-1} \\ &\leq 1 - \left[e^{\frac{(\int_\sigma^\theta \max\{w(\eta)k_j, k(\eta)k_j\})^\alpha}{\varrho\tau}} \right]^{-1} \\ &\leq 1 - \sup_{\eta \in [\sigma, \theta]} \left[e^{\frac{(\varrho \max\{w(\eta)k_j, k(\eta)\})^\alpha (\int_\sigma^\theta k_j)^\alpha}{\varrho\tau}} \right]^{-1} \\ &= 1 - \sup_{\eta \in [\sigma, \theta]} \left[e^{\frac{(\max\{w(\eta)k_j, k(\eta)\})^\alpha}{\tau}} \right]^{-1} \\ &= H(w(\eta), k(\eta), \tau), \end{aligned}$$

Hence, ξ is an \perp -contraction. Let $\{w_n\}$ be an O -sequence in Ξ O -converging to $w \in \Xi$. Because ξ is an \perp -preserving, then $\{\xi w_n\}$ is an O -sequence for each $n \in \mathbb{N}$. We have

$$G(\xi w_n(\eta), \xi w, \varrho\tau) \geq G(w_n(\eta), w(\eta), \tau) \tag{49}$$

and

$$H(\xi w_n(\eta), \xi w, \varrho\tau) \leq H(w_n(\eta), w(\eta), \tau) \quad (50)$$

As $\lim_{n \rightarrow \infty} G(\xi w_n(\eta), \xi w, \varrho\tau) = 1$ and $\lim_{n \rightarrow \infty} H(\xi w_n(\eta), \xi w, \varrho\tau) = 0$ for all $\tau > 0$, it is clear that

$$\lim_{n \rightarrow \infty} G(\xi w_n(\eta), \xi w, \varrho\tau) = 1, \quad (51)$$

$$\lim_{n \rightarrow \infty} H(\xi w_n(\eta), \xi w, \varrho\tau) = 0, \quad (52)$$

Hence, ξ is \perp -continuous. Therefore, all conditions of Theorem 3.13 are satisfied. Hence, the operator ξ has a unique FP. That is, the Fredholm type integral equation (46) has a unique solution. \square

5 Conclusion

In this study, we established the concept of an OIFBMS as a generalization of an IFBMS. We established some fixed point theorems and solved some non-trivial examples with an application to Fredholm integral equations. This work is extendable in the structure of orthogonal neutrosophic b-metric spaces, and orthogonal intuitionistic fuzzy controlled metric spaces and we can increase self mappings to get new results.

Acknowledgements: We would like to thank the reviewers for their thoughtful comments and efforts towards improving our manuscript.

Conflict of Interest: The authors declare no conflicts of interest.

References

- [1] Alaca C, Turkoglu D, Yildiz C. Fixed points in intuitionistic fuzzy metric spaces. *Chaos, Solitons & Fractals*. 2006; 29(5): 1073-1078. DOI: <https://doi.org/10.1016/j.chaos.2005.08.066>
- [2] Bakhtin IA. The contraction mapping principle in quasimetric spaces. *Functional analysis*. 1989; 30: 26-37. DOI: <https://doi.org/10.4236/oalib.1104657>
- [3] Baleanu D, Rezapour S, Mohammadi H. Some existence results on nonlinear fractional differential equations. *Philosophical Transaction of the Royal Society A: Mathematical, Physical and Engineering Sciences*. 2013; 371(1990): 1-7. DOI: <https://doi.org/10.1098/rsta.2012.0144>
- [4] Czerwil S. Contraction mappings in b-metric spaces. *Acta Mathematica et Informatica Universitatis Ostraviensis*. 1993; 1(1): 5-11. DOI: <http://dml.cz/dmlcz/120469>
- [5] Deng Z. Fuzzy pseudo-metric spaces. *Journal of Mathematical Analysis and Applications*. 1982; 86(1): 74-95. DOI: [https://doi.org/10.1016/0022-247X\(82\)90255-4](https://doi.org/10.1016/0022-247X(82)90255-4)
- [6] Doenovic T, Javaheri A, Sedghi S, Shobe N. Coupled fixed point theorem in b-fuzzy metric spaces. *Novi Sad J. Math*. 2017; 47(1): 77-88. DOI: <https://doi.org/10.1515/fascmath-2018-0015>
- [7] Eshaghi Gordji M, Habibia H. Fixed point theory in generalized orthogonal metric space. *Journal of Linear and Topological Algebra*. 2017; 6(03): 251-260. DOI: <https://dorl.net/dor/20.1001.1.22520201.2017.06.03.7.7>
- [8] Eshaghi Gordji M, Ramezani M, De La Sen M, Cho YJ. On orthogonal sets and Banachs fixed point theorem. *Fixed Point Theory*. 2017; 18(2): 569-578. DOI: <https://doi.org/10.24193/fpt-ro.2017.2.45>

-
- [9] George A, Veeramani P. On some results in fuzzy metric spaces. *Fuzzy Sets and Systems.* 1994; 64(3): 395-399. DOI: [https://doi.org/10.1016/0165-0114\(94\)90162-7](https://doi.org/10.1016/0165-0114(94)90162-7)
- [10] George A, Veeramani P. On some results of analysis for fuzzy metric spaces. *Fuzzy Sets and Systems.* 1997; 90(3): 365-368. DOI: [https://doi.org/10.1016/S0165-0114\(96\)00207-2](https://doi.org/10.1016/S0165-0114(96)00207-2)
- [11] Hezarjaribi M. Fixed point result in orthogonal fuzzy metric space. *Jordan Journal of Mathematics and Statistics.* 2018; 11(4): 295-308.
- [12] Javed K, Uddin F, Aydi H, Arshad M, Ishtiaq U, Alsamir H. On fuzzy b-metric-like spaces. *Journal of Function Spaces.* 2021; 2021: 1-9. DOI: <https://doi.org/10.1155/2021/6615976>
- [13] Konwar N. Extension of fixed results in intuitionistic fuzzy b-metric spaces. *Journal of Intelligent & Fuzzy Systems.* 2020; 39(5): 7831-7841. DOI: <https://doi.org/10.3233/JIFS-201233>
- [14] Kramosil I, Michlek j. Fuzzy metric and statistical metric spaces. *Kybernetika.* 1975; 11(5): 336-344. DOI: <http://dml.cz/dmlcz/125556>
- [15] Mehmood F, Ali R, Ionescu C, Kamran T. Extended fuzzy b-metric spaces. *Journal of Mathematical Analysis.* 2017; 8(6): 124-131.
- [16] Mohamad A. Fixed-point theorems in intuitionistic fuzzy metric spaces. *Chaos, Solitons & Fractals.* 2007; 34(5): 1689-1695. DOI: <https://doi.org/10.1016/j.chaos.2006.05.024>
- [17] Ndban S. Fuzzy b-metric spaces. *International Journal of Computers Communications & Control.* 2016; 11(2): 273-281. DOI: <https://doi.org/10.15837/IJCCC.2016.2.2443>
- [18] Park JH. Intuitionistic fuzzy metric spaces. *Chaos, Solitons & Fractals.* 2004; 22(5): 1039-1046. DOI: <https://doi.org/10.1016/j.chaos.2004.02.051>
- [19] Rafi M, Noorani MSM. Fixed point theorem on intuitionistic fuzzy metric space. *Iranian Journal of Fuzzy Systems.* 2006; 3(1): 23-29. DOI: <https://doi.org/10.22111/IJFS.2006.428>
- [20] Rakic D, Mukheimer A, Doenovic T, Mitrovic ZD, Radenovic S. On some new fixed point results in fuzzy b-metrics paces. *Journal of Inequalities and Applications.* 2020; 2020:1-14. DOI: <https://doi.org/10.1186/s13660-020-02371-3>
- [21] Schweizer B, Sklar A. Statistical metric spaces. *Pacific Journal of Mathematics.* 1960; 10(1): 313-334. DOI: <http://dx.doi.org/10.2140/pjm.1960.10.313>
- [22] Sedghi S, Shobe N. Common fixed point theorem in b-fuzzy metric space. *Nonlinear Functional Analysis and Applications.* 2012; 17: 349-359. DOI: <https://doi.org/10.1155/2021/6615976>
- [23] Senapati T, Dey LK, Damjanovic B, Chanda A. New fixed point results in orthogonal metric Space with an application. *Kragujevac Journal of Mathematics.* 2018; 42(4): 505-516. DOI: <https://doi.org/10.5937/KgJMath1804505S>
- [24] Shukla S, Abbas M. Fixed point results in fuzzy metric-like spaces. *Iranian Journal of Fuzzy Systems.* 2014; 11(5): 81-92. DOI: <https://doi.org/10.22111/ijfs.2014.1724>
- [25] Sintunavarat W, Kumam P. Fixed theorems for a generalized intuitionistic fuzzy contraction in intuitionistic fuzzy metric spaces. *Thai Journal of Mathematics.* 2012; 10(1): 123-135.

- [26] Sudsutad W, Tariboon J. Boundary value problems for fractional differential equations with three-point fractional integral boundary conditions. *Advances in Difference Equations*. 2012; 2012(93): 1-10. DOI: <https://doi.org/10.1186/1687-1847-2012-93>
- [27] Uddin F, Park C, Javed K, Arshad M, Lee JR. Orthogonal m-metric spaces and an application to solve integral equations. *Advance in Difference Equations*. 2021; 2021: 1-15. DOI: <https://doi.org/10.1186/s13662-021-03323-x>
- [28] Zadeh LA. Fuzzy sets. *Information and Control*. 1965; 8(3): 338-353. DOI: [https://doi.org/10.1016/S0019-9958\(65\)90241-X](https://doi.org/10.1016/S0019-9958(65)90241-X)

Fahim Uddin

Abdus Salam School of Mathematical Sciences
Government College University
Lahore, Pakistan.
E-mail: fahamiiu@gmail.com

Muhammad Saeed

Department of Mathematics
University of Management and Technology
Lahore, Pakistan.
E-mail: muhammad.saeed@umt.edu.pk

Khaleel Ahmad

Department of Mathematics
University of Management and Technology
Lahore, Pakistan.
E-mail: khalil7066616@gmail.com

Umar Ishtiaq

Department of Mathematics
Quaid-i-Azam University
Islamabad, Pakistan.
E-mail: umarishtiaq000@gmail.com

Salvatore Sessa

Department of Mathematics
University of Naples Federico II
Naples, Italy.
E-mail: salvatore.sessa2@unina.it