



# The Effects of high-intensity Resistance-Endurance Training on Physical Fitness, Levels of serum Cortisol, Testosterone, and insulin-like growth factor1 (IGF1) among Adolescent Taekwondo players

Mahshid Talesh Sasani<sup>1</sup>, Fatemeh Tavana<sup>2</sup>

<sup>1</sup> M.A. In Exercise Physiology, Department of Physical Education and Sport Sciences, Rasht Branch, Islamic Azad University, Rasht, Iran.

<sup>2</sup> Ph.D. Candidate in Exercise Physiology, Department of Physical Education and Exercise training Science, Rasht Branch, Islamic Azad University, Rasht, Iran.

## Keywords

Taekwondo, Resistance and endurance training, Cortisol, Testosterone, insulin like growth factor1.

## Correspondence

E-mail address: [fatemehtavana@gmail.com](mailto:fatemehtavana@gmail.com)

**Received:** 1 Sept 2024; **Revised:** 2 Oct 2024; **Accepted:** 30 Nov 2024

**DOI:** <https://doi.org/10.71878/jpah.2024.1185990>

## Abstract

**Introduction:** Strength and endurance training has experienced a surge in popularity in recent years and is currently one of the most commonly practiced training methods. This study investigated the effects of high-intensity resistance, and endurance training on physical fitness and levels of serum cortisol, testosterone, and insulin-like growth factor1 (IGF1) among adolescent athletes' taekwondo players.

**Material & Methods:** This quasi-experimental study was based on the pre and post-test design with a control group. The first group engaged in moderate-intensity resistance endurance training along with regular taekwondo training (CG). CG performed a total of 95 minutes of training, including warm-up, cool-down, technical and tactical exercises, 50 minutes of moderate-intensity resistance (60-70% of one repetition maximum; 1RM), and endurance (60-70% of target heart rate; THR) training. During this time, the experimental group (EG) performed a total of 80 minutes of the same training, except that they performed high-intensity resistance (70-80% of 1RM) and endurance (70-80% of THR) training for 35 minutes. Participants included 20 male taekwondo players who were randomly assigned to two equal EG and CG groups with more than two years of training experience. They were evaluated for age, height, weight, and some physical fitness parameters such as aerobic capacity (shuttle run test) and anaerobic capacity (RAST test). CG performed conventional moderate-intensity strength and endurance exercises; the EG performed these exercises at high-intensity training for 80 minutes. The variables were measured before and after 8 weeks of training. Moreover, 10 ml of blood was taken from the arm vein of participants both before and after the study to measure the levels of cortisol, testosterone, and IGF1.

**Results:** Results showed a significant increase in the cortisol level and anaerobic capacity in both CG and EG ( $p < 0.05$ ). In addition, the resting metabolic rate (RMR) increased in the EG, and cortisol level decreased in the CG ( $p < 0.05$ ), but aerobic capacity was promoted in both groups ( $p < 0.05$ ). In the comparison between groups, only anaerobic capacity increased significantly in the experimental group, and no significant differences were observed in other variables.

**Conclusion:** Considering the positive effects of high-intensity aerobic and resistance training on both aerobic and anaerobic power, it is recommended that taekwondo coaches incorporate these exercises into their athletes' training programs. However, further studies are needed to confirm the effects of these exercises on hormonal and bioenergetic changes in the body. The findings underscore the importance of tailored training regimens that consider individual athlete needs and the specific demands of taekwondo.

## 1. Introduction

Taekwondo (TKD) is a martial art characterized by techniques including kicking, blocking, and punching for

offensive or defensive purposes. This sport is an explosive activity requiring athletes to execute various striking techniques with high levels of energy, muscular strength, endurance, flexibility, speed, and agility (1, 2).

Muscle strength enhancement positively influences the rate of force development, thereby improving jumping, sprinting, and agility (3, 4). Strength training aims to enhance an athlete's strength, power, muscular endurance, and muscle hypertrophy. These adaptations may manifest both neurologically and physiologically (5, 6). Resistance training enhances strength and power while mitigating injury risk, serving as a crucial supplementary training modality for elite, amateur (7), young (8), and old (9) athletes, as well as athletes participating in individual (10) and group (11) sports. Moreover, resistance training is regarded as a fundamental training modality in sports such as bodybuilding, weightlifting, powerlifting, and CrossFit (12, 13). Resistance training causes the release of anabolic hormones such as testosterone and growth hormone, and in return, endurance training helps reduce cortisol levels, which is a stress-related hormone. The combination of these two types of training not only improves sports performance but also has positive effects on people's mental health and quality of life (14). By examining the acute and chronic physiological changes of the endocrine system, it was evaluated how nutrition, age, sex, and exercise program can affect the physiological adaptation caused by strength training

By examining the acute and chronic physiological changes of the endocrine system, it was evaluated how nutrition, age, sex, and exercise programs can affect the physiological adaptation caused by strength training. Hormonal surges induced by strength training manifest in four primary ways: acute adaptations during and after exercise; chronic and acute alterations in the training stimulus; improvement in muscle receptors, and chronic alterations in resting levels (6, 15).

Testosterone and GH are both anabolic hormones that increase protein synthesis and prevent protein degradation in muscles (16). Testosterone is produced through the secretion of luteinizing hormone by the anterior pituitary gland. Luteinizing hormone activates Leydig cells to synthesize testosterone. GH is released by the anterior pituitary gland and facilitates protein synthesis in muscles (15, 17). Serum-free testosterone concentrations in both men and women are markedly elevated following a course of resistance training (18, 19). Various components of resistance training can amplify acute responses in testosterone levels. The highest increase in testosterone levels occurs during exercises that engage large muscle groups, such as Olympic lifts that include cleans, squats, and deadlifts. Other factors included high volume and intensity along with short rest periods (20). Testosterone levels peak approximately 15 minutes after exercise and revert to baseline within 1 hour (21).

As a catabolic hormone, cortisol is the principal glucocorticoid hormone (22). Cortisol is synthesized by the adrenal cortex and constitutes 95% of total glucocorticoid activity. Cortisol influences the metabolism of various sources of energy, including glucose, protein, and free fatty acids. Cortisol facilitates the catabolism of proteins into amino acids and promotes growth hormone activity. If cortisol remains unregulated, it may inhibit protein synthesis and lead to protein degradation (23, 24).

The cortisol level follows a circadian rhythm marked by elevated levels upon awakening, a rapid decrease 30-45 minutes thereafter, and a gradual decline until bedtime (25). Brunsden (24) showed that resistance training significantly increases acute cortisol levels, especially in exercise

regimens that concurrently enhance testosterone and GH. The highest increase in serum cortisol levels is observed after high-intensity exercise programs with short rest intervals. Additionally, concurrent training increases cortisol levels more significantly than strength training alone, as physical stress is correlated with an increase in this hormone. Nevertheless, men's cortisol levels experience a decline after 8 weeks of training and revert to their baseline levels after 16 weeks.

Strength training significantly elevates acute levels of GH, with this elevation contingent upon variables including exercise protocol, muscle mass, and overall training volume. Moreover, increasing the volume and intensity of exercises, minimizing rest intervals, and employing multiple sets (supersets) rather than single sets significantly contribute to this elevation. Studies have demonstrated that rest intervals of 30 to 60 seconds between sets can effectively increase the GH level (26). The testosterone-to-cortisol concentration ratio (T/C) serves as an indicator of exercise-induced stress levels. Variations in the levels of these hormones are responsible for regulating numerous exercise-induced responses, such as hypertrophy and enhanced strength (27, 28). Brief rest intervals combined with heightened intensity and volume of exercises can increase lactate levels in the blood. The increased blood lactate concentration results in a heightened acute level of GH. The increased H<sup>+</sup> ions generated from lactic acid have been proposed as the primary mechanism of GH secretion (28). Hormonal adaptations resulting from exercise training in children and adolescents markedly differ from those in adults, likely due to varying stages of maturation (29). High intensity strength and endurance training in athletes can potentially lead to significant reduction in subcutaneous fat. Decreased serum leptin and glucose levels and improved insulin sensitivity can improve the taekwondo player's performance (30). Given the lack of evidence regarding the effects of high-intensity training in adolescents and young adults, this study sought to determine the effect of a high-intensity strength-endurance training on physical fitness and levels of serum cortisol, testosterone, and insulin-like growth factor1 (IGF1) among adolescent athletes' taekwondo players.

## 2. Methodology

### 2.1. Materials and methods

This was a quasi-experimental study based on a pretest-posttest design that was conducted in Rasht, Iran. The control group (CG) performed a total of 95 minutes of training, including warm-up, cool-down, technical and tactical exercises, 50 minutes of moderate-intensity resistance (60-70% of one repetition maximum; 1RM), and endurance (60-70% of target heart rate; THR) training. During this time, the experimental group (EG) performed a total of 80 minutes of the same training, except that they performed high-intensity resistance (70-80% of 1RM) and endurance (70-80% of THR) training for 35 minutes.

### 2.2. Participants

The participants included 20 male taekwondo players who were randomly selected from 40 healthy athletes (with no injury) aged 15-20 years with more than two years of experience in taekwondo training. All participants filled out a specialized questionnaire on physical activity level and

provided their medical records before entering the study; written consent forms were obtained from all participants. The exclusion criteria were injury during the training sessions and absence in more than three consequent sessions. After the general health of participants was ensured, they were invited to a laboratory to brief them on the research objectives and procedures. Participants were randomly assigned to the CG (regular moderate-intensity training group) and EG (high-intensity resistance and endurance training).

### 2.3. Measurements

Participants were evaluated for age, height, weight, and some physical fitness parameters such as aerobic capacity (shuttle run test) and anaerobic capacity (Running-Based Anaerobic Sprint Test; RAST).

**Shuttle run test:** The test is an incremental intermittent running test between two separate lines at a distance of 20 m(31). The initial speed, marked by acoustic signals, is 8.5 km/h, increasing by 0.5 km/h every minute; so, the time that the participants have to cover the distance of 20 m decreases over time until the participant can no longer reach the designated speed (31).  $VO_{2max}$  estimations were based on regression equations, established by Leger et al. The formula used for this calculator is the last stage number used to predict maximal oxygen uptake ( $VO_{2max}$ ,  $ml.kg^{-1}.min^{-1}$ ) from the speed (X,  $km.hr^{-1}$ ) corresponding to that stage (speed =  $8 + 0.5 \times \text{stage no.}$ ) and age (A, years):  $VO_{2max} (ml.kg^{-1}.min^{-1}) = 31.025 + 3.238X - 3.248A + 0.1536AX$

This formula is suitable for boys and girls aged 8-19 years. For adults, similar measurements indicated that the same equation could be used keeping the age constant at 18.

**Running-Based Anaerobic Sprint Test; RAST:** RAST is a testing protocol designed to measure anaerobic power and capacity. The test involves six sprints over a 35-metre distance, with a 10-second recovery between each sprint. The formula for calculating Peak Power Output (PPO) is:

$$PPO = \text{Body mass} * \text{Distance}^2 \div \text{Time}^3$$

Anaerobic capacity (AC) is the total work completed during the test duration.

$$AC = \text{Sum of all six sprint PPOs}(32)$$

**Resting metabolism raten(RMR):** To measure resting metabolic rate (RMR), using an indirect consumption calculation method that involves assessing carbon dioxide consumption and production at rest (33). The Harris-Benedict formula for calculating RMR in men is as follows:

$$RMR = 66.47 + (13.75 \times \text{body mass in kg}) + (5.003 \times \text{height, in cm}) - (6.775 \times \text{age, in years})(34).$$

**Laboratory tests:** 10 ml of blood was taken from the brachial vein of participants both before and after the study to measure the levels of cortisol, testosterone, and IGF1. To prevent blood clotting, the samples were immediately transferred into FL tubes (Italy). Then they were centrifuged (Orom Tajhiz) at 3,000 rpm for 5 minutes to separate the plasma; finally, the samples were kept at  $-20^{\circ}C$  until the end of the study. Testosterone and cortisol levels were assessed utilizing the Roche kit (Germany) and with the use of the electro-chemiluminescence (ECL) technique.

### 2.4. Intervention

The training program for teenage taekwondo athletes consisted of 3 weekly training sessions for 8 weeks. Each training session included a 10-minute warm-up with

stretching and light jogging, and 5 minute cool-down (stretching and deep breathing). It was accompanied by 30 minutes of technical and tactical training. The control group performed 50 minutes of moderate-intensity strength (25 minutes with 60% - 70% one Repetition maximum), which included basic movements such as squats, chest presses, deadlifts, and leg presses, with 3 sets of 10 repetitions, and endurance (25 minutes with 60% - 70% target heart rate) training. The target heart rate was calculated using the Karvenon formula ( $\text{Target HR} = ((\text{Maximum HR} - \text{Resting HR}) \times \text{Intensity \%}) + \text{Resting HR}$ ). Attention to correct form and sufficient rest between sets (60-90 seconds) were important points of this program. The experimental group performed 35 minutes of intense resistance endurance training. They performed 35 minutes of aerobic-strength training, including 15 minutes of endurance training at an intensity of 70-80% of target heart rate and 20 minutes of strength training at an intensity of 70-80% of one repetition maximum (1RM).

### 2.5. Statistical Methods

The normal distribution of the data obtained was examined using the Shapiro-wilk test. The independent t-test was then employed to compare the CG and EG in terms of the data that followed a normal distribution pattern. All statistical tests were performed in SPSS version 23 at a significance level of 0.05.

### 3. Results

Table 1 Shows the individual characters of the groups. The results showed that in the CG (age  $17.70 \pm 2.26$  years, weight  $63.85 \pm 7.04$  kg), and EG (age  $16.9 \pm 1.7$  years, weight  $61.35 \pm 11.61$  kg). The values are presented in the table.

**Table 1.** Individual characteristics of participants

Parameter	Group	Pre-test	Post- test
		Mean $\pm$ SD	Mean $\pm$ SD
Age (years)	CG	17.70 $\pm$ 2.26	---
	EG	16.9 $\pm$ 1.72	---
Height (cm)	CG	169 $\pm$ 0.06	---
	EG	171 $\pm$ 0.09	---
Weight (kg)	CG	63.85 $\pm$ 7.04	64.27 $\pm$ 7.67
	EG	61.35 $\pm$ 11.61	62.60 $\pm$ 12.30

CG: control group; EG: Experimental group

The paired t-test results indicated the significant effects of training intervention on the physiological parameters of the participants. The participants in the EG showed a significant promotion in RMR and testosterone levels ( $p < 0.05$ ), whereas these parameters demonstrated no significant change in the CG. The significant increase in cortisol levels in the CG ( $p < 0.01$ ). The data revealed no significant change in GH in either of the groups. Furthermore, both groups exhibited a significant reduction in aerobic and anaerobic capacity ( $p < 0.01$ ), which can be attributed to the influence of exercise training. These results suggest that exercise training can improve some physiological indicators and positively affect general health (Table 2).

**Table 2.** Results of the paired *t*-test comparing the situation before and after the intervention in the control group (CG) and the EG (*n*=10 in each group)

Parameter	Group	Mean $\pm$ SE (Pre-test)	Mean $\pm$ SE (Post-test)	Mean difference	SEM	<i>t</i> value	df	Sig.
RMR	CG	1198.15 $\pm$ 41.60	1199.55 $\pm$ 43.31	-1.39	1.01	-1.37	9	0.20
	EG	1200.48 $\pm$ 67.44	1204.85 $\pm$ 68.07	-4.36	1.74	-2.50	9	0.03*
Testosterone (ng/ml)	CG	7.34 $\pm$ 2.73	8.85 $\pm$ 2.73	-1.39	0.59	-4.55	9	0.00*
	EG	7.27 $\pm$ 1.83	8.30 $\pm$ 2.30	-1.03	0.3	-3.45	9	0.01*
Cortisol ( $\mu$ g/dL)	CG	17.27 $\pm$ 3.12	12.97 $\pm$ 3.48	4.30	0.92	4.67	9	0.01*
	EG	13.85 $\pm$ 2.77	11.78 $\pm$ 2.67	2.07	1.15	-1.79	9	0.1
IGF1 (ng/ml)	CG	6.55 $\pm$ 3.04	5.72 $\pm$ 21.82	1.60	45.77	-0.46	9	0.65
	EG	13.85 $\pm$ 2.77	11.78 $\pm$ 2.76	2.07	1.15	-1.79	9	0.10
Aerobic capacity (ml/kg/min)	CG	57.84 $\pm$ 9.71	59.02 $\pm$ 8.74	2.32	0.86	10.33	9	0.00*
	EG	56.85 $\pm$ 8.56	61.42 $\pm$ 10.61	5.43	1.78	6.46	9	0.00*
Anaerobic capacity (wat)	CG	466 $\pm$ 15.77	491 $\pm$ 17.28	-24.2	3.07	-8.13	9	0.04*
	EG	467 $\pm$ 14.94	544 $\pm$ 10.74	-77.87	5.35	-16.46	9	0.00*

\* Significant level at pretest and posttest; CG: control group; EG: Experimental group; RMR, Resting Metabolic Test

The study results also indicated that RMR and testosterone levels increased significantly in both groups. In addition, the cortisol level showed a significant reduction in the CG, but this reduction was not observed in the EG. On the other hand, aerobic capacity significantly increased in both groups, but no significant difference was observed between the two groups. However, compared to the control group, only anaerobic power increased significantly in the experimental group. (Table 3). No significant differences were observed between the groups in other variables.

**Table 3.** Results of independent *t*-test comparing post-test between control group (10 subjects) and exercise group

	Mean difference	F	t	df	Sig.
RMR	-2.97	2.57	-1.47	18	.15
Testosterone (ng/ml)	.48	1.52	.9	18	.38
Cortisol ( $\mu$ g/dL)	-2.23	2.11	-1.51	18	.12
IgF1 (ng/ml)	16.70	3.96	.3	18	.76
Aerobic capacity (ml/kg/min)	2.61	2.64	1.31	18	0.20
Anaerobic capacity (wat)	-62.00	4.93	-9.99	18	0.01 *

#### 4. Discussion

The study results indicated that high-intensity resistance-strength training significantly increased the resting metabolic rate (RMR) in male taekwondo players, although there was no significant difference between two groups. This aligned with Sirithienthad (2006), who reported a 10% increase in RMR within 21 hours after resistance exercises. Additionally, Potteiger et al. (2008) demonstrated that RMR significantly increased following 16 months of aerobic training, supporting the notion that consistent training positively impacts metabolic rates. Furthermore, Nouri et al. (2012) reported a significant increase in RMR in the resistance training group, suggesting that resistance training can effectively boost metabolic efficiency among athletes. The findings suggest that the type and intensity of exercises play a crucial role in influencing RMR. Factors such as heredity, body size, age, gender, and diet also significantly affect RMR, indicating a complex interplay between various elements that influence metabolic rates.

Conversely, several studies present contradictory results regarding RMR. Haghighi et al. (2013) found that resistance training at 75% of maximum repetition did not

significantly change RMR(38), and Hunter et al. (2006) reported that 40 minutes of high-intensity resistance training failed to alter resting energy expenditure (REE) (39). Antunes et al. (2005) further demonstrated a significant reduction in RMR after six months of aerobic training, along with declines in thyroid hormones T3 and T4, which may contribute to decreased RMR(41). These inconsistencies highlight the need for further research to clarify the effects of different training modalities on RMR. Changes in RMR are less influenced by physical activity in athletes compared to sedentary individuals, suggesting that regular exercise may lead to a more stable metabolic rate. Moreover, aerobic exercise has been shown to reduce plasma thyroxine levels, potentially leading to decreased RMR (37).

The investigation into cortisol levels revealed a significant reduction in the control group, but no significant difference between the EG and CG. This finding is consistent with previous studies that reported mixed results regarding exercise's effects on cortisol. For example, (42). Soheili et al. (2024) noted a significant increase in cortisol following moderate-intensity endurance training, while Farzanaki et al. (2008) documented increased cortisol concentrations with higher exercise intensities (43). Kraemer et al. (2008) found similar results in marathon cycling races (44), whereas Reeves et al. (2006) showed no significant change in cortisol levels with resistance training at 70% of maximum repetition (45). These discrepancies highlight that exercise intensity and duration are critical factors influencing cortisol secretion. Intense physical activity can elevate ACTH secretion, leading to increased cortisol levels (39). Sebghati Shiraz et al. in their study, revealed that both sprint and small-size game training individually can have desirable effects on testosterone, cortisol, and T/C ratio in teenage soccer players(46).

The results indicated a significant increase in testosterone levels in both groups following high-intensity resistance and endurance training, but no significant difference between the groups. This finding aligns with previous studies, such as Jabloo et al. (2012), which reported significant changes in testosterone levels after similar training protocols (47). However, West et al. (2009) and Reeves et al. (2006) found no significant changes in testosterone levels under specific training conditions, indicating that various factors—including exercise duration, muscle activity intensity, and individual characteristics—affect hormonal responses. Rigorous training targeting large muscle groups can lead to acute elevations in testosterone, possibly due to adrenaline stimulation and the effects of lactate (49). and his colleagues pointed out that high-intensity training in Taekwondo athletes probably has a positive effect on physical fitness factors, cortisol, and testosterone hormones and improves the performance of these athletes (50).

Investigating IGF-1 levels revealed no significant differences within or between groups. Previous studies, including Irwin et al. (2009), noted a decrease in IGF-1 following moderate-intensity aerobic activities (51). Hasani-Ranjbar et al. (2012) documented an elevation in IGF-1 after rigorous resistance training, suggesting that the type and intensity of exercise significantly influence IGF-1 responses (52). The mechanisms governing IGF-1 secretion involve growth hormone-releasing hormone (GHRH), which stimulates GH secretion and subsequently promotes IGF-1 synthesis in the liver. Various factors, including training type, intensity, and duration, impact IGF-1 levels. Babaei et al.

showed that serum testosterone levels may increase slightly after high-intensity strength training and then high-intensity aerobic training. Serum cortisol levels decrease after high-intensity aerobic training but increase after high-intensity strength training, and it is possible that insulin-like growth factor-1 (Igf1) serum levels decrease after high-intensity strength training(53).

The study findings indicated a significant reduction in aerobic capacity in both groups, while only the CG exhibited a significant increase in anaerobic capacity. This highlights the varying effects of exercise protocols on bioenergetics indicators. Marković et al. (2005) noted that successful taekwondo athletes have higher speed and anaerobic thresholds (54). Melhim et al. (2001) reported that taekwondo training improved anaerobic capacity but did not affect aerobic capacity (55). The demands of taekwondo, characterized by rapid movements and powerful kicks, necessitate a focus on anaerobic capabilities. However, enhancing aerobic fitness is essential for adequate recovery between training sessions and competition intervals (56-58). Contrastingly, Chara et al. (2005) found improvements in maximal oxygen consumption after simultaneous resistance and endurance training in non-athletic men. Nouri et al. (2013) confirmed the positive effects of exercise training on VO<sub>2</sub> max (37). However, Murtag et al. (2005) concluded that physical activity had no significant effect on aerobic capacity (59). These discrepancies may stem from variations in baseline VO<sub>2</sub>max, exercise intensity, and monitoring methods during training (60).

This study faces several limitations, including a small sample size that may affect the generalizability of the results and a lack of control over environmental factors that can influence athletic performance and hormonal responses. Additionally, the absence of psychological impact assessments and the lack of studies on female athletes are notable limitations. To improve future research, it is suggested that studies be conducted with larger sample sizes and standardized training protocols. Furthermore, the simultaneous effects of resistance and aerobic training, as well as the role of diet and supplements, should be investigated. Utilizing advanced technologies for more accurate measurement of hormonal and metabolic changes, along with designing long-term studies, could enhance our understanding of the sustained effects of training on metabolism and hormones.

The findings of this research indicate that high-intensity resistance and strength training significantly enhance the resting metabolic rate (RMR) among male taekwondo players, although no significant differences were observed between the test and control groups. These results align with previous studies that suggest resistance training positively impacts metabolic efficiency. Additionally, while cortisol levels decreased in the CG, no significant changes were found in the EG, highlighting the complex relationship between exercise intensity and hormonal responses. Overall, the study underscores the importance of exercise type and intensity in influencing metabolic rates and hormonal balances, while also identifying the

need for further research to clarify these relationships and their implications for athletic performance.

## 5. Conclusion

In conclusion, the research contributes valuable insights into the effects of high intensity aerobic resistance training can affect metabolic and hormonal responses in male taekwondo players, emphasizing the need for standardized protocols and larger sample sizes in future studies to enhance understanding in this field.

## 6. Acknowledgment

Researchers hereby thank and appreciate all the women for their participation in this research.

**Conflict of interests:** The authors declare that they have no conflict of interest relating to the publication of this manuscript.

## References

1. Bridge CA, Ferreira da Silva Santos J, Chaabene H, Pieter W, Franchini E. Physical and physiological profiles of taekwondo athletes. *Sports Medicine*. 2014;44:713-33. DOI: 10.1007/s40279-014-0135-2
2. Mathunjwa ML, Djarova-Daniels T, Shaw I, Mugandani S, Shaw BS. Short duration high-intensity interval taekwondo training substantially improves body composition and physical fitness in previously-trained individuals: a proof-of-concept study. *Arch Budo*. 2020;16:221-6.
3. Bompa T, Buzzichelli C. *Periodization of Strength Training for Sports*. Human Kinetics Publishers; 2021.
4. Folland JP, Williams AG. Morphological and neurological contributions to increased strength. *Sports Medicine*. 2007;37:145-68. DOI: 10.2165/00007256-200737030-00002
5. Khurana D, Dutta N, Malik S, Sharma S, Minhaj M, Sharma RK, et al. Blood flow restriction therapy with exercise is no better than exercise alone in improving athletic performance, muscle strength, and hypertrophy: A systematic review and meta-analysis. *Somatossensory & Motor Research*. 2024;41(2):97-114. DOI: 10.1080/08990220.2024.2038295
6. Zavanela PM, Crewther BT, Lodo L, Florindo AA, Miyabara EH, Aoki MS. Health and fitness benefits of a resistance training intervention performed in the workplace. *Journal of Sports Science & Medicine*. 2012;26(3):811-7.
7. García-Pinillos F, Lago-Fuentes C, Jaén-Carrillo D, Bujalance-Moreno P, Latorre-Román PÁ, Roche-Seruendo LE, et al. Strength training habits in amateur endurance runners in Spain: Influence of athletic level. *International Journal of Environmental Research and Public Health*. 2020;17(21):8184. DOI: 10.3390/ijerph17218184
8. Lloyd RS, Oliver JL. The youth physical development model: A new approach to long-term athletic development. *Journal of Sports Sciences*. 2012;34(3):61-72. DOI: 10.1080/02640414.2012.725640
9. Delvecchio L, Reaburn PJ. Mixed methods strength training for the masters athlete: A review. *Journal of Australian Strength and Conditioning*. 2013;21(4).
10. Reid M, Schneiker K. Strength and conditioning in tennis: current research and practice. *Journal of Sports Medicine*. 2008;11(3):248-56.
11. Bradley WJ, Cavanagh B, Douglas W, Donovan TF, Twist C, Morton JP, et al. Energy intake and expenditure assessed 'in-season' in an elite European rugby union squad. *Journal of Sports Sciences*. 2015;15(6):469-79. DOI: 10.1080/02640414.2015.1038501
12. Claudino JG, Gabbett TJ, Bourgeois F, Souza HD, Miranda RC, Mezêncio B, et al. CrossFit overview: systematic review and meta-analysis. *Sports*. 2018;4:1-14. DOI: 10.3390/sports4010003
13. Mitchell L, Slater G, Hackett D, Johnson N, O'Connor H. Physiological implications of preparing for a natural male bodybuilding competition. *European Journal of Sport Science*. 2018;18(5):619-29. DOI: 10.1080/17461391.2017.1415730
14. Sperlich B, Hahn L-S, Edel A, Behr T, Helmprobst J, Leppich R, et al. A 4-week intervention involving mobile-based daily 6-minute micro-sessions of functional high-intensity circuit training improves strength and quality of life, but not cardio-respiratory fitness of young untrained adults. *Frontiers in Physiology*. 2018;9:423. DOI: 10.3389/fphys.2018.00423



15. Buresh R, Berg K, French J. The effect of resistive exercise rest interval on hormonal response, strength, and hypertrophy with training. *Journal of Strength & Conditioning Research*. 2009;23(1):62-71. DOI: 10.1519/JSC.0b013e31818f8b47
16. Gharahdaghi N, Phillips BE, Szewczyk NJ, Smith K, Wilkinson DJ, Atherton PJ. Links between testosterone, oestrogen, and the growth hormone/insulin-like growth factor axis and resistance exercise muscle adaptations. *Frontiers in Physiology*. 2021;11:621226. DOI: 10.3389/fphys.2020.621226
17. Jambassi Filho JC, Gurjão ALD, Ceccato M, Prado AKG, Gallo LH, Gobbi SR. Effect of different rest intervals between sets on growth hormone concentrations in trained older women. *Revista Brasileira de Ciência e Movimento*. 2015;17(2):216-25.
18. Schwanbeck SR, Cornish SM, Barss T, Chilibeck PD. Effects of training with free weights versus machines on muscle mass, strength, free testosterone, and free cortisol levels. *The Journal of Strength & Conditioning Research*. 2020;34(7):1851-9. DOI: 10.1519/JSC.0000000000001941
19. Bell G, Syrotuik D, Socha T, Maclean I, Quinney HA. Effect of strength training and concurrent strength and endurance training on strength, testosterone, and cortisol. *The Journal of Strength & Conditioning Research*. 1997;11(1):57-64.
20. Rahimi R, Rohani H, Ebrahimi M. Effects of very short rest periods on testosterone to cortisol ratio during heavy resistance exercise in men. *Apunts Medicina de l'Esport*. 2011;46(171):145-9.
21. Urban RJ. Growth hormone and testosterone: anabolic effects on muscle. *Hormone Research in Paediatrics*. 2011;76(Suppl. 1):81-3. DOI: 10.1159/000331633
22. Kamgang VW, Murkwe M, Wankeu-Nya M. Biological Effects of Cortisol. 2023.
23. Tarpenning KM. Influence of weight training exercise and modification of hormonal response on skeletal muscle growth: University of Southern California; 1997.
24. Brunsden T. Hormonal responses to resistance training and its effects on strength adaptations: A brief overview. *Journal of Medicine*. 2023;10(11).
25. Pruessner KM. Housing Discrimination-Selected Problems in Filing a Housing Complaint with HUD under Section 810 of Title VIII, 42 USC 3610. *University of Kansas Law Review*. 1977;26:255.
26. Sheikholeslami-Vatani D, Ahmadi S, Chehri B, Tadibi V. The effect of changes in concentric-eccentric contraction time ratio on hormonal response to resistance exercise in trained men. *Science & Sports*. 2018;33(3):164-8. DOI: 10.1016/j.scispo.2017.07.005
27. Gryaznykh A, Kiseleva M, Kolokoltsev M, Vorozheikin A, Belykh V, Romanova E, et al. Cortisol and testosterone content variability in blood serum of professional wrestlers under different conditions of physical activity. *Journal of Physical Education and Sport*. 2022;22(2):448-54. DOI: 10.7752/jpes.2022.02055
28. Uchida MC, Bacurau RFP, Navarro F, Pontes Jr FL, Tessuti VD, Moreau RL, et al. Alteration of testosterone: cortisol ratio induced by resistance training in women. *Revista Brasileira de Medicina do Esporte*. 2004;10:165-8.
29. Jansson D, Lindberg A-S, Lundberg E, Domellöf M, Theos A. Effects of resistance and endurance training alone or combined on hormonal adaptations and cytokines in healthy children and adolescents: a systematic review and meta-analysis. *Sports Medicine-Open*. 2022;8(1):81. DOI: 10.1186/s40798-022-00428-2
30. Abasspour Mojdehi A, Shabani R, Fadaei Chafy M. The effect of high-intensity strength and endurance training on body fat index, glucose homeostasis, and serum leptin in taekwondo players aged 15 to 20 years. *Metabolism and Exercise*. 2017;7(1):69-82.
31. Leger LA, Mercier D, Gadoury C, Lambert J. The multistage 20 metre shuttle run test for aerobic fitness. *Journal of Sports Sciences*. 1988;6(2):93-101. DOI: 10.1080/02640418808729800
32. Maciel O, Martins R, Nakamura FY, Figueiredo P, Afonso J, Baptista I. The effect of speed endurance versus core training on the repeated sprint ability of youth male soccer players: a randomized controlled trial. *Journal of Sports Science & Medicine*. 2024;23(4):907.
33. Mifflin MD, St Jeor ST, Hill LA, Scott BJ, Daugherty SA, Koh YO. A new predictive equation for resting energy expenditure in healthy individuals. *The American Journal of Clinical Nutrition*. 1990;51(2):241-7. DOI: 10.1093/ajcn/51.2.241
34. Nieman DC, Austin MD, Benezra L, Pearce S, McInnis T, Unick J, et al. Validation of Cosmed's FitMate™ in measuring oxygen consumption and estimating resting metabolic rate. *Research in Sports Medicine*. 2006;14(2):89-96. DOI: 10.1080/15438620600794996
35. Sirithienthad P. A comparison of the effects of post-exercise basal metabolic rate among continuous aerobic, intermittent aerobic, and resistance exercise: Implications for weight control. The Florida State University. 2006.
36. Potteiger JA, Kirk EP, Jacobsen DJ, Donnelly JE. Changes in resting metabolic rate and substrate oxidation after 16 months of exercise training in overweight adults. *International Journal of Sport Nutrition and Exercise Metabolism*. 2008;18(1):79-95. DOI: 10.1123/ijnsn.18.1.79
37. Nouri Y, Rahmani F, Mirzaie B, Arazi H. The effect of resistance and endurance training on resting metabolic rate and body composition in sedentary males. *Journal of Advances in Medical and Biomedical Research*. 2013;21(89):51-63.
38. Haghighi AH, Mahmoudi M, Delgosha H. Hormonal responses to two programs of exhaustive resistance training of different intensities in male bodybuilders. *Iranian Journal of Endocrinology and Metabolism*. 2012;14(3):267-74.
39. Hunter GR, Byrne NM, Gower BA, Sirikul B, Hills AP. Increased resting energy expenditure after 40 minutes of aerobic but not resistance exercise. *Obesity*. 2006;14(11):2018-25. DOI: 10.1038/oby.2006.236
40. Antunes HK, Santos RF, Boscolo RA, Bueno OF, Mello MT. Analysis of resting metabolic rate and body composition in elderly males before and after six months of endurance exercise. *Revista Brasileira de Medicina do Esporte*. 2005;11:71-5.
41. Antunes PE, Silva R, de Oliveira JF, Antunes MJ. Left ventricular aneurysms: early and long-term results of two types of repair. *European Journal of Cardio-Thoracic Surgery*. 2005;27(2):210-5. DOI: 10.1016/j.ejcts.2004.10.013
42. Soheili S, Sori R, Rezaian N. Hormonal adaptations to moderate-intensity endurance training in sedentary obese men. *Koomesh*. 2024;14(2):181-91.
43. Farzanaki P, Azarbayjani M, Rasaee M, Jourkesh M, Ostojic S, Stannard S. Salivary immunoglobulin A and cortisol response to training in young elite female gymnasts. *Brazilian Journal of Biomotricity*. 2008;2(4):252-8.
44. Kraemer R, Durand R, Acevedo E, Johnson L, Kraemer G, Hebert E, et al. Rigorous running increases growth hormone and insulin-like growth factor-I without altering ghrelin. *Experimental Biology and Medicine*. 2004;229(3):240-6. DOI: 10.1177/153537020422900304
45. Reeves GV, Kraemer RR, Hollander DB, Clavier J, Thomas C, Francois M, et al. Comparison of hormone responses following light resistance exercise with partial vascular occlusion and moderately difficult resistance exercise without occlusion. *Journal of Applied Physiology*. 2006;101(6):1616-22. DOI: 10.1152/jappphysiol.00505.2006
46. Sebhathi Shiraz M, Shabani R, Mohammadi M. Comparison of small-sided games and sprint training program on the testosterone, cortisol, blood cell count, and physical fitness indices in teenage soccer players. 1403.
47. Jabloo DS, Hosseini SRA, Zanjani DS, Ahmadi A. Effects of resistance and endurance exercises on androgens, cortisol, and lactate in elderly women. *Tehran University Medical Journal*. 2012;70(2).
48. West DW, Kujbida GW, Moore DR, Atherton P, Burd NA, Padzik JP, et al. Resistance exercise-induced increases in putative anabolic hormones do not enhance muscle protein synthesis or intracellular signalling in young men. *The Journal of Physiology*. 2009;587(21):5239-47. DOI: 10.1113/jphysiol.2009.179693
49. Sourati Jabloo D, Attarzadeh Hosseini SR, Sayadpour Zanjani D, Ahmadi A, Mansouri J. Comparison of resistance and endurance exercises on testosterone to cortisol ratio in post-menopausal women. *Daneshvar Medicine*. 2020;19(6):69-78.
50. Moravej SR, Nazari M, Shabani R. The effect of high-intensity strength and endurance training on cortisol, testosterone, and physical fitness of 15-20-year-old male taekwondo athletes in Rash. *Journal of Physical Activity and Hormones*. 2024;5(2):-
51. Irwin ML, Varma K, Alvarez-Reeves M, Cadmus L, Wiley A, Chung GG, et al. Randomized controlled trial of aerobic exercise on insulin and insulin-like growth factors in breast cancer survivors: the Yale Exercise and Survivorship study. *Cancer Epidemiology, Biomarkers & Prevention*. 2009;18(1):306-13. DOI: 10.1158/1055-9965.EPI-08-0638
52. Hasani-Ranjbar S, Soleymani Far E, Heshmat R, Rajabi H, Kosari H. Time course responses of serum GH, insulin, IGF-1, IGFBP1, and IGFBP3 concentrations after heavy resistance exercise in trained and untrained men. *Endocrine*. 2012;41(1):144-51. DOI: 10.1007/s12020-011-9497-5
53. Babaei K, Dadmanesh M. Comparison of two methods of high-intensity aerobic and strength training on the levels of testosterone, cortisol, insulin-like growth hormone (IGF1), and immunoglobulin (IgG) in professional taekwondo athletes. 1403.

54. Marković G, Misigoj-Duraković M, Trinić S. Fitness profile of elite Croatian female taekwondo athletes. *Collegium Anthropologicum*. 2005;29(1):93-9.
55. Melhim A. Aerobic and anaerobic power responses to the practice of taekwondo. *British Journal of Sports Medicine*. 2001;35(4):231-4. DOI: 10.1136/bjsm.35.4.231
56. Gaeini AA, Mahmoudi Y-o-l, Moradian K, Fallahi AA. The relationship between anthropometric, physiological, and body composition characteristics of male elite taekwondo athletes with their success. *Journal of Sport Biosciences*. 2010;2(4):5-20.
57. Arazı H, Hosseinzadeh Z, Izadı M. Relationship between anthropometric, physiological, and physical characteristics with success of female taekwondo athletes. *Turkish Journal of Sport and Exercise*. 2016;18(2):69-75.
58. Can E, Kutlay E, Quintana MS, Bridge CA. Anthropometric characteristics of elite male taekwondo athletes according to weight category and performance level. *Scientific Journal of Sport and Performance*. 2023;2(1):16-27.
59. Murtagh EM, Boreham CA, Nevill A, Hare LG, Murphy MH. The effects of 60 minutes of brisk walking per week, accumulated in two different patterns, on cardiovascular risk. *Preventive Medicine*. 2005;41(1):92-7. DOI: 10.1016/j.ypmed.2004.12.006
60. Schmidt WD, Biwer CJ, Kalscheuer LK. Effects of long versus short bout exercise on fitness and weight loss in overweight females. *Journal of the American College of Nutrition*. 2001;20(5):494-501. DOI: 10.1080/07315724.2001.10719201