

Dietary Supplementation of Zinc Oxide Nanoparticles and Yeast Culture: Effect on Nutrient Digestibility, Growth Performance, and Blood Metabolites in Ghanduzi Male Lambs

Research Article

E. Ibrahimi Khoram Abadi^{1*}¹ Department of Animal Sciences, Faculty of Agriculture and Animal Science, University of Torbat-e Jam, Torbat-e Jam, Iran

Received on: 2 Aug 2024

Revised on: 11 Oct 2024

Accepted on: 15 Nov 2024

Online Published on: Dec 2024

*Correspondence E-mail: el.ebrahimi@tjamcaas.ac.ir

© 2010 Copyright by Islamic Azad University, Rasht Branch, Rasht, Iran

Online version is available on: www.ijas.ir

ABSTRACT

The worldwide prohibition of antibiotics has heightened interest in alternative feed additives, like paraprobiotics and nanominerals, to improve livestock performance. This study investigates the effects of zinc oxide nanoparticles (ZnO-Nps) and yeast culture (YC) on nutrient digestibility, growth performance, and blood metabolites in Ghanduzi male lambs. Sixteen Ghanduzi male lambs, with an average body weight of 32 ± 4.2 kg, were randomly assigned to one of four dietary treatments: a control diet without supplements; a diet supplemented with 50 mg/kg ZnO-Nps; a diet containing 5 g/kg YC; and a diet combining both ZnO-Nps and YC. The trial lasted 28 days, during which daily dry matter intake (DMI), nutrient digestibility, and growth performance metrics were recorded. Blood samples were collected for biochemical analysis. Lambs fed with the diet combining ZnO-Nps and YC showed the highest daily dry matter intake (DMI). This diet significantly enhanced the digestibility of dry matter (DM), crude protein (CP), neutral detergent fiber (NDF), and organic matter (OM) compared to other groups ($P < 0.001$). The final body weight (BW) and average daily gain (ADG) were also significantly higher in the ZnO-Nps + YC group (43.42 kg and 0.417 kg/day, respectively) compared to the control and YC groups ($P < 0.001$). Blood metabolites showed no significant differences among treatment groups, although trends indicated improved nitrogen utilization in the supplemented groups. Combining ZnO-Nps and yeast culture substantially enhances growth performance and nutrient digestibility in Ghanduzi male lambs without adversely affecting blood metabolites. These results advocate for the utilization of ZnO-Nps and YC as potent alternatives to conventional growth promoters in sheep farming.

KEY WORDS feed additives, nanominerals, paraprobiotics, sheep farming.

INTRODUCTION

Since antibiotics were prohibited worldwide, farmers have been increasingly using other substances in sheep feed to help their animals grow faster (Valenzuela-Grijalva *et al.* 2017; He *et al.* 2020). Paraprobiotics represent an alternative feed additive employed in sheep farming systems, serving as replacements for traditional growth-promoting agents (Liu *et al.* 2019; Plaza-Díaz *et al.* 2019; Direkvandi *et al.*

2020). Paraprobiotics, which are characterized by non-viable microbial cells that offer health benefits, can potentially enhance the host's well-being. Yeast culture, a type of paraprobiotic, is a natural product obtained through yeast fermentation (Liu *et al.* 2019). Yeast culture comprises dead yeast cells, nutrients they used to grow, and substances they produced while alive. These substances include proteins, sugars, fats, vitamins, small proteins, amino acids, nucleotides, organic acids, and substances that pro-

fect cells from damage. Supplementing ruminants' diets with paraprobiotics, such as YC, offers multiple benefits: it has antimicrobial effects, inhibits the growth of enteric pathogens, regulates rumen pH, boosts feed intake and digestion, enhances the production of volatile fatty acids (VFA), and optimizes rumen function. These benefits collectively lead to improved animal performance (Tripathi and Karim, 2011; Zhang *et al.* 2016; Dias *et al.* 2018; Nawab *et al.* 2019; Mitchell and Heinrichs, 2020).

A new approach in livestock farming involves using nanominerals, particularly zinc (Zn), to integrate these essential elements into the animals' bodies. This technique enables targeted delivery of active compounds to specific organs, minimizing their rapid breakdown and enhancing health benefits (Fesseha *et al.* 2020; Reddy *et al.* 2020). Research suggests that the use of nanominerals in animal production could significantly enhance both immunity and reproduction. Zinc oxide nanoparticles provide a superior surface-to-volume ratio compared to larger ZnO particles or zinc methionine supplements and can be used in small quantities, yielding better outcomes than conventional zinc sources (Geetha *et al.* 2020; Hussein *et al.* 2020). Furthermore, ZnO-Nps not only enhance productivity but also serve as antibacterial agents, positively impacting animal immunity and breeding (Hussein *et al.* 2020).

Feed additives can significantly enhance livestock health and performance. Antioxidants like vitamin E, C, proanthocyanidins, and curcumin, along with trace elements like selenium and zinc, can reduce oxidative stress and improve overall well-being (Sun *et al.* 2023). Additionally, probiotics and prebiotics have been shown to promote growth and strengthen the gastrointestinal immune system (Garcia *et al.* 2019). Studies suggest that combining different feed additives often yields better results than using them individually (Sun *et al.* 2023). For instance, Zn-enriched yeast has been shown to benefit lambs by reducing intestinal damage and improving overall performance (NRC, 2007). However, the effects of combining zinc oxide nanoparticles (ZnO-Nps) with yeast culture on fattening lambs are less well-understood. The study focused on examining how adding zinc oxide nanoparticles, yeast culture, or a combination of both to the diet of male Ghanduzi lambs affected their ability to digest nutrients, their growth, and their blood chemistry.

MATERIALS AND METHODS

Experimental feed additives

The ZnO-Nps were purchased from Nano-Scale Borhan, a knowledge-based company in Mashhad, Iran. The YC was obtained from Kavoshbiotech company in Tehran, Iran. Figure 1 shows a scanning electron microscopy (SEM) im-

age and the corresponding energy dispersive X-ray (EDX) spectrum of the ZnO-Nps used in this study.

Animals, diet, and treatments

The research was carried out at the Torbat-e Jam livestock farm in Iran. The lambs were housed, according to the Iranian Council on Animal Care (1995). Sixteen Ghanduzi male lambs (32±4.2 kg body weight) were randomly divided into four groups and fed different experimental diets. The diets included a control diet without supplements, a diet with ZnO-Nps (50 mg/kg), a diet with YC (5 g/kg), and a diet with both ZnO-Nps and YC (50 mg/kg and 5 g/kg, respectively).

The diets were formulated based on the NRC (2007) recommendations for a forage-to-concentrate ratio of 15:85. Before the main trial, the lambs were allowed to adjust to their new diets for 20 days. The trial then lasted for 28 days. The lambs were housed individually (2 m×2 m) (Iranian Council on Animal Care, 1995) and provided with unlimited feed (2% refusals, twice daily: 7 a.m. and 7 p.m.) and fresh water.

Animal sampling

Daily DMI and orts were recorded daily. Feed samples were dried in a forced-air oven (Behdad Co., Iran) at 65 °C for 48 hours and stored in plastic bags for analysis. Lambs were weighed daily before morning feeding and weekly thereafter. Fresh fecal samples were collected one hour after feeding for seven days (days 41-48) using leather bags. Nutrient digestibility was assessed by measuring DMI and refusals. Lambs were given a two-day adjustment period to the fecal collection bags. Bags were emptied completely twice daily. Collected feces were thoroughly mixed, weighed, and a portion was stored at -20 °C for later analysis. Fecal samples were dried in a forced-air oven at 65 °C for 48 hours and then ground to a particle size of 1 mm using a Wiley mill before analysis. Nutrient digestibility was calculated based on nutrient intake and excretion data, using the analytical procedures described in the analytical procedures section. Blood samples were collected over five consecutive days during week four, three hours after the morning feeding, using ethylenediaminetetraacetic acid (EDTA) tubes through the jugular vein (5-10 mL). The samples were centrifuged at 3000 × g for 10 minutes, and the plasma supernatant was transferred into sterile 1.5 mL micro-tubes, stored at -80 °C for subsequent analysis.

Analytical procedures

The AOAC protocol (AOAC, 2005) was used to analyze the following: of dry matter (method 930.15), ether extract (EE, method 991.36), ash (method 942.05), and crude protein (CP), (Kjeldahl method, N × 6.25, method 954.01).




Figure 1 The company's SEM image and EDX spectrum about the purchased ZnO-Nps

For the determination of neutral detergent fiber (NDF), the procedure described by Van Soest *et al.* (1991) and the protocol of Ankom Technology (2006) was utilized, with the analysis conducted using the Ankom fiber analyzer (ANKOM, model A2001, New York, USA). The levels of total protein (T.P), albumin (Alb), globulin (Glo), blood urea nitrogen (BUN), creatinine (Cre), glucose (Glu), triglyceride (TG), cholesterol (Chol), alanine transaminase (ALT), aspartate transferase (AST), creatine kinase (CPK), gamma-glutamyltransferase (GGT), and alkaline phosphatase (ALP) were measured using an auto-analyzer (Mindray BS-200 Chemistry Analyzer, China).

Statistical analysis

All data were analyzed using the PROC GLM procedure in SAS (2013) with the following model:

$$Y_{ij} = \mu + T_i + e_{ij}$$

Where:

Y_{ij} : value of each observation.

μ : denotes the overall mean.

T_i : signifies the treatment effect.

e_{ij} : experimental error.

The statistical significance of the treatment differences was assessed using the Tukey test ($P < 0.05$).

RESULTS AND DISCUSSION

Dry matter intake and nutrient digestibility

Table 2 presents the effects of individually or combined dietary supplementation of ZnO-Nps with YC on DMI and nutrient digestibility in Ghanduzi male lambs.

Table 1 Major ingredients and chemical composition of the experimental basal diet (%)

| | |
|--|-------|
| Alfalfa hay | 10.00 |
| Wheat straw | 5.00 |
| Barley grain | 40.00 |
| Corn grain | 29.80 |
| Wheat bran | 12.00 |
| Soybean meal | 10.00 |
| Vitamins and minerals mixture ¹ | 5.00 |
| Urea | 2.50 |
| Fat powder | 0.70 |
| Chemical composition | |
| Dry matter (% of fresh weight) | 90.00 |
| Metabolizable energy (Mcal/kg DM) | 2.83 |
| Crude protein (%) | 14.85 |
| Neutral detergent fiber (%) | 26.12 |
| Non-fiber carbohydrates (%) | 48.06 |
| Ether extract (%) | 3.80 |
| Ash (%) | 7.17 |

Mineral and vitamin mixture (per kg): vitamin E: 1000 IU; vitamin A: 100,000 IU; vitamin D₃: 40,000 IU; Ca: 147 g; P: 5 g; Mg: 20 g; Na: 40 g; S: 4 g; Cu: 100 mg; I: 20 mg; Co: 20 mg; Mn: 800 mg; Zn: 800 mg; Fe: 400 mg; Se: 20 mg and Antioxidant: 1000 mg.

The findings demonstrate that the group receiving ZnO-Nps and YC had the highest DMI compared to the control and individual supplementation groups ($P < 0.001$). Additionally, lambs fed a diet containing ZnO-Nps exhibited significantly higher nutrient digestibility for DM, CP, NDF, and organic matter (OM) compared to the control and YC groups ($P < 0.001$). Including ZnO-Nps at levels of 100 and 200 mg/kg has been shown to enhance VFAs production, microbial CP, and OM breakdown during the 6th and 12th hours of an *in vitro* fermentation study (Chen *et al.* 2011). Similarly, in an *in vitro* fermentation study, an increase in microbial biomass production and a reduction in methane emissions were observed when 20 mg of zinc in the form of

ZnNps was administered, compared to higher zinc levels (40 and 60 mg/kg dry matter) (Riyazi *et al.* 2019). Incorporating ZnNps into lamb diets improved diet digestibility and feeding value, as evidenced by increased feed utilization efficiency in the experimental dietary treatments (Mohamed *et al.* 2015). Positive results have been observed in Holstein calves. Supplementation of 50 mg Zn/kg DM, either as ZnO or ZnO-Nps, has led to an increase in DM digestibility (Rajendran *et al.* 2013). The beneficial impacts of ZnO-Nps on DMI and nutrient digestibility can be attributed to their unique properties. These include a large surface area-to-volume ratio, nanoscale size, rapid and targeted movement, and catalytic efficiency. These properties enhance the bioavailability of zinc for absorption in the gastrointestinal tract (GIT) as noted by (Zhan *et al.* 2007; Seven *et al.* 2018). The growth-promoting effects of nanominerals are likely due to their ability to positively influence the gut's structure and function. They can modify the gut microarchitecture in animals (Adegbeye *et al.* 2019) and improve rumen fermentation, particularly in terms of fiber digestion and redox balance (Murthy, 2013; Zhai *et al.* 2017).

Yeast culture, a safe and environmentally friendly dietary supplement, has been shown to enhance nutrient digestibility in sheep (Wang *et al.* 2022). Studies by Malekkhahi *et al.* (2015) and Song *et al.* (2021) found that adding YC to high-concentrate diets significantly improved the digestibility of both CP and NDF. Research on beef cattle and dairy cows by Lei *et al.* (2013) and Jiang *et al.* (2017) also revealed that YC can increase fiber digestibility. This consistent improvement is likely due to an increase in cellulose-degrading bacteria or the presence of metabolites in YC that stimulate these bacteria. The positive impact of YC on fiber digestibility is particularly beneficial for ruminants fed high-concentrate diets, as shown by Dias *et al.* (2018). This may explain the higher dry matter intake (DMI) and nutrient digestibility observed in the ZnO-Nps + YC group compared to the other groups in the study.

Growth performance

Initial body weights were comparable across all treatments ($P=0.080$) (Table 3). However, significant differences emerged in final body weights ($P=0.006$) (Table 3). The ZnO-Nps + YC treatment group demonstrated the highest final body weight, followed by the ZnO-Nps group. Moreover, the ZnO-Nps + YC group exhibited notably higher average daily gains (ADG) on days 14 ($P=0.050$) and 28 ($P=0.003$) compared to the control and YC groups (Table 3). The cumulative ADG and total body weight gains (BWG) were significantly superior in the ZnO-Nps + YC group ($P<0.001$) (Table 3). Additionally, the feed conversion ratio (FCR) was markedly improved in the ZnO-Nps +

YC group relative to the control and YC groups ($P=0.007$) (Table 3).

Studies have shown that supplementing livestock with ZnO-Nps can lead to significant improvements in growth and gut health. Anil *et al.* (2019) found that calves given 10 mg/kg of ZnO-Nps experienced substantial increases in total and live weight. Similarly, Zaboli *et al.* (2013) observed that kids supplemented with 20-40 ppm of ZnO-Nps had significantly higher average daily gains, compared to the control group. Additionally, ZnO supplementation at lower doses (below 50-100 mg/kg of dry matter) has been shown to increase the overall number of bacteria in the gut, including beneficial intestinal bacteria. This, in turn, leads to a significant enhancement in VFAs, which play a crucial role in gut health and nutrient absorption (Pieper *et al.* 2012). Garg *et al.* (2008) further demonstrated that zinc supplements can improve food conversion efficiency by making zinc more readily available in the digestive system. This increased zinc availability contributes to a higher bacterial count and better utilization of ingested feed.

Studies on yeast products have shown benefits for livestock, including: increased feed efficiency, improved daily weight gain, enhanced overall animal health and well-being (Kim *et al.* 2000; Haddad and Goussous, 2005; Rozeboom *et al.* 2005; Ding *et al.* 2008).

Research has also demonstrated that supplementing ruminants with yeast products can positively influence DMI, rumen pH, and overall nutrient digestibility (Callaway and Martin, 1997; Dann *et al.* 2000). Yeast and yeast products may enhance animal performance by supporting animal health through positive effects on bacterial colonization in the GIT, and potentially increasing nutrient digestibility (Harris and Webb, 1990). The impact of dietary treatments on blood metabolites in Ghanduzi male lambs is summarized in Table 4. While no significant differences were found among the various treatment groups, certain trends emerged, particularly in BUN levels. Lambs fed the ZnO-Nps + YC diet exhibited the highest BUN levels, followed by those receiving ZnO-Nps alone.

Table 5 outlines the effects of individual or combined dietary supplementation of ZnO-Nps with YC on liver enzymes and creatine phosphokinase in Ghanduzi male lambs. Although ALT, AST, CPK, GGT, and ALP were unaffected by the inclusion of ZnO-Nps and YC in the basal diet, some trends were observed, especially in GGT and ALP levels. Blood biochemistry values are crucial to the body's physiological state (El-Kholy *et al.* 2017). Previous research has indicated that adding ZnO-Nps to ruminant diets doesn't significantly impact specific biochemical parameters (Zaboli *et al.* 2013; Swain *et al.* 2019; Abd El Rahim *et al.* 2023).

Table 2 Effects of individual or combined dietary supplementation of zinc oxide nanoparticles (ZnO-Nps) with yeast culture (YC) on dry matter intake and nutrient digestibility

| Items | Treatments | | | | SEM | P-value |
|------------------------|-------------------|--------------------|--------------------|-------------------|-------|---------|
| | Control | ZnO-Nps | YC | ZnO-Nps + YC | | |
| DMI (kg/day) | 1.25 ^a | 1.51 ^b | 1.36 ^c | 1.56 ^d | 0.024 | < 0.001 |
| Nutrient digestibility | | | | | | |
| DM | 0.71 ^a | 0.75 ^{bc} | 0.72 ^{ab} | 0.76 ^c | 0.005 | < 0.001 |
| CP | 0.73 ^a | 0.75 ^{bc} | 0.74 ^{ab} | 0.78 ^c | 0.005 | < 0.001 |
| NDF | 0.55 ^a | 0.62 ^{bc} | 0.58 ^{ab} | 0.64 ^c | 0.008 | < 0.001 |
| OM | 0.73 ^a | 0.77 ^{bc} | 0.75 ^{ab} | 0.78 ^c | 0.005 | < 0.001 |

DMI: dry matter intake; CP: crude protein; NDF: neutral detergent fiber and OM: organic matter.

Control: basal diet without supplements; ZnO-Nps: 50 mg/kg DM basal diet; YC: 5 g/kg DM basal diet and ZnO-Nps + YC: 50 mg and 5g/kg DM basal diet, respectively. The means within the same row with at least one common letter, do not have significant difference (P>0.05).

SEM: standard error of the means.

Table 3 Effects of individual or combined dietary supplementation of zinc oxide nanoparticles (ZnO-Nps) with yeast culture (YC) on growth performance

| Items | Treatments | | | | SEM | P-value |
|---------------------------------|-------------------|--------------------|-------------------|-------------------|------|---------|
| | Control | ZnO-Nps | YC | ZnO-Nps + YC | | |
| Initial BW (kg) | 27.9 | 35.2 | 32.1 | 32.5 | 1.05 | 0.080 |
| Final BW (kg) | 33.9 ^a | 44.1 ^{ab} | 39.4 ^a | 43.4 ^b | 1.31 | 0.006 |
| Average daily gain (ADG) (kg/d) | | | | | | |
| Day 07 | 0.2 | 0.3 | 0.3 | 0.2 | 0.02 | 0.200 |
| Day 14 | 0.2 ^a | 0.3 ^{bc} | 0.3 ^{ab} | 0.4 ^b | 0.02 | 0.050 |
| Day 21 | 0.1 | 0.2 | 0.1 | 0.2 | 0.28 | 0.670 |
| Day 28 | 0.1 ^a | 0.3 ^{bc} | 0.2 ^{ab} | 0.4 ^c | 0.41 | 0.003 |
| Total ADG | 0.2 ^a | 0.3 ^b | 0.2 ^{ab} | 0.3 ^c | 0.01 | < 0.001 |
| Total BWG (kg) | 6.0 ^a | 8.8 ^b | 7.3 ^{ab} | 10.8 ^c | 0.49 | < 0.001 |
| FCR | 7.6 ^a | 5.2 ^b | 7.1 ^a | 4.6 ^b | 0.40 | 0.007 |

BW: body weight; BWG: body weight gains and FCR: feed conversion ratio.

Control: basal diet without supplements; ZnO-Nps: 50 mg/kg DM basal diet; YC: 5 g/kg DM basal diet and ZnO-Nps + YC: 50 mg and 5g/kg DM basal diet, respectively. The means within the same row with at least one common letter, do not have significant difference (P>0.05).

SEM: standard error of the means.

Table 4 Effects of individual or combined dietary supplementation of zinc oxide nanoparticles (ZnO-Nps) with yeast culture (YC) on blood metabolites

| Items | Treatments | | | | SEM | P-value |
|--------------|------------|---------|------|--------------|------|---------|
| | Control | ZnO-Nps | YC | ZnO-Nps + YC | | |
| T.P (g/dL) | 8.0 | 8.5 | 7.6 | 9.1 | 0.37 | 0.57 |
| Alb (g/dL) | 2.4 | 2.3 | 2.4 | 2.1 | 0.05 | 0.30 |
| Glo (g/dL) | 5.5 | 7.1 | 5.3 | 7.8 | 0.62 | 0.48 |
| Alb/Glo | 0.4 | 0.3 | 0.4 | 0.3 | 0.02 | 0.15 |
| Cre (mg/dL) | 0.7 | 0.6 | 0.7 | 0.6 | 0.03 | 0.84 |
| Glu (mg/dL) | 99.0 | 88.3 | 102. | 107.6 | 3.52 | 0.27 |
| TG (mg/dL) | 28.6 | 31.3 | 19.3 | 28.3 | 2.20 | 0.24 |
| Chol (mg/dL) | 53.0 | 56.3 | 64.3 | 69.6 | 3.84 | 0.45 |
| BUN (mg/dL) | 6.3 | 11.0 | 7.0 | 13.6 | 1.16 | 0.06 |

T.P: total protein; Alb: albumin; Glo: globulin; Cre: creatinine; Glu: glucose; TG: triglyceride; Chol: cholesterol and BUN: blood urea nitrogen.

Control: basal diet without supplements; ZnO-Nps: 50 mg/kg DM basal diet; YC: 5 g/kg DM basal diet and ZnO-Nps + YC: 50 mg and 5g/kg DM basal diet, respectively. The means within the same row with at least one common letter, do not have significant difference (P>0.05).

SEM: standard error of the means.

Table 5 Effects of individual or combined dietary supplementation of zinc oxide nanoparticles (ZnO-Nps) with yeast culture (YC) on alanine transaminase, aspartate transaminase, creatine phosphokinase, gamma glutamyltransferase, and alkaline phosphatase

| Items | Treatments | | | | SEM | P-value |
|-----------|------------|---------|-------|--------------|------|---------|
| | Control | ZnO-Nps | YC | ZnO-Nps + YC | | |
| ALT (U/L) | 55.0 | 51.6 | 51.6 | 55.3 | 1.99 | 0.89 |
| AST (U/L) | 132.3 | 121.0 | 130.3 | 114.0 | 6.59 | 0.79 |
| CPK (U/L) | 488.6 | 307.6 | 413.0 | 303.0 | 42.4 | 0.38 |
| GGT (U/L) | 84.3 | 40.0 | 62.3 | 41.3 | 7.33 | 0.08 |
| ALP (U/L) | 95.6 | 22.0 | 69.1 | 28.9 | 11.8 | 0.06 |

ALT: alanine transaminase; AST: aspartate transaminase; CPK: creatine phosphokinase; GGT: gamma glutamyltransferase and ALP: alkaline phosphatase.

Control: basal diet without supplements; ZnO-Nps: 50 mg/kg DM basal diet; YC: 5 g/kg DM basal diet and ZnO-Nps + YC: 50 mg and 5g/kg DM basal diet, respectively. The means within the same row with at least one common letter, do not have significant difference (P>0.05).

SEM: standard error of the means.

However, the results in Table 4 align with a study by [Belewu and Adewumi \(2021\)](#), which found that West African Dwarf (WAD) goats fed diets with 40 and 80 mg/kg of ZnO-Nps experienced a notable decrease in BUN levels compared to the control group. Additionally, [Novoselec et al. \(2017\)](#) observed that incorporating ZnO-Nps, ZnO, and Zn-Met at 28 mg/kg DM in sheep diets led to lower serum BUN levels. The higher plasma BUN concentrations in the ZnO-Nps + YC group might be due to improved diet digestibility and increased N intake.

Fluctuations in AST and ALT blood levels can signal ZnO-Nps toxicity ([Wang et al. 2017](#)), potentially leading to liver damage ([Kurtz and Travlos, 2017](#)). However, the similar AST and ALT levels in the lambs suggest that including ZnO-Nps in their diet did not cause any liver cell injury. This finding aligns with previous research on sheep fed ZnO-Nps ([Mohamed et al. 2017](#); [Swain et al. 2019](#)), or alternative zinc sources like Zn-methionine, Zn glycine, and Zn-protein (Alimohamady et al. 2019).

The addition of YC did not affect serum metabolite level. These findings are consistent with previous research on lambs ([Malekkhahi et al. 2015](#); [Mahmoud et al. 2020](#)) and dairy cows ([Dias et al. 2018](#)). Blood urea nitrogen is a by-product of protein metabolism and indicates the amount of dietary protein and its utilization by the body. Increased DMI typically leads to higher CP intake. Excess protein is metabolized, producing urea, which is excreted in urine, raising BUN levels ([Song et al. 2021](#)). Liver function parameters ALT, AST, and CPK were unaffected by YC in this study. Previous research by [Abou Elenin et al. \(2016\)](#) and [Song et al. \(2021\)](#) suggest that YC has no adverse or beneficial effects on liver metabolism.

In the present study, ZnO-Nps treatment was found to lower GGT and ALP levels in livestock compared to other groups. Livestock can experience various stressors throughout their lives, including disease exposure, environmental challenges, handling stress, nutritional problems, relocation, and new surroundings. These stressors can activate the hypothalamic-pituitary-adrenal (HPA) axis and potentially the immune system, leading to decreased performance ([Burdick et al. 2011](#)). Probiotics, like yeast and yeast-derived products, can boost immunity and health by modifying the gut microbiome ([White et al. 2002](#); [Mullins et al. 2013](#); [AlZahal et al. 2014](#)). A healthy gut flora improves nutrient absorption and immune function ([Bauer et al. 2006](#); [Czarnecki-Maulden, 2008](#)). Additionally, yeast cell wall polysaccharides may prevent harmful bacteria from attaching to the gut, reducing the risk of infections ([Bäumler et al. 1997](#)).

CONCLUSION

Combining zinc oxide nanoparticles and yeast culture significantly boosted growth performance and nutrient digestibility in Ghanduzi male lambs without negatively impacting blood metabolites. These results suggest that ZnO-NPs and YC could be effective alternatives to traditional growth promoters in sheep farming. Future research should focus on long-term effects of ZnO-NPs and YC on lamb health and performance, determining the optimal dosages of these supplements, understanding the underlying mechanisms that contribute to the observed benefits, and assessing the implications for animal health and welfare within livestock production systems.

ACKNOWLEDGEMENT

We thank the University of Torbat-e Jam for providing financial and technical support for this project. We also acknowledge Mr. Yousef Saeedi for his valuable contributions and assistance throughout this study.

REFERENCES

- Abd El Rahim S.A., Arafa M.M., Abdel Hamid H.Y. and Mohamed A.E.A. (2023). Effect of Zinc Oxide Nanoparticles on some biochemical parameters and body weight in Barki fattening lambs. *SVU- Int. J. Vet. Sci.* **6(2)**, 88-103.
- Abou Elenin E.I.M., Abd El-Galil E.R., Etman K.E.I. and El-Shabrawy H.M. (2016). Improvement of rumen fermentation and performance of growing lambs by adding natural microbial resources. *Asian J. Anim. Sci.* **10**, 202-212.
- Adegbeye M.J., Elghandour M.M.M.Y., Barbabosa-Pliego A., Monroy J.C., Mellado M., Ravi K.R.P. and Salem A.Z.M. (2019). Nanoparticles in equine nutrition: Mechanism of action and application as feed additives. *J. Equine Vet. Sci.* **78**, 29-37.
- Alimohamady R., Aliarabi H., Bruckmaier R.M. and Christensen R.G. (2019). Effect of different sources of supplemental zinc on performance, nutrient digestibility, and antioxidant enzyme activities in lambs. *Biol. Trace Elem. Res.* **189**, 75-84.
- AlZahal O., Dionissopoulos L., Laarman A., Walker N. and McBride B. (2014). Active dry *Saccharomyces cerevisiae* can alleviate the effect of subacute ruminal acidosis in lactating dairy cows. *J. Dairy Sci.* **97**, 7751-7763.
- Anil T.S.V., Venkata Seshaiiah C., Ashalatha P. and Sudhakar K. (2019). Effect of dietary supplementation of nano zinc oxide (nZnO) on growth performance in crossbred calves. *Int. J. Curr. Microbiol. Appl. Sci.* **8(12)**, 1852-1856.
- Ankom Technology. (2006). Neutral Detergent Fiber in Feeds-filter Bag Technique. Available at:

- https://www.ankom.com/sites/default/files/document-files/Method_6_NDF_A200.pdf
- AOAC. (2005). Official Methods of Analysis. 18th Ed. Association of Official Analytical Chemists, Gaithersburg, MD, USA.
- Bauer E., Williams B.A., Smidt H., Verstegen M.W. and Mosenthin R. (2006). Influence of the gastrointestinal microbiota on development of the immune system in young animals. *Curr. Issues Intest. Microbiology*. **7**, 35-52.
- Bäumler A., Tsohis R. and Heffron F. (1997). Mechanisms in the Pathogenesis of Enteric Diseases. Springer, Berlin, Germany.
- Belewu A. and Adewumi D. (2021). Effect of green synthesized nano zinc oxide on performance characteristics and hemato-biochemical profile of West African Dwarf goats. *Anim. Res. Int.* **18(1)**, 3938-3946.
- Burdick N.C., Randel R.D., Carroll J.A. and Welsh T.H. (2011). Interactions between temperament, stress, and immune function in cattle. *Int. J. Zool.* **2011**, 1-10.
- Callaway E.S. and Martin S.A. (1997). Effects of a *Saccharomyces cerevisiae* culture on ruminal bacteria that utilize lactate and digest cellulose. *J. Dairy Sci.* **80**, 2035-2044.
- Chen J., Wei W. and ZhiSheng W. (2011). Effect of nano-zinc oxide supplementation on rumen fermentation *in vitro*. *Chin. J. Anim. Nutr.* **23**, 1415-1421.
- Czarnecki-Maulden G. (2008). Effect of dietary modulation of intestinal microbiota on reproduction and early growth. *Theriogenology*. **70**, 286-290.
- Dann H., Drackley J., McCoy G., Hutjens M. and Garrett J. (2000). Effects of yeast culture (*Saccharomyces cerevisiae*) on prepartum intake and postpartum intake and milk production of Jersey cows. *J. Dairy Sci.* **83**, 123-127.
- Dias J.D.L., Silva R.B., Fernandes T., Barbosa E.F., Graças L.E.C., Araujo R.C., Pereira R.A.N. and Pereira M.N. (2018). Yeast culture increased plasma niacin concentration, evaporative heat loss, and feed efficiency of dairy cows in a hot environment. *J. Dairy Sci.* **101**, 5924-5936.
- Ding J., Zhou Z.M., Ren L.P. and Meng Q.X. (2008). Effect of monensin and live yeast supplementation on growth performance, nutrient digestibility, carcass characteristics and ruminal fermentation parameters in lambs fed steam-flaked corn-based diets. *Asian-Australasian J. Anim. Sci.* **21**, 547-554.
- Direkvandi E., Mohammadabadi T. and Salem A.Z.M. (2020). Oral administration of lactate producing bacteria alone or combined with *Saccharomyces cerevisiae* and *Megasphaera elsdenii* on performance of fattening lambs. *J. Appl. Anim. Res.* **48**, 235-243.
- El-Kholy M.S., El-Hindawy M.M., Alagawany M., Abd El-Hack M.E. and ElSayed S.A.A. (2017). Dietary supplementation of chromium can alleviate the negative impacts of heat stress on performance, carcass yield, and some blood hematology and chemistry indices of growing Japanese quail. *Biol. Trace Elem. Res.* **179(1)**, 148-157.
- Fesseha H., Degu T. and Getachew Y. (2020). Nanotechnology and its application in animal production: A review. *Vet. Med. Open J.* **52**, 43-50.
- Garcia G.R., Dogi C.A. and Poloni V.L. (2019). Beneficial effects of *Saccharomyces cerevisiae* RC016 in weaned piglets: *In vivo* and *ex vivo* analysis. *Benef. Microbes.* **10**, 33-42.
- Garg A.K., Mudgal V. and Dass R.S. (2008). Effect of organic zinc supplementation on growth, nutrient utilization and mineral profile in lambs. *Anim. Feed Sci. Technol.* **144**, 82-96.
- Geetha K., Chellapandian M., Arulnathan N. and Ramanathan A. (2020). Nano zinc oxide - An alternate zinc supplement for livestock. *Vet. World.* **13(1)**, 121-126.
- Haddad S.G. and Goussous S.N. (2005). Effect of yeast culture supplementation on nutrient intake, digestibility, and growth performance of Awassi lambs. *Anim. Feed Sci. Technol.* **118**, 343-348.
- Harris B.J. Jr. and Webb D. (1990). The effect of feeding a concentrated yeast culture product to lactating dairy cows. *J. Dairy Sci.* **73**, 1-10.
- He Y., Yuan Q., Mathieu J., Stadler L., Senehi N., Sun R. and Alvarez P.J. (2020). Antibiotic resistance genes from livestock waste: Occurrence, dissemination, and treatment. *Npj Clean Water.* **3**, 4-11.
- Hussein M.A., Ismail Z.S.H. and Abdel Wareth A.A.A. (2020). Application of zinc oxide nanoparticles on productive performance in rabbit nutrition: A Review. *SVU- Int. J. Vet. Sci.* **2(2)**, 278-290.
- Iranian Council of Animal Care. (1995). Guide to the Care and Use of Experimental Animals. Isfahan University of Technology, Isfahan, Iran.
- Jiang Y., Ogunade I.M., Arriola K.G., Qi M., Vyas D., Staples C.R. and Adesogan A.T. (2017). Effects of the dose and viability of *Saccharomyces cerevisiae*. 2. Ruminal fermentation, performance of lactating dairy cows, and correlations between ruminal bacteria abundance and performance measures. *J. Dairy Sci.* **100**, 8102-8118.
- Kim J., Hyun Y., Sohn K., Kim T., Woo H. and Han I. (2000). Effects of mannan-oligosaccharide and protein levels on growth performance and immune status in pigs weaned at 21 days of age. *Korean J. Anim. Sci.* **42**, 489-498.
- Kurtz D.M. and Travlos G.S. (2017). The Clinical Chemistry of Laboratory Animals. CRC Press, Taylor & Francis Group, LLC, Boca Raton, FL, USA.
- Lei C.L., Dong G.Z., Jin L., Zhang S. and Zhou J. (2013). Effects of dietary supplementation of montmorillonite and yeast cell wall on lipopolysaccharide adsorption, nutrient digestibility and growth performance in beef cattle. *Livest. Sci.* **158**, 57-63.
- Liu Y.Z., Lang M., Zhen Y.G., Chen X., Sun Z., Zhao W., Zhang X.F., Wang T. and Qin G.X. (2019). Effects of yeast culture supplementation and the ratio of non-structural carbohydrate to fat on growth performance, carcass traits and the fatty acid profile of the longissimus dorsi muscle in lambs. *J. Anim. Physiol. Anim. Nutr.* **103**, 1274-1282.
- Mahmoud M.M., Youssef I.M.I., Abd El-Tawab M.M., Bakr H.A., Eissa N.A. and Hassan M.S. (2020). Influence of probiotic and yeast culture supplementation on selected biochemical and immunological parameters of growing lambs. *Polish J. Vet. Sci.* **23(1)**, 5-12.
- Malekkhahi M., Tahmasbi A.M., Naserian A.A., Danesh Mesgaran M., Kleen J.L. and Parand A.A. (2015). Effects of essential oils, yeast culture and malate on rumen fermentation, blood metabolites, growth performance and nutrient digestibility of baluchi lambs fed high-concentrate diets. *J. Anim.*

- Physiol. Anim. Nutr.* **99**, 221-229.
- Mitchell L.K. and Heinrichs A.J. (2020). Feeding various forages and live yeast culture on weaned dairy calf intake, growth, nutrient digestibility, and ruminal fermentation. *J. Dairy Sci.* **103**, 8880-8897.
- Mohamed A., Mohamed M., Abd-El-Ghany F., Mahgoup A. and Ibrahim K. (2015). Influence of some trace minerals in form of nanoparticles as feed additives on lamb's performance. *J. Anim. Poult. Prod.* **6**, 693-703.
- Mohamed A.H., Mohamed M.Y., Ibrahim K., Abd-El-Ghany F.T.F. and Mahgoup A.A.S. (2017). Impact of nano-zinc oxide supplementation on productive performance and some biochemical parameters of ewes and offspring. *Egyptian J. Sheep Goat Sci.* **12**, 49-64.
- Mullins C., Mamedova L., Carpenter A., Ying Y., Allen M., Yoon I. and Bradford B. (2013). Analysis of rumen microbial populations in lactating dairy cattle fed diets varying in carbohydrate profiles and *Saccharomyces cerevisiae* fermentation product. *J. Dairy Sci.* **96**, 5872-5881.
- Murthy N. (2013). Investigation of Biochemical and Histopathological Modifications Induced by Selenium Toxicity on Albino Mice. LAP Lambert Academic Publishing, Saarbrücken, Germany.
- Nawab A., Liu W., Li G., Ibtisham F., Fox D., Zhao Y., Wo J., Xiao M., Nawab Y. and An L. (2019). The potential role of probiotics (nutraceuticals) in gut health of domestic animals; an alternative to antibiotic growth promoters. *J. Hellenic Vet. Med. Soc.* **69**, 1169-1188.
- Novoselec J., Šperanda M., Klir Ž., Mioc B., Steiner Z. and Antunović Z. (2017). Blood biochemical indicators and concentration of thyroid hormones in heavily pregnant and lactating ewes depending on selenium supplementation. *Acta Vet. Brno.* **86**, 353-363.
- NRC. (2007). Nutrient Requirements: Sheep, Goats, Cervids, and New World Camelids. National Academy Press, Washington, DC, USA.
- Pieper R., Vahjen W., Neumann K., Van Kessel A.G. and Zentek J. (2012). Dose-dependent effects of dietary zinc oxide on bacterial communities and metabolic profiles in the ileum of weaned pigs. *J. Anim. Physiol. Nutr.* **96**, 825-833.
- Plaza-Díaz J., Ruiz-Ojeda F.J., Gil-Campos M. and Gil A. (2019). Mechanism of action of probiotics. *Adv. Nutr.* **10**, 49-66.
- Rajendran D., Kumar G., Ramakrishnan S. and Shibi T. (2013). Enhancing the milk production and immunity in Holstein Friesian crossbred cow by supplementing novel nano zinc oxide. *Res. J. Biotechnol.* **8**, 11-17.
- Riyazi H., Rezaei J. and Rouzbehan Y. (2019). Effects of supplementary nano-ZnO on *in vitro* ruminal fermentation, methane release, antioxidants, and microbial biomass. *Turk. J. Vet. Anim. Sci.* **43**, 737-746.
- Reddy P.R.K., Yasaswini D., Reddy P.P.R., Zeineldin M., Adebeye M.J. and Hyder I. (2020). Applications, challenges, and strategies in the use of nanoparticles as feed additives in equine nutrition. *Vet. World.* **13**, 1685-1696.
- Rozeboom D., Shaw D., Tempelman R., Miguel J., Pettigrew J. and Connolly A. (2005). Effects of mannan oligosaccharide and an antimicrobial product in nursery diets on performance of pigs reared on three different farms. *J. Anim. Sci.* **83**, 2637-2644.
- SAS Institute. (2013). SAS®/STAT Software, Release 9.4. SAS Institute, Inc., Cary, NC, USA.
- Seven P.T., Seven I., Baykalir B.G., Mutlu S.I. and Salem A.Z.M. (2018). Nanotechnology and nano-propolis in animal production and health: An overview. *Italian J. Anim. Sci.* **17**, 921-930.
- Song B., Wu T., You P., Wang H.L., Burke J., Kang K., Yu W., Wang M., Li B., He Y., Huo Q., Li C., Tian W., Li R., Li J., Wang C. and Sun X. (2021). Dietary supplementation of yeast culture into pelleted total mix rations improves the growth performance of fattening lambs. *Front. Vet. Sci.* **8**, 657816-657826.
- Sun H., Chen J., Xiong D. and Long M. (2023). Detoxification of selenium yeast on mycotoxins and heavy metals: A review. *Biol. Trace Elem. Res.* **201(11)**, 5441-5454.
- Swain P.S., Rao S.B.N.D., Poornachandra K.T., Loksha R.E. and Kumar R.D. (2019). Effect of nanozinc supplementation on hematological and blood biochemical profiles in goats. *Int. J. Curr. Microbiol. Appl. Sci.* **8(9)**, 2688-2696.
- Tripathi M.K. and Karim S.A. (2011). Effect of yeast cultures supplementation on live weight change, rumen fermentation, ciliate protozoa population, microbial hydrolytic enzymes status and slaughtering performance of growing lamb. *Livest. Sci.* **135**, 17-25.
- Valenzuela-Grijalva N.V., Pinelli-Saavedra A., Muhlia-Almazan A., Domínguez-Díaz D. and González-Ríos H. (2017). Dietary inclusion effects of phytochemicals as growth promoters in animal production. *J. Anim. Sci. Technol.* **59**, 8-17.
- Van Soest P.J., Robertson J.B. and Lewis B.A. (1991). Methods for dietary fiber, neutral detergent fiber, and non-starch polysaccharides in relation to animal nutrition. *J. Dairy Sci.* **74(10)**, 3583-3597.
- Wang C., Zhang L., Su W., Ying Z., He J., Zhang L., Zhong X. and Wang T. (2017). Zinc oxide nanoparticles as a substitute for zinc oxide or colistin sulfate: effects on growth, serum enzymes, zinc deposition, intestinal morphology and epithelial barrier in weaned piglets. *PLoS One.* **12**, e0181136.
- Wang J., Zhao G., Zhuang Y., Chai J. and Zhang N. (2022). Yeast (*Saccharomyces cerevisiae*) culture promotes the performance of fattening sheep by enhancing nutrients digestibility and rumen development. *Fermentation.* **8**, 719-728.
- White L., Newman M., Cromwell G. and Lindemann M. (2002). Brewers dried yeast as a source of mannan oligosaccharides for weanling pigs. *J. Anim. Sci.* **80**, 2619-2628.
- Zaboli K., Aliarabi H., Bahari A. and Abbasalipourkabar R. (2013). Role of dietary nano-zinc oxide on growth performance and blood levels of mineral: A study on Iranian Angora (Markhoz) goat kids. *J. Pharm. Health Sci.* **2**, 19-26.
- Zhai X., Zhang C., Zhao G., Stoll S., Ren F. and Leng X. (2017). Antioxidant capacities of the selenium nanoparticles stabilized by chitosan. *J. Nanobiotechnol.* **15**, 4-11.
- Zhan X., Wang M., Zhao R., Li W. and Xu Z. (2007). Effects of different selenium source on selenium distribution, loin quality and antioxidant status in finishing pigs. *Anim. Feed Sci. Technol.* **132**, 202-211.

Zhang R., Zhou M., Tu Y., Zhang N.F., Deng K.D., Ma T. and Diao Q.Y. (2016). Effect of oral administration of probiotics on growth performance, apparent nutrient digestibility and

stress-related indicators in holstein calves. *J. Anim. Physiol. Anim. Nutr.* **100**, 33-38.
