

Detection of *bla*_{CTX-M}, *bla*_{TEM} and *bla*_{SHV} genes in *Enterobacter cloacae* isolates

Elahe Barzam Dehkordi, Elahe Tajbakhsh*, Hasan Momtaz

Department of Microbiology, Faculty of Basic Sciences, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran

*Corresponding Author: ee_tajbakhsh@yahoo.com

Received: 03/11/2023, Accepted: 24/12/2023

Extended Abstract

Introduction: *Enterobacter cloacae* is a Gram-negative, facultatively anaerobic bacterium commonly found in the human gut but is also associated with a wide range of nosocomial infections, including urinary tract infections (UTIs), pneumonia, and sepsis. The increasing antibiotic resistance among *Enterobacter* species, particularly due to the production of extended-spectrum beta-lactamases (ESBLs), poses a significant public health challenge. ESBL-producing strains hydrolyze a broad range of beta-lactam antibiotics, making treatment difficult. Among the most common ESBL genes are *bla*_{CTX-M}, *bla*_{TEM}, and *bla*_{SHV}, which contribute to multidrug resistance (MDR) in *Enterobacter* species. This study aimed to determine the prevalence of these genes in *Enterobacter cloacae* isolates from patients with UTIs in Shahrekord, Iran, and assess their antibiotic resistance profiles.

Methods: This cross-sectional study was conducted over one year, from June 2019 to June 2020, on 1000 urine samples collected from patients suspected of UTIs in Shahrekord, Iran. The bacterial isolates were identified using standard biochemical and microbiological methods. Antibiotic susceptibility testing was performed using the disk diffusion method following Clinical and Laboratory Standards Institute (CLSI) guidelines. The presence of *bla*_{CTX-M}, *bla*_{TEM}, and *bla*_{SHV} genes was detected using polymerase chain reaction (PCR) with specific primers.

Results and Discussion: Of the 1000 urine samples, 300 were culture-positive for UTI pathogens, among which 65 isolates (21.66%) were identified as *Enterobacter cloacae*. The prevalence of infections was significantly higher in female patients (76.92%) compared to males (23.08%). The highest infection rates were observed in patients aged above 60 years and those aged between 15-30 years. Antibiotic susceptibility testing revealed high resistance rates among the isolates, with the highest resistance observed against trimethoprim-sulfamethoxazole (84.62%) and the lowest resistance against nitrofurantoin (23.08%). Multidrug resistance (MDR) was detected in over 50% of the isolates. The phenotypic confirmatory test for ESBL production showed that 25 (38.26%) isolates were ESBL producers. PCR analysis detected *bla*_{CTX-M} in 28% of isolates, *bla*_{TEM} in

32%, and *blaSHV* in 10%. Some isolates harbored multiple resistance genes, with co-occurrence of *blaTEM* and *blaSHV* in 4%, *blaSHV* and *blaCTX-M* in 8%, and *blaCTX-M* and *blaTEM* in 16% of isolates. The presence of ESBL-producing *Enterobacter cloacae* in UTIs is concerning due to its association with treatment failures and increased morbidity. The high prevalence of resistance genes suggests that empirical antibiotic treatment strategies should be revised to account for MDR strains. The results also highlight the necessity for routine surveillance and stringent antimicrobial stewardship programs to mitigate the spread of resistant strains.

Conclusion: This study provides evidence of a high prevalence of ESBL-producing *Enterobacter cloacae* strains in Shahrekord, Iran, with a significant presence of *blaCTX-M*, *blaTEM*, and *blaSHV* genes. The increasing resistance to commonly used antibiotics underscores the urgent need for infection control measures and judicious use of antimicrobials. Implementing molecular surveillance and incorporating novel therapeutic strategies will be crucial in controlling the dissemination of resistant strains.

Keywords: *Enterobacter cloacae*, Extended-Spectrum Beta-Lactamases, Antibiotic Resistance, Urinary Tract Infections