

## Catalyst free synthesis of a of triazothiazole using multicomponent reaction of isothiocyanates

Narges Ghasemi\*

National Petrochemical Company (NPC), petrochemical Research and Technology Company, Arak Center, Iran

Received: November 2021; Revised: December 2021; Accepted: January 2022

**Abstract:** Triazothiazole derivatives are obtained in good to excellent yields by proceeding through a simple, mild, and efficient procedure utilizing isothiocyanate, activated acetylenic compounds, ammonium acetate and hydrazoyl chloride in water at room temperature.

**Keywords:** Triazothiazole, Activated acetylenes, Ammonium acetate, Water, Arylisothiocyanate.

### Introduction

Multicomponent reactions (MCRs), with three or more reactants join in a one-pot procedure to afford a single product [1-3]. They are economically and environmentally useful because multi-step synthesis produce large amounts of trash frequently because of complex isolation actions frequently involving comfortable, toxic, and hazardous solvents after each step [4-7]. MCRs are absolutely suited for combinatorial library synthesis and increased utilize in the finding procedure for new drugs and agrochemicals [8]. They supply a dominant tool toward the one-pot synthesis of diverse and complex compounds as well as small and drug-like heterocycles [9]. Green chemistry move towards hold out significant potential not only for reduction of byproducts, waste produced, and lowering of energy but also in the expansion of new methodologies toward before exclusive materials, using existing technologies [10]. Between existing part of chemistry, medicinal and pharmaceutical chemistry are possibly developed for greening [11]. Hence, we investigated a simple three-component reaction between activated acetylenic compounds,

primary amines and isothiocyanates in the presence of *N*-formylmorpholine as solvent-free conditions at room temperature which afforded 1,3-thiazolane derivatives **4** in good isolated yields. With propargylic esters yield of reactions is low (Scheme 1).


### Results and discussion

The  $^1\text{H}$  NMR spectrum of **5a** displayed signals for vicinal methine protons at  $\delta = 4.78$  and  $4.92$ , which appeared as two set of doublets with  $^3J_{\text{HH}}$  values of  $12.4$  Hz. The methoxy groups showed two separate singlet at  $\delta = 3.78$  and  $3.85$ . Observation of  $^3J_{\text{HH}} = 12.4$  Hz for the vicinal methine protons in **5a** indicates the dominance of anti arrangement. The carbonyl groups resonances in the  $^{13}\text{C}$  NMR spectra of **5a** are appeared at  $172.5$  (C=O),  $173.7$  (C=O) ppm. Also the mass spectra of **5a** displayed the molecular ion peak in the appropriate  $m/z$  values. A proposed mechanism for the formation of compound **5** is shown in Scheme 2. Apparently, the zwitterionic intermediate **6** which formed from the reaction of triethylamine ( $\text{X}_3\text{N}$ ) and electron deficient acetylenic ester **3** is protonated by the intermediate **5** that was generated in situ from the reaction of primary amine **2** and isothiocyanate **1**, produce intermediates **7** and **8**. Nucleophilic attack of the conjugate base **7** on intermediate **8** leads to adduct **9** which undergo proton shifts to afford new


\*Corresponding author: Tel: 0098-8633677201-9; Fax: 0098-8633677203, E-mail: [naghasemi.16@gmail.com](mailto:naghasemi.16@gmail.com)

zwitterionic **10**. Finally, intramolecular cyclization of **10** with elimination of triethylamine produces

compound **12** that with react by hydrazoyl chloride **4** produced desired compounds **5**.



**Scheme 1:** Reaction of activated acetylenes, isothiocyanates and primary aminins



**Scheme 2:** Proposed mechanism for the formation of **5**.

## Conclusion

In conclusion, we found that the reaction of activated acetylenic compounds with isothiocyanates and ammonium acetate in the presence of catalytic amount of triethylamine to a facile synthesis of some functionalized triazothiazoles under solvent-free conditions without using any catalyst.

## Experimental

All chemicals used in this work were prepared from Fluka (Buchs, Switzerland) and were used without further purification. Electrothermal 9100 apparatus is employed for measuring of melting points of products. Elemental analyses for C, H, and N were performed with Heraeus CHN-O-Rapid analyzer. Mass spectra were recorded on a FINNIGAN-MAT 8430 spectrometer operating at an ionization potential of 70 eV. Measurement of IR spectra was performed by Shimadzu IR-460 spectrometer.  $^1\text{H}$ , and  $^{13}\text{C}$  NMR spectra were evaluated with a BRUKER DRX-500 AVANCE spectrometer at 500.1 and 125.8 MHz, respectively.  $^1\text{H}$ , and  $^{13}\text{C}$ , spectra were obtained for solutions in  $\text{CDCl}_3$  using TMS as internal standard or 85%  $\text{H}_3\text{PO}_4$  as external standard.

### General procedure for preparation of compounds 5

To a magnetically stirred mixture of activated acetylenes **3** (2 mmol) and triethylamine (5 mol%) was added mixture of isothiocyanates **1** and ammonium acetate **3** (2 mmol) and hydrazoyl chloride **4** at room temperature. The reaction mixture was then stirred. After completion of the reaction [TLC (AcOEt/hexane 1:7) monitoring], 15 mL  $\text{H}_2\text{O}$  was poured into the reaction mixture. The solid residue was filtered and washed by cold diethyl ether to afforded pure compounds **4** and **12**.

**Dimethyl 2-(methylimino)-3-phenyl-1,3-thiazolane-4,5-dicarboxylate (5a):** Yellow powder, m.p. 156-158°C, yield: 0.46 g (75%). IR (KBr) ( $\nu_{\text{max}}/\text{cm}^{-1}$ ): 1745, 1738, 1698, 1657, 1574, 1467, 1382, 1215  $\text{cm}^{-1}$ . Anal. Calcd for  $\text{C}_{14}\text{H}_{16}\text{N}_2\text{O}_4\text{S}$  (308.35): C, 54.53; H, 5.23; N, 9.08. Found: C, 54.62; H, 5.34; N, 9.23%.  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  2.83 (3 H, s, NMe), 3.78 (3 H, s, MeO), 3.85 (3 H, s, MeO), 4.78 (1 H, d,  $^3J = 12.4$ , CH), 4.92 (1 H, d,  $^3J = 12.4$ , CH), 7.23 (1 H, t,  $^3J = 7.4$ , CH), 7.35 (2 H, d,  $^3J = 7.6$ , 2 CH), 7.54 (2 H, t,  $^3J = 7.6$ , 2 CH) ppm.  $^{13}\text{C}$  NMR (125.7 MHz,  $\text{CDCl}_3$ ):  $\delta$  34.6 (NMe), 42.7 (CH), 51.6 (MeO), 52.4 (MeO), 58.4

(CH), 122.8 (CH), 128.3 (2 CH), 129.6 (2 CH), 139.8 (C), 163.4 (C=N), 172.5 (C=O), 173.7 (C=O) ppm. MS,  $m/z$  (%): 308 ( $\text{M}^+$ , 15), 277 (86), 77 (64), 31 (100).

**Dimethyl 2-(ethylimino)-3-(4-methoxyphenyl)-1,3-thiazolane-4,5-dicarboxylate (5b):** Pale yellow powder, m.p. 168-170 °C, yield: 0.59 g (87%). IR (KBr) ( $\nu_{\text{max}}/\text{cm}^{-1}$ ): 1742, 1736, 1686, 1632, 1525, 1487, 1325, 1219  $\text{cm}^{-1}$ . Anal. Calcd for  $\text{C}_{16}\text{H}_{20}\text{N}_2\text{O}_5\text{S}$  (352.41): C, 54.53; H, 5.72; N, 7.95. Found: C, 54.64; H, 5.80; N, 8.10%.  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.24 (3H, t,  $^3J = 7.3$ ,  $\text{CH}_3$ ), 3.27 (2 H, q,  $^3J = 7.3$ ,  $\text{CH}_2$ ), 3.70 (3 H, s, MeO), 3.76 (3 H, s, MeO), 3.87 (3 H, s, MeO), 4.75 (1 H, d,  $^3J = 12.2$ , CH), 4.87 (1 H, d,  $^3J = 12.2$ , CH), 7.14 (2 H, d,  $^3J = 7.8$ , 2 CH), 7.28 (2 H, d,  $^3J = 7.6$ , 2 CH) ppm.  $^{13}\text{C}$  NMR (125.7 MHz,  $\text{CDCl}_3$ ):  $\delta$  14.2 ( $\text{CH}_3$ ), 41.5 ( $\text{CH}_2$ ), 43.7 (CH), 51.5 (MeO), 52.6 (MeO), 55.4 (MeO), 59.3 (CH), 111.2 (2 CH), 130.3 (2 CH), 134.8 (C), 154.2 (C), 160.7 (C=N), 171.8 (C=O), 172.6 (C=O) ppm. MS,  $m/z$  (%): 352 ( $\text{M}^+$ , 10), 321 (64), 108 (96), 31 (100).

**Diethyl 2-(buthylimino)-3-(4-methoxyphenyl)-1,3-thiazolane-4,5-dicarboxylate (5c):** White powder, m.p. 162-164 °C, yield: 0.70 g (83%). IR (KBr) ( $\nu_{\text{max}}/\text{cm}^{-1}$ ): 1740, 1738, 1687, 1645, 1438, 1357, 1256  $\text{cm}^{-1}$ . Anal. Calcd for  $\text{C}_{20}\text{H}_{28}\text{N}_2\text{O}_5\text{S}$  (408.51): C, 58.80; H, 6.91; N, 6.86. Found: C, 58.92; H, 6.98; N, 6.90%.  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.19 (3H, t,  $^3J = 7.2$ ,  $\text{CH}_3$ ), 1.22 (3H, t,  $^3J = 7.4$ ,  $\text{CH}_3$ ), 1.28 (3H, t,  $^3J = 7.3$ ,  $\text{CH}_3$ ), 1.68 (2 H, q,  $^3J = 7.3$ ,  $\text{CH}_2$ ), 1.78 (2 H, m,  $\text{CH}_2$ ), 2.83 (2 H, t,  $^3J = 6.8$ ,  $\text{NCH}_2$ ), 3.75 (3 H, s, MeO), 4.12 (2 H, q,  $^3J = 7.3$ ,  $\text{CH}_2\text{O}$ ), 4.23 (2 H, q,  $^3J = 7.3$ ,  $\text{CH}_2\text{O}$ ), 4.62 (1 H, d,  $^3J = 11.7$ , CH), 5.02 (1 H, d,  $^3J = 11.7$ , CH), 7.12 (2 H, d,  $^3J = 7.6$ , 2 CH), 7.32 (2 H, d,  $^3J = 7.6$ , 2 CH) ppm.  $^{13}\text{C}$  NMR (125.7 MHz,  $\text{CDCl}_3$ ):  $\delta$  13.3 ( $\text{CH}_3$ ), 13.8 ( $\text{CH}_3$ ), 14.3 ( $\text{CH}_3$ ), 21.4 ( $\text{CH}_2$ ), 32.5 ( $\text{CH}_2$ ), 43.6 (CH), 54.8 (MeO), 59.5 (CH), 61.2 ( $\text{CH}_2\text{O}$ ), 62.0 ( $\text{CH}_2\text{O}$ ), 62.7 ( $\text{NCH}_2$ ), 114.5 (2 CH), 130.8 (2 CH), 135.4 (C), 156.7 (C), 161.2 (C=N), 172.3 (C=O), 174.2 (C=O) ppm. MS,  $m/z$  (%): 408 ( $\text{M}^+$ , 8), 363 (84), 108 (68), 45 (100).

**Diethyl 2-(tert-butylimino)-3-(4-methylphenyl)-1,3-thiazolane-4,5-dicarboxylate (5d):** yellow powder, m.p. 164-166 °C, yield: 0.59 g (75%). IR (KBr) ( $\nu_{\text{max}}/\text{cm}^{-1}$ ): 1736, 1732, 1694, 1587, 1467, 1346, 1238  $\text{cm}^{-1}$ . Anal. Calcd for  $\text{C}_{20}\text{H}_{28}\text{N}_2\text{O}_4\text{S}$  (392.51): C, 61.20; H, 7.19; N, 7.14. Found: C, 61.32; H, 7.25; N, 7.22%.  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.25 (3H, t,  $^3J = 7.4$ ,  $\text{CH}_3$ ), 1.32 (3H, t,  $^3J = 7.4$ ,  $\text{CH}_3$ ), 1.35 (9H, s,  $\text{Me}_3\text{C}$ ), 2.28 (3 H, s,  $\text{CH}_3$ ), 4.15 (2 H, q,  $^3J = 7.4$ ,  $\text{CH}_2\text{O}$ ), 4.28

(2 H, q,  $^3J = 7.4$ , CH<sub>2</sub>O), 4.73 (1 H, d,  $^3J = 11.5$ , CH), 4.96 (1 H, d,  $^3J = 11.5$ , CH), 7.24 (2 H, d,  $^3J = 7.5$ , 2 CH), 7.36 (2 H, d,  $^3J = 7.6$ , 2 CH) ppm. <sup>13</sup>C NMR (125.7 MHz, CDCl<sub>3</sub>):  $\delta$  13.8 (CH<sub>3</sub>), 14.2 (CH<sub>3</sub>), 22.4 (CH<sub>3</sub>), 28.7 (Me<sub>3</sub>C), 44.3 (CH), 48.7 (Me<sub>3</sub>C), 58.7 (CH), 61.4 (CH<sub>2</sub>O), 62.3 (CH<sub>2</sub>O), 129.4 (2 CH), 130.2 (C), 131.4 (2 CH), 140.7 (C), 160.4 (C=N), 172.5 (C=O), 175.3 (C=O) ppm. MS, *m/z* (%): 392 (M<sup>+</sup>, 20), 377 (84), 91 (84), 45 (100).

**Dimethyl 2-(methyylimino)-3-(4-bromophenyl)-1,3-thiazolane-4,5-dicarboxylate (5e):** yellow crystals, m.p. 183-185 °C, yield: 0.62 g (80%). IR (KBr) ( $\nu_{\max}/\text{cm}^{-1}$ ): 1737, 1732, 1695, 1587, 1485, 1436, 1342, 1225 cm<sup>-1</sup>. Anal. Calcd for C<sub>14</sub>H<sub>15</sub>BrN<sub>2</sub>O<sub>4</sub>S (387.25): C, 43.42; H, 3.90; N, 7.23. Found: C, 43.53; H, 3.95; N, 7.32%. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  3.12 (3H, s, NMe), 3.75 (3 H, s, MeO), 3.82 (3 H, s, MeO), 4.83 (1 H, d,  $^3J = 11.8$ , CH), 4.92 (1 H, d,  $^3J = 11.8$ , CH), 7.10 (2 H, d,  $^3J = 7.8$ , 2 CH), 7.54 (2 H, d,  $^3J = 7.8$ , 2 CH) ppm. <sup>13</sup>C NMR (125.7 MHz, CDCl<sub>3</sub>):  $\delta$  34.5 (NCH<sub>3</sub>), 44.2 (CH), 51.2 (MeO), 51.8 (MeO), 60.3 (CH), 116.7 (C), 129.7 (2 CH), 132.6 (2 CH), 139.4 (C), 162.3 (C=N), 172.4 (C=O), 173.8 (C=O) ppm. MS, *m/z* (%): 387 (M<sup>+</sup>, 15), 356 (78), 156 (64), 31 (100).

## References

- [1] Domling, A. *Comb. Chem. High Throughput Screening* **1998**, *1*, 1.
- [2] Domling, A.; Ugi, I. *Angew. Chem., Int. Ed.* **2000**, *39*, 3169.
- [3] Weber, L. *Drug Discovery Today* **2002**, *7*, 143.
- [4] Multicomponent Reactions; Zhu, J., Bienayme, H., Eds.; Wiley-VCH: Weinheim, Germany, **2005**.
- [5] Wipf, P.; Kendall, C. *Chem. Eur. J.* **2002**, *8*, 1779–1784.
- [6] Balme, G.; Bossharth, E.; Monteiro, N. *Eur. J. Org. Chem.* **2003**, 4101–4111.
- [7] Jacobi von Wangelin, A.; Neumann, H.; Gordes, D.; Klaus, S.; Strubing, D.; Beller, M. *Chem. Eur. J.* **2003**, *9*, 4286–4294.
- [8] (a) Domling, A.; Ugi, I. *Angew. Chem., Int. Ed.* **2000**, *39*, 3168; (b) Ugi, I.; Domling, A. *Endeavour* **1994**, *18*, 115; (c) Heck, S.; Domling, A. *Synlett* **2000**, 424.
- [9] Weber, L. *Curr. Med. Chem.* **2002**, *9*, 2085.
- [10] Cave, G. W. V.; Raston, C. L.; Scott, J. L. *Chem. Commun.* **2001**, 2159-2169.
- [11] Sheldon, R. A. *Chem. Ind.* **1997**, 12