

Effects of Feeding Different Levels of Whole and Ground Flaxseed in Late Pregnancy on the Performance of Karakul Ewes

S.H. Hosseini¹, M.R. Dehghani^{1*}, A. Karimi², M. Yousef Elahi¹ and M.J. Abarghuei²

¹ Department of Animal Science, Faculty of Agriculture, University of Zabol, Zabol, Iran

² Department of Animal Science, Fars Agricultural and Natural Resources research and Education Center, Agricultural Research, Education and Extension Organization (AREEO), Shiraz, Iran

*Correspondence Tel: 98-915-5422024, E-mail: Mr.dehghani@uoz.ac.ir

ABSTRACT

This study were conducted to determine the effect of different levels of flaxseed, whole and ground (8 and 15%), in late pregnancy on the performance of ewes. 30 Karakul ewes (ages between 2 and 2.5 years old, mean weight of 55 ± 2.5 kg) were used in a completely randomised design with five treatments and six replicates. Treatments include: 1) control diet, a diet without flaxseed; 2) WFS-L, a diet containing 8% whole flaxseed; 3) WFS-H, a diet containing 15% whole flaxseed; 4) GFS-L, a diet containing 8% ground flaxseed; and 5) GFS-H, a diet containing 15% ground flaxseed. The addition of flaxseed (whole and ground) had no significant effect on dry matter intakes or body weight changes in ewes and lambs ($P>0.05$), but in comparison, crude protein and ether extract digestibility was significantly ($P<0.05$) higher in ewes fed whole and ground flaxseed compared to those fed the control diet. The yield and composition of colostrum were not affected by experimental treatments ($P>0.05$), except for colostrum protein, which increased by using treatments with whole flaxseed ($P<0.05$). Glucose and cholesterol in blood plasma were significantly increased in ewes fed whole and ground flaxseed compared to the control treatment. ($P<0.05$), and the highest blood glucose and cholesterol levels were observed in the treatment of 15% whole flaxseed. The research showed that whole flaxseed (15%) can be added to sheep's diets, and it can be concluded that processing flaxseed for ewes in late pregnancy might not be necessary to improve performance.

Key words apparent digestibility, flaxseed, performance. pregnancy, Karakul ewe.

INTRODUCTIN

Given that 80% of fetus growth takes place in the last two months of pregnancy, the nutrient requirements of ewes increase significantly during this period (Dawson *et al.* 1999). The nutrition management of ewes during late pregnancy plays a crucial role in determining various factors, such as lambs' birth weight and good-quality colostrum, and early detection of colostrum quality in post-lambing is also extremely important for lamb health and performance (Karakus and Atmac, 2016). Petit and Benchaar (2007) observed that feeding cows with flaxseed as a source of omega-3 is an effective way to improve feed intake and energy balance. Flaxseed (*Linum usitatissimum*) contains approximately 20% crude protein (CP) and 40% oil on a dry matter (DM) basis (Petit, 2010), and it is also a rich source of essential fatty acids, of which 50–70% is alpha-linolenic acid (ALA), a rich source of n-3 polyunsaturated fatty acids (PUFA) (Xu *et al.* 2013). Greco *et al.* (2015) stated that reducing the ratio of n-6 to n-3 FA in the diet of dairy cows increased dry matter intake (DMI). Rosa Velazquez *et al.* (2020) also observed that supplementation of the diet with omega-3 (10.1 g/kg dry matter) resources such as flaxseed has beneficial effects on the development and growth of the lambs. Moreover, some studies showed that when flaxseed was used in the ruminant diet, disease resistance increased due to the effect of ALA on the immune system. In

other words, the level of immunity against diseases in livestock increased (Nudda *et al.* 2015). Farmer *et al.* (2010) reported that feeding flaxseed meals in late gestation supplementation improved the post weaning growth of piglets, possibly due to increased carcass glycogen on day 1. Thus, we hypothesized that supplementing pregnant ewes' diets with flaxseeds in late gestation may enhance the performance of them. There is also little research on direct comparisons of the use of whole and ground flaxseed in late pregnancy on the performance of Karakul ewes. Therefore, the main objective of the present study was to investigate the effects of feeding diets with flaxseed (whole or ground) during late pregnancy on performance, nutrient digestibility, colostrum yield, colostrum composition, and some plasma metabolites of Karakul ewes.

MATERIALS AND METHODS

Animals, experimental treatments, and nutrition management

This experiment was carried out in the Fars Agricultural and Natural Resources Research and Training Center in the fall of 2018 for 8 weeks before calving (including 2 weeks of habituation and 6 weeks of testing period) and 48 hours after calving. In this experiment, 30 Karakul ewes (ages between 2 and 3 years old and mean of weight of 55 ± 2.5 kg) were used in a completely randomized design with five treatments and six replicates. Treatments include 1) control diet, a diet without flaxseed; 2) WFS-L, a diet containing 8% whole flaxseed; 3) WFS-H, a diet containing 15% whole flaxseed; 4) GFS-L, a diet containing 8% ground flaxseed; and 5) GFS-H, a diet containing 15% ground flaxseed. Experimental diets were balanced based on NRC, 2001 to meet the recommended nutrient requirements and adjusted based on DM using small ruminant feeding system software (SRNS, 2010). The body weight of ewes was recorded and they were randomly assigned to one of five experimental treatments inside an individual cage. The experiment was conducted during the last two months of pregnancy. During the experiment, the ewes were nourished twice a day (8.00 and 16.00 h) and had free access to fresh drinking water. Ingredients and chemical composition of diets and flaxseed during late pregnancy are shown in Tables 1 and 2, respectively.

Table 1 Ingredients and composition of the experimental diets

Ingredients (%)	Treatments				
	Control	WFS-L	WFS-H	GFS-L	GFS-H
Alfalfa hay	33.34	33.34	33.34	33.34	33.34
Wheat straw	27.01	27.01	27.01	27.01	27.01
Flaxseed	-	8	15	8	15
Wheat bran	16	16	16	16	16
Barley grains	9.33	8.53	8.13	8.53	8.13
Soybean meal	7.33	4.93	2.53	4.93	2.53
Corn grain	5.33	5.33	5.33	5.33	5.33
CaCO ₃	0.67	0.67	0.67	0.67	0.67
Salt	0.33	0.33	0.33	0.33	0.33
Sodium bicarbonate	0.33	0.33	0.33	0.33	0.33
Mineral and vitamin supplement ^b	0.33	0.33	0.33	0.33	0.33
Chemical Composition					
Metabolizable energy (Mcal / kg)	2.14	2.17	2.19	2.17	2.19
Dry matter (%)	91.28	91.32	91.74	91.32	91.74
Crude protein (%)	16.86	16.47	16.04	16.47	16.04
Ether extract (%)	3.09	4.02	4.83	4.02	4.83
Neutral detergent fiber (%)	39.11	39.46	39.77	39.46	39.77
Acid detergent fiber (%)	24.32	24.63	24.89	24.63	24.89
Ca (%)	0.88	0.80	0.82	0.80	0.82
P (%)	0.31	0.30	0.30	0.30	0.30

^a Each kilogram of vitamin–mineral premix contained: vitamin A (50,000 IU), vitamin D3 (10,000 IU), vitamin E (1000 IU), Ca (196 g), P (96 g), Na (71 g), Mg (19 g), Fe (3 g), Cu (0.3 g), Mn (2 g), Zn (3 g), Co (0.1 g), I (0.1 g) and Se (0.001 g); ME, Metabolism Energy (Mcal/kg DM).

Table 2 Compounds of flaxseed

Ingredients	(% DM)
DM (% of fresh weight)	93
Crude protein (% of DM)	19.3
Ether extract (% of DM)	35
Crude fiber (% of DM)	26
Ash (% of DM)	2.92

Performance ewes

The amount of feed intake and daily residue were recorded to calculate the DMI. The ewes of each treatment were weighed at intervals two months before lambing and one hour after lambing. The newborn lambs were allowed to be licked and dried by the ewes and within one hour were weighed, their sex was determined, and ear-tagged (Hashemi *et al.* 2008).

Apparent digestibility by the acid-insoluble ash method

A digestibility trial was performed during late pregnancy (7 days before the expected birth date). On three consecutive days, the faeces were collected using the faecal grab samples method in all ewes three times a day (7.00, 14.00, and 18.00) (Meteb *et al.* 2020). Acid-insoluble ash was used as an internal marker to estimate faecal output and nutrient digestibility. The digestibility coefficient of a given nutrient was calculated according to the following formula (Van Keulen and Young, 1977).

$$\text{Digestibility} = 100 - \frac{\% \text{ indicator in the feed}}{\% \text{ indicator in the feces}} \times \frac{\% \text{ nutrient in the feces}}{\% \text{ nutrient in the feed}}$$

Colostrum production

The measure of colostrum was determined by measuring the difference between the pre- and post-sucking weights of lambs. The rest of the colostrum was milked and weighed. The colostrum yield can be calculated by adding the colostrum consumed by the lambs to the milked colostrum. The colostrum yield of each ewe was recorded at 12, 24, and 48 h after birth (Ocak *et al.* 2005).

Blood metabolites

Blood samples were taken from the jugular vein using heparinized plastic syringes 3 hours after the morning feeding (Ababakri *et al.* 2021), 30 days after the beginning of the experiment (Hashemi *et al.* 2008). Blood samples were withdrawn from the jugular vein into vacutainer tubes containing EDTA. The plasma was isolated by centrifuging at 3000 rpm for 20 minutes and frozen at -20°C until the following analysis. Blood serum samples were analyzed using Pars Azmon kits and spectrophotometer (Jenway 6300, UK).

Determination of chemical composition of feed and feces

Sample of feed and feces were oven-dried at 60°C for 48 h and milled to pass through a 1-mm mesh (Wiley mill, Swedesboro, NJ). Samples of diet were analyzed for dry matter (DM), Crude protein (CP), Ether extract (EE), organic matter (OM) Ash and acid detergent fiber (ADF) according to AOAC (2000) procedures. Also, the determination of neutral detergent fiber (NDF) was measured by Van Soest *et al.* (1991) method.

Statistical Analysis

The data underwent analysis of variance (ANOVA) using a completely randomized design. Mean comparisons were performed using Duncan's 1995 test, with a statistical significance level of up to 5%, utilizing the PROC MIXED module in SAS 9.4 software (SAS, 1998). The analysis followed the model outlined below:

$$\hat{Y}_{ij} = \mu + T_i + \epsilon_{ij}$$

\hat{Y}_{ij} : represents the observed value of the dependent variable, μ : is the overall mean, T_i : signifies the effect of the treatment, ϵ_{ij} : accounts for the experimental random error associated with each observation.

RESULTS

The results of the effect of different levels of whole and ground flaxseed on feed intake, the ewes' weight, and the birth weight of their lambs are given in Table 3. During the experimental period, different levels of whole and ground flaxseed supplementation in late pregnancy had no significant effect on feed intake, change the ewes' weight or birth weight of their lambs ($P > 0.05$).

Results from Table 4 revealed that whole and ground flaxseed supplementation in late pregnancy had no significant effect on DM, OM, ADF, and NDF digestibility ($P > 0.05$), but the CP digestibility was significantly ($P < 0.05$) higher in diets containing whole and ground flaxseed compared to the control diet. The CP digestibility in treatments with whole flaxseed (especially the 15%) was significantly better than in treatments with ground flaxseed. According to the results in Table 4, the EE digestibility in ewes fed whole and ground flaxseed diets improved compared to those fed the control diet ($P < 0.05$). In the treatment with 15% whole flaxseed, the highest amount of EE digestibility in ewes was observed.

The effects of the levels and physical form of flaxseed on the blood metabolites of pregnant ewes are inserted in Table 5. According to Table 5, the blood glucose and cholesterol levels of ewes were affected by experimental treatments ($P < 0.05$), so that the highest blood glucose and cholesterol levels were observed in the treatment of 15% whole flaxseed. The blood triglycerides and albumin levels of ewes were not affected by experimental treatments ($P > 0.05$). The application of whole or ground flaxseed had no significant effect on ewes' yield, fat, protein, or solids non-fat (SNF) of colostrum in ewes supplemented with whole flaxseed or ground flaxseed ($P > 0.05$; Table 6), but colostrum protein was significantly increased in the ewes fed with whole flaxseed compared to other animals ($P < 0.05$). The highest colostrum protein was observed in the treatment of 15% whole flaxseed.

Table 3 The effect of feeding whole flaxseed (WFS) or ground flaxseed (GFS) on feed intake, ewe's body weight change of ewes in late pregnancy, and lambs birth weight

Parameter	Treatments					SEM	P-value
	Control	WFS-L	WFS-H	GFS-L	GFS-H		
Feed intake ewes	1.66	1.77	1.80	1.73	1.76	0.05	0.56
Ewe live weight (kg)							
56 days before lambing	56.50	54.15	55.25	57	55.70	2.65	0.94
1 h after lambing	54.75	52.10	53.67	55.37	54.15	2.58	0.90
Ewe weight loss (kg)	1.75	2.05	1.57	1.62	1.55	0.69	0.98
Lamb birth weight (kg)	4.80	5.42	5.55	5	5.13	0.20	0.21

SEM: Standard error.

NS: Non-significant.

Table 4 Effects of feeding whole flaxseed (WFS) or ground flaxseed (GFS) on apparent digestibility of ewes in late pregnancy

Parameter	Treatments					SEM	P-value
	Control	WFS-L	WFS-H	GFS-L	GFS-H		
Digestibility (% DM)							
DM	69.72	68.68	75.41	72.65	70.40	2.33	0.32
OM	69.62	71.68	73.72	71.50	70.92	2.09	0.73
CP	69.68 ^c	83.48 ^a	85.10 ^a	74.92 ^b	75.35 ^b	1.52	0.001
EE	66.29 ^b	77.06 ^a	80.61 ^a	74.57 ^a	76.56 ^a	2.27	0.01
NDF	44.46	53.44	46.47	43.50	40.18	3.86	0.24
ADF	34.46	40.11	36.47	33.50	30.84	3.32	0.41

SEM: Standard error.

NS: Non-significant.

Table 5 Effects of fed diets containing whole flaxseed or ground flaxseed on blood plasma biochemical metabolites of ewes in late pregnancy

Parameter (mg/dl)	Treatments					SEM	P-value
	Control	WFS-L	WFS-H	GFS-L	GFS-H		
Glucose	60.75 ^b	70.62 ^a	72.09 ^a	66.87 ^a	68.51 ^a	1.67	0.006
Triglycerides	34.53	35.86	37.84	36.98	37.32	1.24	0.39
Cholesterol	69.59 ^b	75.13 ^a	78.84 ^a	75.07 ^a	76.00 ^a	1.17	0.003
Albumin	3.77	3.54	3.69	3.64	3.53	0.09	0.38

SEM: Standard error.

NS: Non-significant.

Table 6 Effects of fed diets containing whole flaxseed or ground flaxseed on yield and composition of colostrum collected 1 h after lambing of ewes

Parameter	Treatments					SEM	P-value
	Control	WFS-L	WFS-H	GFS-L	GFS-H		
yield (g/d)	290	295	298.25	292.25	290.50	34.34	0.99
Fat (%)	11.87	13.42	14.12	12.12	12.25	1.26	0.67
Protein (%)	13.25 ^b	16.75 ^a	17.75 ^a	14.50 ^b	14.75 ^b	0.54	0.001
SNF (%)	25.72	27.48	28.36	27.87	27.68	1.25	0.63
Total fat (g/d)	34.52	40.25	42.84	32.89	35.08	5.09	0.61
Total protein (g/d)	38.82	49.56	53.32	42.72	42.49	6.09	0.45
SNF (g/d)	57.44	63.01	67.75	61.89	67.73	4.45	0..50

SEM: Standard error.

NS: Non-significant.

SNF: solids non-fat.

DISCUSSION

Consistent with our results, Danesh Mesgaran and Stern (2005) reported that fat in the type of oil seeds had no effect on DMI. Also, similar to our results, in a study conducted by Bell *et al.* (2006), fish oil supplementation as another rich source of PUFA, administered 21 days before lambing up to 10 days after lambing, had no effect on DMI in ewes. Gandra *et al.* (2016) reported that the DMI of cows fed whole flaxseed was greater than that of those fed no flaxseed during the prepartum period. Also, Meteab *et al.* (2020) observed that the application of flaxseed in diet increased DMI during the lactation period and suggested that this may be attributed to the increment in nutrient digestibility, which promotes rumen discharge and consequently forces the animal to eat a lot. Grummer *et al.* (2004) stated that fat supplementation increases dietary energy density, and if the DMI is not altered, the NEL intake would be greater and, consequently, could diminish the negative energy balance of early lactating cows (Gandra *et al.* 2016).

In the present study, flaxseed supplementation (especially the whole form) improved CP digestibility. The protein in flaxseed is a matrix that surrounds the fat droplets, leading to protection of the protein from deterioration

in the rumen and, therefore, greater availability of protein in the intestinal tract, thus increasing its digestibility (Khorasani et al. 1992). Also, oilseeds have been shown to increase the secretion and activity of pancreatic enzymes, enhancing digestion and absorption of proteins in the small intestine and the efficiency of protein utilisation (Mir et al. 2000).

The higher digestibility of fat increased in the current study with the inclusion of dietary flaxseeds. These results are similar to the results of Micek et al. (2004), who reported higher ether extract intake in groups containing 10% crushed linseed supplemental as compared to the control groups. Also, this result was similar to the finding of Alcalde et al. (2011) who reported that nutrient intake increased in goat kids fed flaxseed, sunflower, or canola. Although in the current study flaxseed supplementation had no effect on the digestibility of NDF and ADF, there was a tendency to increase the digestibility of these nutrients. This may be attributed to the influence of flaxseed inclusion, which is characterised by a small, flat, and oval-shaped (2×5 mm); consequently, it has a higher possibility of escaping from mastication as well as an increased passage rate from the rumen (Metiab et al. 2019). Moreover, the flaxseed fat contents are protected by the seed coats, which gives partial protection against a microbial attack or limits the negative effects of oil on ruminal microbes or both, leading to a negligible effect on the digestion of fibres (Kim et al. 2004). Similarly, another's studies showed that the reduction in fibre digestibility in the animals supplemented with 10 and 5% full-fat soybean compared to the animals supplemented with 10% flaxseed may have been that the fats in full-fat soy are not protected and have a negative effect on rumen function and cellulolytic bacteria, which led to decreased fibre digestion, whereas on the contrary, for flaxseed, the fat is protected (Khorasani et al. 1992; Kim et al. 2004). In agreement with our findings, Wachira et al. (2000) concluded that diets with 10.5% flaxseed had no significant effect on NDF digestion in ewes. In contrast with the results, da Silva et al. (2007) reported that cows fed whole flaxseed showed higher digestibility of ADF but lower CP and EE digestibility than treatments with ground flaxseed. These differences may be due to the animal's species used, flaxseed level, and diet composition.

Increases in blood glucose concentrations in ewes fed flaxseed are in agreement with experiments. Ababakri *et al.* (2021) observed that the ewes fed flaxseed had a higher glucose concentration than other groups during pre-lambing periods, and Gandra *et al.* (2016) reported that feeding whole flaxseed to cows increased blood glucose in transition dairy cows. Glycerol from fat hydrolysis in oilseeds was converted to propionate, which increased serum glucose through the process of gluconeogenesis (Hess *et al.* 2008). Also, Qin *et al.* (2018) reported that the increase in glucose concentrations in fat-supplemented groups might be a consequence of higher somatotropin concentrations, as somatotropin was found to stimulate hepatic gluconeogenesis to supply the energy demand of the lactating mammary gland (Knapp *et al.* 1992). The findings of the present study showed that feeding whole and ground flaxseed increased cholesterol levels in ewes. In agreement with the results of this study, Ye *et al.* (2009) reported that the inclusion of 2% flaxseed in Holstein dairy cows resulted in an increase in cholesterol and triglyceride concentrations. The increased cholesterol concentrations could be due to a decrease in the circulation of low-density lipoprotein cholesterol in the blood via flaxseed supplementation (Pan *et al.* 2009). In contrast with the results of this study, Metiab *et al.* (2019) reported that flaxseed supplementation in goat diets during the gestation period reduced cholesterol and triglyceride levels.

In the present study, the application of whole flaxseed increased colostrum protein concentration in experimental ewes. Similar to our results, Nudda *et al.* (2013) reported that flaxseed supplementation in goats led to increased milk protein concentrations as a result of higher protein availability in the intestine. Also, Metiab *et*

al. (2020) suggested that milk protein concentration was increased in goats fed 10% flaxseed and could be due to the improved digestibility. Mir et al. (2000) stated that oil seeds increase the secretion and activity of pancreatic enzymes, enhancing digestion and absorption of proteins in the small intestine and the efficiency of protein utilization. Petit (2003) reported no change in milk yield with the feeding of whole flaxseed. In addition, Ababakri et al. (2021) observed that during pre-lambing periods, the ewes fed whole and extruded flaxseed had no effect on the yield, fat, or protein content of the colostrum.

CONCLUSIONS

In general, the results of this study showed that ewes fed 15% whole flaxseed had the best CP and EE digestibility and the highest glucose and cholesterol in the blood plasma. Also, 15% whole flaxseed increases protein concentration in colostrum ewes, which may be beneficial to immunity in their offspring. Therefore, 15% whole flaxseed can be added to sheep's diets in late pregnancy.

ACKNOWLEDGEMENT

The authors thank all the teams who worked on the experiments and provided results during this study.

CONFLICT OF INTERESTS

The authors report no declarations of interest.

REFERENCES

- Ababakri R., Dayani O., Khezri A. and Naserian, A.A. (2021). Influence of flaxseed with rumen undegradable protein level on milk yield, milk fatty acids and blood metabolites in transition ewes. *J. Anim. Sci. Technol.* **63** (3), 475-490.
- Alcalde C.R., Grande P.A., Lima L.S., Macedo F.A.F., Zeoula L.M. and Paula, M.C. (2011). Oilseeds in feeding for growing and finishing $\frac{3}{4}$ Boer + $\frac{1}{4}$ Saanen goat kids. *Revista Brasileira de Zootecnia.* **40**, 1753-1757.
- AOAC. (2000). Official Methods of Analysis. 17th Ed. Association of Official Analytical Chemists, Arlington, Washington, DC., USA.
- Bell J.A., Griinari J.M. and Kennelly J.J. (2006). Effect of Safflower Oil, Flaxseed Oil, Monensin, and Vitamin E on Concentration of Conjugated Linoleic Acid in Bovine Milk Fat. *J. Dairy Sci.* **89**, 733-748.
- Da Silva D.C., Santos G.T.D., Branco A.F., Damasceno J.C., Kazama R., Matsushita M., Horst J.A., Dos Santos W.B.R. and Petit H.V. (2007). Production performance and milk composition of dairy cows fed whole or ground flaxseed with or without monensin. *J. Dairy Sci.* **90**, 2928-2936.
- Danesh Mesgaran M. and Stern M.D. (2005). Ruminal and post-ruminal protein disappearance of various feeds originating from Iranian plant varieties determined by the in situ mobile bag technique and alternative methods. *Anim Feed Sci. Technol.* **118**, 31-46.
- Dawson L.E.R., Carson A.F. and Kilpatrick D.J. (1999). The effect of digestible undegradable protein concentration of concentrates and protein source offered to ewes in late pregnancy on colostrum production and lamb performance. *J. Anim. Sci. Technol.* **82**, 21-36.

- Farmer C., Giguere A. and Lessard M. (2010). Dietary supplementation with different forms of flax in late gestation and lactation: Effects on sow and litter performances, endocrinology, and immune response. *J. Anim. Sci.* **88**, 225-237.
- Gandra J.R., Barletta R.V., Mingoti R.D., Verdurico L.C., Freitas Jr J.E., Oliveira L.J., Takiya C.S., Kfoury Jr J.R., Wiltbank M.C. and Renno F.P. (2016). Effects of whole flaxseed, raw soybeans, and calcium salts of fatty acids on measures of cellular immune function of transition dairy cows. *J. Dairy Sci.* **99**, 4590–4606.
- Greco L.F., Neves Neto J.T., Pedrico A., Ferrazza R.A., Lima F.S., Bisinotto R.S., Martinez N., Garcia M., Ribeiro E.S., Gomes G.C., Shin J.H., Ballou M.A., Thatcher W.W., Staples C.R. and Santos J.E.P. (2015). Effects of altering the ratio of dietary n-6 to n-3 fatty acids on performance and inflammatory response to a lipopolysaccharide challenge in lactation Holstein cows. *J. Dairy Sci.* **98**, 602–617.
- Grummer R. R., Mashek D. G. and Hayirli A. (2004). Dry matter intake and energy balance in the transition period. *Vet. Clin. N. Am. Food Anim.* **20**, 447–470.
- Hashemi M., Zamiri M.J. and Safdarian, M. (2008). Effects of nutritional level during late pregnancy on c colostral production and blood immunoglobulin levels of Karakul ewes and their lambs. *Small Rumin. Res.* **75**, 204-209.
- Hess B.W., Moss G.E. and Rule D.C. (2008). A decade of developments in the area of fat supplementation research with beef cattle and sheep. *J. Anim. Sci.* **86**, 188-204.
- Karakus F. and Atmac M. (2016). The effect of ewe body condition at lambing on growth of lambs and colostral specific gravity. *Arch. Anim. Breed.* **59**, 107-112.
- Khorasani G.R., De Boer G., Robinson P.H. and Kennelly J.J. (1992). Effect of canola fat on ruminal and total tract digestion, plasma hormones, and metabolites in lactating dairy cows. *J. Dairy Sci.* **75**, 492-501.
- Kim C.M., Kim J.H., Chung T.Y. and Park K.K. (2004). Effects of flaxseed diets on fattening response of hanwoo cattle: 2. fatty acid composition of serum and adipose tissues. *J. Anim. Sci.* **17(9)**, 1246-1254.
- Knapp J.R., Freetly H.C., Reis B.L., Calvert C.C. and Baldwin R.L. (1992). Effects of somatotropin and substrates on patterns of liver metabolism in lactating dairy cattle. *J. Dairy Sci.* **75**, 1025-1035.
- Meteab M., El-Sayed H.M., Abeer M.E.E., Nassar M.S. and El-Bordeny N.E. (2019). Impact of flaxseed inclusion in damascus goats ration on digestibility, some blood parameters and reproductive performance. *Arab Univ. J. Agric. Sci.* **5 (1)**, 2347-2357.
- Meteab M.I., El-Sayed H.M., Abeer M.E.E., Nassar M.S. and El-Bordeny N.E. (2020). Effect of dietary incremental levels of flaxseed supplementation on productive performance of lactating Damascus goats. *J. Environ. Agric. Biotech.* **5**, 181-191.
- Meteab M., El-Sayed H.M., Abeer M.E.E., Nassar M.S. and El-Bordeny N.E. (2019). Impact of flaxseed inclusion in damascus goats ration on digestibility, some blood parameters and reproductive performance. *Arab Univ. J. Agric. Sci.* **5(1)**, 2347-2357.
- Micek P., Borowiec F. and Marciński M. (2004). Linseed-based diets for sheep. 1. Nutrient digestibility, N retention and rumen fermentation. *J. Anim. Feed Sci.* **2**, 15-18.
- Mir P.S., Mears G.J., Okine E.K., Entz T., Ross C.M., Husar S.D. and Mir Z. (2000). Effects of increasing dietary grain on viscosity of duodenal digesta and plasma hormone, glucose and amino acid concentrations in steers. *Can. J. Anim. Sci.* **80**, 703-712.
- NRC. 2001. Nutrient Requirements of Dairy Cattle. 7th rev. ed. Natl. Acad. Sci., Washington, DC.

- Nudda A., Correddu F., Marzano A., Battacone G., Nicolussi P., Bonelli P. and Pulina G. (2015). Effects of diets containing grape seed, linseed, or both on milk production traits, liver and kidney activities, and immunity of lactating dairy ewes. *J. Dairy Sci.* **98**, 1157-1166.
- Nudda G., Battacone A., Atzori S., Dimauro C., Rattu S.P.G., Nicolussi P., Bonelli P. and Pulina G. (2013). Effect of extruded linseed supplementation on blood metabolic profile and milk performance of Saanen goats. *J. Animal consortium.* **7 (9)**, 1464-1471.
- Ocak N., Cam M.A. and Kuran M. (2005). The effect of high dietary protein levels during late gestation on colostrum yield and lamb survival rate in singleton-bearing ewes. *Small Rumin. Res.* **56**, 89-94.
- Pan A., Yu D., Demark-Wahnefried W., Franco O.H. and Lin X. (2009). Meta-analysis of the effects of flaxseed interventions on blood lipids. *The American Journal of Clinical Nutrition (AJCN)*. **90 (2)**, 288-297.
- Petit H.V. (2003). Digestion, milk production, milk composition and blood composition of dairy cows fed formaldehyde treated flaxseed or sunflower seed. *J. Dairy Sci.* **86**, 2637-2646.
- Petit H.V. (2010). Review: feed intake, milk production and milk composition of dairy cows fed flaxseed. *Can. J. Anim. Sci.* **90**, 115-27.
- Petit H.V. and Benchaar C. (2007). Milk production, milk composition, blood composition and conception rate of transition dairy cows fed different profiles of fatty acids. *Can. J. Anim. Sci.* **87**, 591-600.
- Qin N., Bayat A.R., Trevisi E., Minuti A., Kairenius P., Viitala S., and et al. (2018). Dietary supplement of conjugated linoleic acids or polyunsaturated fatty acids suppressed the mobilization of body fat reserves in dairy cows at early lactation through different pathways. *J. Dairy. Sci.* **101**, 7954–7970.
- Rosa Velazquez M., Fernanda Batistel F. Rodriguez J.M.P. and Relling A.E. (2020). Effects of maternal dietary omega-3 polyunsaturated fatty acids and methionine during late gestation on fetal growth, DNA methylation, and mRNA relative expression of genes associated with the inflammatory response, lipid metabolism and DNA methylation in placenta and offspring's liver in sheep. *J. Anim. Sci. Biotechnol.* **11 (111)**, 2-15.
- SAS, 1998. SAS/STAT guide to personal computers, version 7. Statistical Analysis System Institute, Cary.
- SRNS. (2010). Small ruminant nutrition system model is the result of a joint collaboration among Texas A and M University, Cornell University.
- Van Keulen J. and Young B.A. (1977). Evaluation of acid-insoluble ash as a natural marker in ruminant digestibility studies. *J. Anim. Sci.* **44**, 282-287.
- Van Soest, P. J., Robertson J. B. and Lewis B. A. (1991). Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *J. Dairy Sci.* **74**:3583–3597.
- Wachira A.M., Sinclair L.A., Wilkinson R.G., Hallett K., Enser M. and Wood J.D. (2000). Rumen biohydrogenation of n-3 polyunsaturated fatty acids and their effects on microbial efficiency and nutrient digestibility in sheep. *J. Agric. Sci.* **135**, 419-428.
- Xu J., Gao H., Song L., Yang W., Chen C., Deng Q., Huang Q., Yang J. and Huang F. (2013). Flaxseed oil and α -lipoic acid combination ameliorates hepatic oxidative stress and lipid accumulation in comparison to lard. *Lipids Health Dis.* **1**, 12-58.
- Ye J., Wang C., Wang H., Ye H., Wang B., Liu H., Wang Y., Yang Z. and Liu J. (2009). Milk production and fatty acid profile of dairy cows supplemented with flaxseed oil, soybean oil, or extruded soybeans. *Acta Agric. Scand. A. Anim.* **59 (2)**, 121-129.

