



New lower bound for numerical radius for off-diagonal 2×2 matrices

B. Moosavi^{a,*}, M. Shah Hosseini^b

^aDepartment of Mathematics, Safadasht Branch, Islamic Azad University, Tehran, Iran.

^bDepartment of Mathematics, Shahr-e-Qods Branch, Islamic Azad University, Tehran, Iran.

Received 30 November 2023; Revised 26 January 2024; Accepted 30 January 2024.

Communicated by Vishnu Narayan Mishra

Abstract. New norm and numerical radius inequalities for operators on Hilbert space are given. Among other inequalities, we prove that if $A, B \in B(H)$, then

$$\|A\| - \frac{3\|A - B^*\|}{2} \leq \omega \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right).$$

Moreover, $\omega(AB) \leq \frac{3}{2}\|Im(A)\| \|B\| + D_B \omega(A)$. In particular, if A is self-adjointable, then $\omega(AB) \leq D_B \|A\|$, where $D_B = \inf_{\lambda \in \mathbb{C}} \|B - \lambda I\|$.

Keywords: Bounded linear operator, Hilbert space, norm inequality, numerical radius.

2010 AMS Subject Classification: 47A12, 47A30, 47A63.

1. Introduction and preliminaries

Let $B(H)$ denote the C^* -algebra of all bounded linear operators on a complex Hilbert space H with inner product $\langle \cdot, \cdot \rangle$. The numerical radius of $A \in B(H)$ is defined by $\omega(A) = \sup\{ |\langle Ax, x \rangle| : \|x\| = 1 \}$. In [11], Yamazaki proved for any $A \in B(H)$ that $\omega(A) = \sup_{\theta \in \mathbb{R}} \|Re(e^{i\theta} A)\|$. It is well known that $\omega(\cdot)$ is a norm on $B(H)$ which is equivalent to the usual operator norm $\|\cdot\|$. In fact, for all $A \in B(H)$,

$$\frac{\|A\|}{2} \leq \omega(A) \leq \|A\|. \quad (1)$$

*Corresponding author.

E-mail address: baharak_moosavie@yahoo.com (B. Moosavi); mohsen_shahhosseini@yahoo.com (M. Shah Hosseini).

The first inequality becomes an equality if $A^2 = 0$. The second inequality becomes an equality if A is normal. Several numerical radius inequalities improving the inequalities in (1) have been recently given in [2, 6, 8, 9].

If A and B are operators in $B(H)$, we write the direct sum $A \oplus B$ for the 2×2 operator matrix $\begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix}$, regarded as an operator on $H \oplus H$. Thus, $\omega(A \oplus B) = \max(\omega(A), \omega(B))$. Also,

$$\|A \oplus B\| = \left\| \begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right\| = \max(\|A\|, \|B\|).$$

Some numerical radius inequalities for certain 2×2 operator matrices is obtained in [4]. More precisely,

$$\sqrt[n]{\max(\omega((AB)^n), \omega((BA)^n))} \leq \omega \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) \leq \frac{\|A\| + \|B\|}{2}$$

for $n = 1, 2, \dots$. In [5], Holbrook showed for any $A, B \in B(H)$ that $\omega(AB) \leq 4\omega(A)\omega(B)$. In the case $AB = BA$, then $\omega(AB) \leq 2\omega(A)\omega(B)$. In [3], it is shown for any $A, B \in B(H)$ that

$$\omega(A^*B \pm BA) \leq 2\|A\|\omega(B). \quad (2)$$

Let $D_A = \inf_{\lambda \in \mathbb{C}} \|A - \lambda I\|$ and let R_A denote the radius of the smallest disk in the complex plane containing $\sigma(A)$ (the spectrum of A). Stampfli in [10] proved that if $A \in B(H)$ and A is normal, then $D_A = R_A$.

The question about the best constant k such that the inequality

$$\omega(AB) \leq k\|A\|\omega(B) \quad (3)$$

holds for all operators $A, B \in B(H)$ is still open.

Concerning the inequality (3), it is shown in [1] that if $A, B \in B(H)$, then $\omega(AB) \leq (\|A\| + D_A)\omega(B)$ and

$$\omega(AB) \leq \|A\|\omega(B) + \frac{1}{2}\omega(B^*A^* - AB^*). \quad (4)$$

Also, if $A > 0$, then $\omega(AB) \leq \frac{3}{2}\|A\|\omega(B)$.

In Section 2, we introduce some inequalities between the operator norm and the numerical radius of operators on Hilbert spaces. Particularly, we establish lower bound for the numerical radius of the off-diagonal parts of 2×2 operator matrices.

2. Main results

In order to derive our main results, we need the following lemma. The lemma, which can be found in [7], gives new numerical radius inequalities for products of two Hilbert space operators.

Lemma 2.1 Let $A, B \in \mathbb{B}(\mathbb{H})$. Then

$$w(AB) \leq \omega(A)\omega(B) + D_A D_B. \tag{5}$$

The following result may be as well.

Theorem 2.2 If $A, B, C \in \mathbb{B}(\mathbb{H})$, then

$$\|Re(CA)\| \leq \frac{3\|A - B^*\|\|C\|}{4} + \frac{1}{2}\omega(CA + BC^*).$$

Proof. Clearly, $\|Re(AB)\| = \omega(Re(AB))$. Then

$$\begin{aligned} \|Re(AB)\| &= \omega(Re(AB)) \\ &= \omega\left(\frac{AB + B^*A^*}{2}\right) \\ &\leq \frac{1}{2}\omega(A(B + B^*)) + \frac{1}{2}\omega(AB - BA^*) \\ &\leq \frac{\|B + B^*\|\omega(A)}{2} + \frac{1}{2}D_A D_{B+B^*} + \frac{1}{2}\omega(AB - BA^*) \quad (\text{by (5)}) \\ &\leq \frac{\|B + B^*\|}{2}(\omega(A) + \|A\|) + \frac{1}{2}\omega(AB - BA^*). \end{aligned}$$

Hence,

$$\|Re(AB)\| \leq \frac{\|B + B^*\|}{2}(\omega(A) + \|A\|) + \frac{1}{2}\omega(AB - BA^*). \tag{6}$$

Now, let $A_1 = \begin{bmatrix} 0 & C \\ 0 & 0 \end{bmatrix}$ and $B_1 = \begin{bmatrix} 0 & -B \\ A & 0 \end{bmatrix}$. By (6), we have

$$\|Re(A_1 B_1)\| \leq \frac{\|B_1 + B_1^*\|}{2}(\omega(A_1) + \|A_1\|) + \frac{1}{2}\omega(A_1 B_1 - B_1 A_1^*)$$

and so

$$\begin{aligned} \|Re(CA)\| &= \left\| Re\left(\begin{bmatrix} CA & 0 \\ 0 & 0 \end{bmatrix}\right) \right\| \\ &= \|Re(A_1 B_1)\| \\ &\leq \frac{1}{2} \left\| \begin{bmatrix} 0 & A^* - B \\ A - B^* & 0 \end{bmatrix} \right\| (\omega(A_1) + \|A_1\|) + \frac{1}{2}\omega\left(\begin{bmatrix} CA + BC^* & 0 \\ 0 & 0 \end{bmatrix}\right) \\ &= \frac{1}{2}\|A^* - B\|(\omega(A_1) + \|C\|) + \frac{1}{2}\omega(CA + BC^*). \end{aligned}$$

Consequently,

$$\|Re(CA)\| \leq \frac{1}{2}\|A^* - B\|(\omega(A_1) + \|C\|) + \frac{1}{2}\omega(CA + BC^*). \tag{7}$$

Since $A_1^2 = 0$ and $\omega(A_1) = \frac{\|C\|}{2}$, the result follows from (7). ■

The following result may be as well.

Corollary 2.3 If $A, B \in \mathbb{B}(\mathbb{H})$, then

$$\|A\| - \frac{3\|A - B^*\|}{2} \leq \omega \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right).$$

Proof. Replacing C by I in Theorem 2.2 gives

$$\|Re(A)\| \leq \frac{3\|A - B^*\|}{4} + \frac{1}{2}\omega(A + B). \quad (8)$$

Now, let

$$A_1 = \begin{bmatrix} 0 & A \\ 0 & 0 \end{bmatrix} \quad \text{and} \quad B_1 = \begin{bmatrix} 0 & 0 \\ B & 0 \end{bmatrix}.$$

By (8), we have

$$\begin{aligned} \frac{\|A\|}{2} &= \|Re(A_1)\| \\ &\leq \frac{3}{4}\|A_1 - B_1^*\| + \frac{1}{2}\omega(A_1 + B_1) \\ &\leq \frac{3}{4}\left\| \begin{bmatrix} 0 & A - B^* \\ 0 & 0 \end{bmatrix} \right\| + \frac{1}{2}\omega \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) \end{aligned}$$

and so

$$\|A\| \leq \frac{3\|A - B^*\|}{2} + \omega \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right).$$

This completes the proof. ■

As a natural application of the above Corollary in providing upper bounds for the nonnegative quantity $\|A\| - \omega(A)$, $A \in \mathbb{B}(\mathbb{H})$, we can state the following result:

Corollary 2.4 If $A, B \in \mathbb{B}(\mathbb{H})$, then $\|A\| - \omega(A) \leq 3\|Im(A)\|$.

Proof. Replacing B by A in Theorem 2.3, we deduce the desired result. ■

The following result may be as well.

Theorem 2.5 If $A, C \in \mathbb{B}(\mathbb{H})$, then $\omega(CA) \leq \frac{3}{2}\|Im(A)\|\|C\| + D_C \omega(A)$.

Proof. By Theorem 2.2,

$$\|Re(CA)\| \leq \frac{3\|A - B^*\|\|C\|}{4} + \frac{1}{2}\omega(CA + BC^*).$$

Replacing B by A in the last inequality gives

$$\begin{aligned} \|Re(CA)\| &\leq \frac{3\|A - A^*\| \|C\|}{4} + \frac{1}{2}\omega(CA + AC^*) \\ &= \frac{3}{2}\|Im(A)\| \|C\| + \frac{1}{2}\omega(CA + AC^*). \end{aligned}$$

Therefore,

$$\|Re(CA)\| \leq \frac{3}{2}\|Im(A)\| \|C\| + \frac{1}{2}\omega(CA + AC^*).$$

Let $\alpha_0 = \frac{\bar{z}_0}{|z_0|}$, where $z_0 \in \mathbb{C}$ is such that $\|C - z_0I\| = D_C$. Replacing C by α_0C in the inequality (6) gives

$$\begin{aligned} \|Re(\alpha_0CA)\| &\leq \frac{3}{2}\|Im(A)\| \|C\| + \frac{1}{2}\omega(\alpha_0CA - \bar{\alpha}_0CA^*) \\ &\leq \frac{3}{2}\|Im(A)\| \|C\| + \frac{1}{2}\omega(\alpha_0(C - z_0I)A - \bar{\alpha}_0A(C - z_0I)^*) \\ &\leq \frac{3}{2}\|Im(A)\| \|C\| + \|C - z_0I\|\omega(A). \end{aligned} \tag{by (2)}$$

Thus,

$$\|Re(\alpha_0CA)\| \leq \frac{3}{2}\|Im(A)\| \|C\| + D_C \omega(A). \tag{9}$$

On the other hand, there exists $\theta_0 \in \mathbb{R}$ such that $\alpha_0 = e^{i\theta_0}$. Let $\theta \in R$ and replacing C by $e^{i\theta}C$ in the inequality (9) gives

$$\|Re(e^{i(\theta+\theta_0)}CA)\| \leq \frac{3}{2}\|Im(A)\| \|C\| + D_C \omega(A).$$

Taking the supremum over $\theta \in R$ gives

$$\omega(CA) \leq \frac{3}{2}\|Im(A)\| \|C\| + D_C \omega(A),$$

which is exactly the desired result. ■

The following corollary are immediate consequences of Theorem 2.5.

Corollary 2.6 If $A, C \in \mathbb{B}(\mathbb{H})$ and A is self-adjointable, then $\omega(CA) \leq D_C\|A\|$.

The following theorem is a considerable improvement of the inequality (4).

Theorem 2.7 If $A, B \in \mathbb{B}(\mathbb{H})$, then $\omega(AB) \leq D_A\omega(B) + \frac{1}{2}\omega(AB - BA^*)$.

Proof. Let $\theta \in \mathbb{R}$. We have

$$\begin{aligned} \|Re(e^{i\theta} AB)\| &= \omega(Re(e^{i\theta} AB)) \\ &= \omega\left(\frac{e^{i\theta} AB + e^{-i\theta} B^* A^*}{2}\right) \\ &= \omega\left(\frac{A(e^{i\theta} B + e^{-i\theta} B^*)}{2} + \frac{e^{-i\theta}(B^* A^* - AB^*)}{2}\right) \\ &\leq \frac{1}{2} D_A \|e^{i\theta} B + e^{-i\theta} B^*\| + \frac{1}{2} \omega(B^* A^* - AB^*) \quad (\text{by Corollary(2.6)}) \\ &= D_A \|Re(e^{i\theta} B)\| + \frac{1}{2} \omega(B^* A^* - AB^*). \end{aligned}$$

Consequently,

$$\|Re(e^{i\theta} AB)\| \leq D_A \|Re(e^{i\theta} B)\| + \frac{1}{2} \omega(B^* A^* - AB^*).$$

Taking the supremum over $\theta \in R$ gives $\omega(AB) \leq D_A \omega(B) + \frac{1}{2} \omega(B^* A^* - AB^*)$, which is exactly the desired result. \blacksquare

The following corollary are immediate consequences of Theorem 2.7.

Corollary 2.8 If $A, B \in \mathbb{B}(\mathbb{H})$ and $AB = BA^*$, then $\omega(AB) \leq D_A \omega(B)$.

Acknowledgments

The authors thank the Editorial Board and referees for their valuable comments that helped to improve the article.

References

- [1] A. Abu-Omar, F. Kittaneh, Numerical radius inequalities for products of Hilbert space operators, J. Operator Theory. 72 (2) (2014), 521-527.
- [2] S. S. Dragomir, Some inequalities for the norm and the numerical radius of linear operators in Hilbert Spaces, Tamkang J. Math. 39 (1) (2008), 1-7.
- [3] C. K. Fong, J. A. Holbrook, Unitarily invariant operator norms, Canad. J. Math. 35 (1983), 274-299.
- [4] O. Hirzallah, F. Kittaneh, K. Shebrawi, Numerical radius inequalities for certain 2×2 operator matrices, Integral Equ. Operator Theory. 71 (2) (2011), 129-147.
- [5] J. A. R. Holbrook, Multiplicative properties of the numerical radius in operator theory, J. Reine Angew. Math. 237 (1969), 166-174.
- [6] F. Kittaneh, Numerical radius inequalities for Hilbert space operators, Studia Math. 168 (1) (2005), 73-80.
- [7] M. Shah Hosseini, B. Moosavi, Some numerical radius inequalities for products of Hilbert space operators, Filomat. 33 (7) (2019), 2089-2093.
- [8] M. Shah Hosseini, B. Moosavi, H. R. Moradi, An alternative estimate for the numerical radius of Hilbert space operators, Math. Slovaca. 70 (1) (2020), 233-237.
- [9] M. Shah Hosseini, M. E. Omidvar, Some inequalities for the numerical radius for Hilbert space operators, Bull. Aust. Math. Soc. 94 (3) (2016), 489-496.
- [10] J. G. Stampfli, The norm of a derivation, Pacific J. Math. 33 (1970), 737-747.
- [11] T. Yamazaki, On upper and lower bounds of the numerical radius and an equality condition, Stud. Math. 178 (2007), 83-89.