



Original Article

A survey on cattle hard ticks fauna in Maragheh city, Iran

Sohrab Rasouli^{1*}, Omid Mohammadpour², Seyyed Mohammad Rahchamani²

¹Department of Pathobiology, Faculty of Veterinary Medicine, Urmia Branch, Islamic Azad University, Urmia, Iran

²Department of Clinical Sciences, Faculty of Veterinary Medicine, Shahid Bahonar University of Kerman, Kerman, Iran

ARTICLE INFO

Received: 12 May 2021

Accepted: 5 December 2021

DOI:10.30495/jbcvm.2021.1930418.1010

KEY WORDS :

Hard tick
Ixodidae family
Hyalomma
Rhipicephalus
Cattle

ABSTRACT

Ticks are the most important ectoparasites which inflict heavy economic losses to livestock and transmit viral, rickettsia, bacterial and protozoal diseases. *Ixodidae* family is one of the most great tick family of domestic animals. So far, 13 different kinds and 650 species in five subgroups have been reported. The aim of this research is to assess quantitatively the extent of cattle infestations to these different ticks and also to identify the diversity of the species found in Maragheh. In this survey, carried out throughout fall 2015 to the end of summer 2016, a total number of 328 ticks were collected and identified including 224 *Hyalomma* spp. (68.30%), *Rhipicephalus* spp. 98 (29.88%), *Dermacentor* spp. 3 (0.91%) and *Boophilus* spp. 3 (0.91%) as the minimum infestation rate. From seasonal infestation prevalence aspect, in spring, 168 ticks, in summer 89 ticks and in autumn and winter 45 and 26 ticks were respectively, found on the cattle. Maximum infestation of cattle in summer and minimum in winter were observed. The maximum rate of the found ticks was observed in groin and the minimum was observed in testicle surfaces.

بررسی فون کنه‌های سخت گاو در شهر مراغه، ایران

سهراب رسولی^{۱*}، امید محمدپور^۲، سید محمد راه چمنی^۲

^۱ گروه پاتوبیولوژی، دانشکده دامپزشکی، دانشگاه آزاد اسلامی واحد ارومیه، ارومیه، ایران

^۲ گروه علوم درمانگاهی، دانشکده دامپزشکی، دانشگاه شهید باهنر کرمان، کرمان، ایران

چکیده

هدف از مطالعه حاضر، ارزیابی گسترش آلودگی گاوهای شهرستان مراغه به گونه‌های مختلف کنه‌های سخت از نظر کمی و همچنین شناسایی تنوع گونه‌ای کنه‌های یافت شده می‌باشد. در این مطالعه که از ابتدای پائیز ۱۳۹۴ تا انتهای تابستان ۱۳۹۵ انجام شده است از کل گاوهای مورد بررسی در شهرستان مراغه، ۳۲۸ کنه جمع‌آوری و شناسایی گردید که شامل ۴ جنس و ۸ گونه بود. هیالوما آناتولیکم با فراوانی ۱۵۱ (۴۶/۰۶ درصد) بعنوان بیشترین عامل آلودگی و رپی سفالوس تورانیکوس با فراوانی ۲ (۰/۶۱ درصد) بعنوان کمترین عامل آلودگی شناسایی شدند در حالی که بعد از هیالوما آناتولیکم آناتولیکم، رپی سفالوس بورس با فراوانی ۵۸ (۱۷/۵۸ درصد)، هیالوما آناتولیکم اکسکواتوم با فراوانی ۴۱ (۱۲/۵۰ درصد)، رپی سفالوس سانگوینوس با فراوانی ۳۸ (۱۱/۵۹ درصد)، هیالوما دتریتوم با فراوانی ۳۲ (۹/۷۶ درصد) و هر یک از درماستور مارژیناتوس و بواقیلوس آنوالتوس با فراوانی ۳ (۰/۹۱ درصد) به ترتیب بالاترین آلودگی بودند. در تحقیق حاضر، از لحاظ شیوع فصلی آلودگی به کنه سخت، در بهار ۱۶۸ کنه، در تابستان ۸۹ کنه و در پاییز و زمستان به ترتیب ۴۵ و ۲۶ کنه یافته شد و بیشترین میزان آلودگی در گاوها در فصل بهار و کمترین میزان آلودگی در فصل زمستان مشاهده شد. بیشترین پراکندگی کنه در کشاله ران و کمترین پراکندگی کنه در روی بیضه‌ها مشاهده شد.

واژه‌های کلیدی: کنه سخت، خانواده ایکسودیده، هیالوما، رپی سفالوس، گاو

* Corresponding author: sohrab_rasouli86@yahoo.com

©2021 Islamic Azad University, Urmia Branch.

This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).



INTRODUCTION

Ticks are hematophagous ectoparasites that are known as pests and vectors of a wide range of diseases of humans, livestock, pets, and wild animals. It has been estimated that about 80% of the world cattle population is infested with ticks [1]. Ticks have a variety of direct and indirect effects on their hosts. Tick infestations can cause considerable irritation in animals and can lead to severe disorders, such as blood loss, general stress, damages to hide and skins, tick paralysis, and tick toxicosis [2, 3]. In Middle East, taxonomically accurate information on tick species is limited, and the tick-borne diseases of this region remain poorly characterized. The relationship between ticks and tickborne pathogens in the region is largely unknown, even though the presence of these pathogens has been recognized for many years and the number of new pathogens discovered in ticks has increased markedly [4-6]. Advances have recently been made in Middle East infectious disease research, but vector-borne diseases are still misdiagnosed and underestimated because of inadequate clinical training and limited surveillance and laboratory capacity. Tick species that occur in Iran have been discussed by Toumanoff in 1944 [7]; Hoogstraal *et al.* as well as Wilson between the 1960s and 1980s [8-13]; Petney and Keirans in the mid-1990s [14-17]; Robbins *et al.* in 1996 [18] and Kernif *et al.* in 2012 [4] the most recent work. Cattle are reared in this area by releasing them into the forest and by periodically moving them close to or under village houses. Such practices may facilitate interactions between ticks and their hosts, including the exchange of ticks between wild and domestic animals. They may also lead to dispersal of the tick population, thereby potentially increasing the risk of tick-borne disease transmission. The goal of this research is to assess quantitatively

the extent of cattle infestation to these different kinds of ticks, family from quantitative aspect and also to identify the diversity of the species found in Maragheh.

MATERIALS AND METHODS

The present study was performed carried out in 48 different rural areas of Maragheh city in Eest Azarbayjan of Iran. It is located at 37°23'21"N 46°14'15"E. Throughout a year, 12 sampling periods (each month were one period) were referenced to 48 designated geographical areas, so that at each stage 4 region (one of each west, east, north and south areas) were examined. In this study, a total number of 384 cattle (198 female and 186 male) from different age groups (less than 1 year, 1–2 years old, 2–3 years old, and over 3 years old) were selected by stratified random sampling over the course of 1 year (October 2015 to September 2016). In each stage, 8 samples were randomly selected for each of the four study areas. The examined cattle were raised under traditional husbandry practices (grazing on pastures during the day) without regular *acaricide* treatment. A total of 328 ticks were collected from cattle. Data for all specimens, including date, sex, age, and number of ticks, were recorded. All of the methods used in this study were confirmed by the Ethics Committee of Islamic Azad University of Urmia, respecting currently accepted animal welfare rules in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Helsinki Declaration of 1975, as revised in 2000 and 2008.

At first, the animal was fastened before any inspection, and then different parts of the body were examined. Due to the location of most

ticks in the animal's low-lying areas, the contractions of the perineal region, groin, under the scapula, breast, Genital organs and cattle neck were reviewed. Ticks were removed from the host with rubbing alcohol pads surrounding the skin and blunt pointed forceps, avoiding damage to the mouthparts of the ticks and the skin of host [19]. The collected specimens were transferred into holding tubes contain 70% ethanol (Merck, Darmstadt, Germany) and transferred to the Parasitology Research Laboratory of Islamic Azad University, Urmia, Iran. Following examinations under a stereomicroscope, ticks were identified by morphological characteristics using the key identification guide [20]. In some samples, 5% potash solution was used to remove sediment and clarification. For data analysis, descriptive statistics for qualitative data with 95% confidence intervals (95% CI) were used.

Collected data were analysed by Excel, v.2013 and SPSS, v.19 and the chi-square χ^2 test was run to determine the relationship between the variables (sex, different parts of the body and season).

RESULTS

According to the results, 68 of the 384 cattle (17.71 %) including 9 (13.24%) male and 59 (68.76%) female were diagnosed as being infected with ticks. 328 ticks were collected and identified. A number of 224 *Hyalomma* spp. (68.30%) determined as maximum infestation, 98 *Rhipicephalus* spp. (29.88%), 3 *Dermacentor* spp. (0.91%), 3 *Boophilus* spp. (0.91%) determined as minimum infestation rate were identified. The prevalence of tick species in cattle is examined in Table 1 and Figure 1. From seasonal infestation prevalence

Table 1. The prevalence of tick species in examined cattle in Maragheh, Iran

Tick species	Frequency of infected cattle						Frequency of tick							
	Male		Female		Total		M. Mature		F. Mature		Nymph&Larvae		Total	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<i>Hyalomma anatolicum anatolicum</i>	4	5.88	24	35.29	28	41.18	64	19.51	79	24.09	8	2.43	151	46.04
<i>Hyalomma anatolicum excavatum</i>	1	1.47	7	10.29	8	11.76	15	4.57	23	7.01	3	0.91	41	12.50
<i>Hyalomma deteritum</i>	-	0.00	7	10.29	7	10.29	10	3.05	22	6.71	-	0.00	32	9.76
<i>Rhipicephalus bursa</i>	3	4.41	8	11.76	11	16.18	18	5.49	38	11.59	2	0.61	58	17.68
<i>Rhipicephalus sanguineus</i>	1	1.47	8	11.76	9	13.24	20	6.10	18	5.48	-	0.00	38	11.59
<i>Rhipicephalus turanicus</i>	-	0.00	1	1.47	1	1.47	-	-	2	0.61	-	0.00	2	0.61
<i>Dermacentor marginatus</i>	-	0.00	3	4.41	3	4.41	-	-	3	0.91	-	0.00	3	0.91
<i>Boophilus annulatus</i>	-	0.00	1	1.47	1	1.47	1	0.30	2	0.61	-	0.00	3	0.91
Total	9	13.24	59	86.76	68	100	128	39.02	187	57.01	13	3.96	328	100




Figure 1. Relative frequency of identified ticks in cattle in Maragheh, Iran.




Figure 2. The relative frequency of the prevalence of ticks in different months in Maragheh, Iran.

Table 2. Distribution of hard tick in different parts of the body of cattle in Maragheh, Iran.

Tick genus	Frequency		Testicles		Breast		Perina		Groin	
	No.	%	No.	%	No.	%	No.	%	No.	%
<i>Hyaloma.spp</i>	224	68.30	12	5.35	31	13.84	82	36.60	99	44.20
<i>Rhipicephalus.spp</i>	98	29.88	4	4.08	13	13.26	42	42.86	39	39.80
<i>Dermacentor.spp</i>	3	0.91	0	0.00	0	0.00	1	33.33	2	66.66
<i>Boophilus.spp</i>	3	0.91	0	0.00	0	0.00	3	100	0	0.00
Total	328	100	16	4.88	44	13.41	128	39.02	140	42.68

aspect, in spring, 168 ticks, in summer 89 ticks and in autumn and winter 45 and 26 ticks were respectively, found on the cattle. Maximum infestation of cattle in summer and minimum in winter was observed. The maximum rate of the found ticks was observed in groin and the minimum were observed in testicle surfaces (Table 2 and Figure 2).

DISCUSSION

There are some studies on the prevalence of tick infestation in cattle in different parts of Iran and other countries, including 32.49%, 75.8%, and 24.63% in the Sari, Golestan, and Kermanshah regions of Iran, respectively as well as 86.1%, 72.9%, and 29.6% in Ethiopia, Pakistan, and Turkey, respectively [3, 21-39]. According to the results, 68 out of the 384 (17.71%) examined cattle were infected with hard ticks including *Hyaloma anatolicum*, *Hyaloma anatolicum excavatum*,

Hyaloma deteritum, *Rhipicephalus bursa*, *Rhipicephalus sanguineus*, *Rhipicephalus turanicus*, *Dermacentor marginatus* and *Boophilus annulatus*. In this investigation, the frequency of tick infestation was lower than that of three recent studies carried out in Turkey and Iran, where 34% and 36.90% of cattle were infested with at least one tick species [40-42]. The variation in the prevalence of tick infestation might be due to geographical distribution, climate condition, and management systems [37]. Furthermore, the methods and some other factors used in the field study could also affect the results. The data analysis indicated that there is a statistically significant difference between species. From 328 collected tick samples, *Hyalomma anatolicum anatolicum* was the most prevalent tick (46.04 %) and *Rhipicephalus turanicus* was the least (0.61%), at $P < 0.05$, whereas *Dermacentor marginatus* and *Boophilus annulatus* with (0.91%) had the very close prevalence to the minimum. It indicates the absolute dominance of the genus *Hyalomma* in particular, *Hyalomma anatolicum anatolicum*. Several research on ixodid ticks revealed that the genus *Hyalomma* was predominant in Iran [2, 43-46]. Ghashghaee *et al.* reported, *R. sanguineus* was the major tick that infested cattle, which was not consistent with our results [42].

In the current investigation, there was a significant difference between prevalence of ticks in different seasons, hard ticks were more prevalent during spring than other seasons, while the fewest were observed in winter ($p < 0.05$). The results of our study are similar to the findings by Sofizadeh *et al.* and Ghashghaee *et al.* [38, 42]. In contrast, Yakhchali and Hosseini [47] reported higher tick prevalence in winter and lower prevalence in summer seasons. It is an established fact

that climate condition and temperature affect tick prevalence [3].

CONCLUSION

In the current study, the frequency distribution of canals collected according to different parts of the body of cattle's body of the cattle was considered. According to the results, ticks related to the genus *Hyaloma* and genus *Rhipicephalus* are found in all four parts of the body of the cattle, while the genus *Dermacentor* was observed on the perineum and groin. The dispersion of the genus *Boophilus* was limited to the perineum. Between the four parts of the body, the groin (with 42.68% average inclination to ticks) had highest variation and contamination with ticks, and the surface of the testicle with 4.88% average tick inclination had the least variation and frequency of ticks ($p < 0.05$). Although many researchers have identified specific hosts for hard ticks in some cases, but there are no such consensus among a few species. Recent studies on the location of the parasite have shown that most tick species prefer specific part of the body to operate. Also, the results of the studies show that head and ear areas, groin, subcutaneous, and perineum are more exposed to *Ixodidae* ticks than other anatomical points [46].

ETHICS

All procedures of the current research have been performed based on the ethical standards.

CONFLICT OF INTEREST

There is no conflict of interest.

REFERENCES

- [1] Minjauw B, McLeod A. Tick-borne diseases and poverty. The impact of ticks and tick borne diseases

- on livestock owners in India and eastern and southern Africa. Edinburgh, UK: Health Program, Center for Tropical Veterinary Medicine, University of Edinburgh; 2003. pp. 24-57.
- [2] Razmi GR, Glinsharifodini M, Sarvi S. Prevalence of ixodid ticks on cattle in Mazandaran province, Iran. *Korean J Parasitol.* 2007; 45: 307-310.
 - [3] Sajid MS, Iqbal Z, Khan MN, Muhammad GH, Khan MK. Prevalence and associated risk factors for bovine tick infestation in two districts of lower Punjab, Pakistan. *Prev Vet Med.* 2009; 92: 386-391.
 - [4] Kernif T, Socolovschi C, Wells K, Lakim MB, Inthalad S, Slesak G, and *et al.* Bartonella and Rickettsia in arthropods from the Lao PDR and from Borneo, Malaysia. *Comparative Immunology, Microbiology and Infectious Diseases.* 2012; 35, 51–57.
 - [5] Kho KL, Koh FX, Tay ST. Molecular evidence of potential novel spotted fever group rickettsiae, Anaplasma and Ehrlichia species in Amblyomma ticks parasitizing wild snakes. *Parasites & Vectors,* 2015; 8, 112.
 - [6] Yu XJ, Liang MF, Zhang SY, Liu Y, Li JD, Sun YL, and *et al.* Fever with thrombocytopenia associated with a novel bunyavirus in China. *New England Journal of Medicine.* 2011; 364, 1523–1532.
 - [7] Toumanoff C. Les tiques (Ixodoidea) de l'Indochine. Institut Pasteur de l'Indochine, S.I.L.I., Saigon. Voltzit, O.V. & Keirans, J.E. A review of Asian Amblyomma species. *Acarina,* 1944; 10, 95–136.
 - [8] Hoogstraal H, Trapido H & Kohls, GM. Studies on Southeast Asian Haemaphysalis ticks (Ixodoidea, Ixodidae), The identity, distribution, and hosts of H. (Kaiseriana) hystricis Supino, *Journal of Parasitology.* 1965; 51: 467–480.
 - [9] Hoogstraal H & Dhanda V. Haemaphysalis (H.) darjeeling sp. n., a member of the H. (H.) birmaniae group (Ixodoidea, Ixodidae) parasitizing artiodactyl mammals in Himalayan forests of India, and in Burma and Thailand. *Journal of Parasitology.* 1970; 56: 169–174.
 - [10] Hoogstraal H, Dhanda V & Kammah, KM. Aborphysalis, a new subgenus of Asian Haemaphysalis ticks and identity, distribution, and hosts of H. aborensis Warburton (resurrected) (Ixodoidea: Ixodidae). *Journal of Parasitology.* 1971; 57: 748–760.
 - [11] Hoogstraal H, & Wassef HY. The Haemaphysalis ticks (Ixodoidea: Ixodidae) of birds. H. (Ornithophysalis) subgen. n.: definition, species, hosts, and distribution in the Oriental, Palearctic, Malagasy, and Ethiopian faunal regions. *Journal of Parasitology* 1973b; 59: 1099–1117.
 - [12] Hoogstraal H, & Wassef HY, Dermacentor (Indocentor) auratus (Acari: Ixodoidea: Ixodidae): hosts, distribution, and medical importance in tropical Asia. *Journal of Medical Entomology,* 1985b; 22: 170–177.
 - [13] Wilson N. New distributional records of ticks from Southeast Asia and the Pacific (Metastigmata: Argasidae, Ixodidae). *Oriental Insects.* 1970; 4: 37-46.
 - [14] Petney TN & Keirans JE. Ticks of the genus Ixodes in South-east Asia. *Tropical Biomedicine.* 1994; 11: 123–134.
 - [15] Petney TN & Keirans JE. Ticks of the genera Amblyomma and Hyalomma in South-east Asia. *Tropical Biomedicine.* 1995; 12: 45–46.
 - [16] Petney TN & Keirans JE. Ticks of the genera Boophilus, Dermacentor, Nosomma and Rhipicephalus (Acari: Ixodidae) in South-east Asia. *Tropical Biomedicine.* 1996a; 13: 73–84.
 - [17] Petney TN & Keirans JE. Ticks of the genus Aponomma in South-east Asia. *Tropical Biomedicine.* 1996b; 13: 167–172.
 - [18] Robbins RG, Karesh WB, Rosenberg S, Schonwalter N & Inthavong C. Two noteworthy collections of ticks (Acari: Ixodida: Ixodidae) from endangered carnivores in the Lao People's Democratic Republic. *Entomological News.* 1996; 108: 60–62.

- [19] Wall R, and Shearer D. Veterinary entomology. 1st ed. Chapman and Hall. 1997; 114-135.
- [20] Walker AR, Bouattour A, Camicas JL, Estrada-Pena A, Horak IG, Latif AA, Pegram RG, Preston MM. Ticks of Domestic Animals in Africa: A Guide to Identification of Species. 1st ed. Edinburgh, UK: Bioscience Reports. 2003; 114-135.
- [21] Parola P, Raoult D. Ticks and tickborne bacterial diseases in humans: an emerging infectious threat. Clin Infect Dis. 2001; 32: 897-928.
- [22] Nourollahi Fard SR, Khalili M. PCR-Detection of *Coxiella burnetii* in Ticks Collected from Sheep and Goats in Southeast Iran. J Arthropod Borne Dis. 2011; 5: 1-6.
- [23] Aktas M, Vatansever Z, Ozubek S. Molecular evidence for trans-stadial and transovarial transmission of *Babesia occultans* in *Hyalomma marginatum* and *Rhipicephalus turanicus* in Turkey. Vet Parasitol. 2014; 204: 369-371.
- [24] Aktas M. A survey of ixodid tick species and molecular identification of tick-borne pathogens. Vet Parasitol. 2014; 200: 276-283.
- [25] Aydin MF, Aktas M, Dumanli N. Molecular identification of *Theileria* and *Babesia* in ticks collected from sheep and goats in the Black Sea region of Turkey. Parasitol Res. 2015; 114: 65-69.
- [26] Psaroulaki A, Ragiadakou D, Kouris G, Papadopoulos B, Chaniotis B, Tselentis Y. Ticks, tickborne rickettsiae, and *Coxiella burnetii* in the Greek Island of Cephalonia. Ann NY Acad Sci. 2006; 1078: 389-399.
- [27] Leung-Shea C, Danaher PJ. Q fever in members of the United States armed forces returning from Iraq. Clin Infect Dis. 2006; 43: 77-82.
- [28] Bailey MS, Trinick TR, Dunbar JA, Hatch R, Osborne JC, Brooks TJ, Green AD. Undifferentiated febrile illnesses amongst British troops in Helmand, Afghanistan. J R Army Med Corps. 2011; 157: 150-155.
- [29] Karabay O, Gozdas HT, Ozturk G, Tuna N, Utku AC. A Q fever case mimicking Crimean-Congo haemorrhagic fever. Indian J Med Microbiol. 2011; 29: 418-419.
- [30] Rolain JM, Gourié F, Brouqui P. Concomitant or consecutive infection with *Coxiella burnetii* and tickborne diseases. Clin Infect Dis. 2005; 40: 82-88.
- [31] Rehacek J, Urvolgyi J, Kocianova E, Sekeyova Z, Vavrekova M, Kovacova E. Extensive examination of different ticks species for infestation with *Coxiella burnetii* in Slovakia. Eur J Epidemiol. 1991; 7: 299-303.
- [32] Bernasconi MV, Casati S, Peter O, Piffaretti JC. *Rhipicephalus* ticks infected with *Rickettsia* and *Coxiella* in Southern Switzerland (Canton Ticino). Infect Genet Evol. 2002; 2: 111-120.
- [33] Walker AR, Bouattour A, Camicas JL, Estrada-Pena A, Horak IG, Latif AA, Pegram RG, Preston MM. Ticks of Domestic Animals in Africa: A Guide to Identification of Species. 1st ed. Edinburgh, UK: Bioscience Reports. 2003. 114-135.
- [34] Parisi A, Fraccalvieri R, Cafiero M, Miccolupo A, Padalino I, Montagna C, Capuano F, Sottili R. Diagnosis of *Coxiella burnetii*-related abortion in Italian domestic ruminants using single-tube nested PCR. Vet Microbiol. 2006; 118: 101-106.
- [35] Mamak N, Gençer L, Özkanlar YE, Özçelik S. Determination of tick species and treatment of cows, sheep and goats in the Sivas-Zara region. Türkiye Parazitoloj Derg. 2006; 30: 209-212 (in Turkish with English abstract).
- [36] Haghi FM, Razmi G, Fakhar M, Mohammadpoor RA. The hard ticks (Ixodidae) fauna of livestock in Sari suburb, Northern Iran. Comp Clin Path. 2013; 22: 5-8.
- [37] Sohrabi S, Yakhchali M, Ghashghaei M. Hard ticks (Acarina: Ixodidae) diversity in the natural habitat of Iranian domestic ruminants: a provincial study in Kermanshah. J Vet Res Tehran Uni. 2013; 68: 39-46.
- [38] Sofizadeh A, Telmadarraiy Z, Rahnama A, Gorganli-Davaji A, Hosseini-Chegeni A. Hard tick species of livestock and their

- bioecology in Golestan province, north of Iran. *J Arthropod Borne Dis.* 2014; 8: 108.
- [39] Werede H, Afera B. Prevalence of ixodid ticks on bovine of Werieleke Wereda, Tigray. *Acta Parasitol.* 2014; 5: 146-150.
- [40] Aktas M, Altay K, Dumanli N. A molecular survey of bovine *Theileria* parasites among apparently healthy cattle and with a note on the distribution of ticks in eastern Turkey. *Vet Parasitol.* 2006; 138: 179-185.
- [41] Ghashghaei O, nourollahi-fard S, khalili M, sharifi H. A survey of ixodid ticks feeding on cattle and molecular detection of *Coxiella burnetii* from ticks in Southeast Iran. *Turk J Vet Anim Sci.* 2017; 41: 46-50.
- [42] Yakhchali M, Rostami A, Esmailzadeh M. Diversity and seasonal distribution of ixodid ticks in the natural habitat of domestic ruminants in north and south of Iran. *Rev Med Vet Toulouse.* 2011; 162: 229-235.
- [43] Hashemzade-Farhang H. Survey of tick species diversity in Tabriz and Suburbs. 2006; 43 (in Persian).
- [44] Mazlum Z. Tick species found in Iran, geographical distribution, seasons of activity and hosts. *Tehran Uni Vet J.* 1971; 2(1): 1-31 (in Persian).
- [45] Davoudi J, Hoghooghi-Rad N, Shahrokhi-Khanghah Sh. Investigation of fauna of cattle infected ticks and their seasonal variations in western Azarbayegan province. *J Daneshe novine keshavarzi.* 2007; 3(8): 15-22. (In Persian)
- [46] Yakhchali M, Hosseini A. Prevalence and ectoparasites fauna of sheep and goats flocks in Urmia suburb, Iran. *Vet Arhiv.* 2006; 76: 431-442.