

# Iranian Journal of Organic Chemistry

IranJOC Vol. 3, No. 4, 2011

## Contents

### Graphical Abstracts



#### Morphology and properties of Nylon6/ poly (butylene terephthalate) in situ microfibrillar reinforced blends: influence of blend ratio

pp 719-724

Alireza Hoseinpour,<sup>a\*</sup> Habib Tayebi,<sup>a</sup> Amir Rastar<sup>a</sup> and Ahmadali Rabienataj Darzi<sup>b</sup>

<sup>a</sup> Department of Textile Engineering, Qaemshahr Branch, Islamic Azad University, Qaemshahr, Iran

<sup>b</sup> Faculty of Mechanical Engineering, Babol University of Technology, Babol, Iran





#### Microwave assisted extraction of benzofurane derivative from *Petasites hybridus* rhizomes

pp 725-727

Faramarz Rostami-Charati<sup>\*a,b</sup>, Seyed Ali Razavi<sup>a</sup>, Mehdi Shahraki<sup>b</sup>, Ahmad Mehraban<sup>b</sup> and Mohammad R. Hosseini-Tabatabaei<sup>b</sup>

<sup>a</sup> Department of Chemistry, Faculty of Science, Gonbad Kavous University, P.O.Box 163, Gonbad, Iran

<sup>b</sup> Department of Chemistry, Islamic Azad University, Zahedan Branch, Zahedan, Iran



#### Synthesis of new 4-thiazolidinone derivatives as potential anti H-pylori


pp 729-731

Sara Kaveh<sup>\*a</sup>, Eskandar Alipour<sup>a</sup>, Alireza Foroumadi<sup>b</sup>, Abbas Shafiee<sup>c</sup> and Nouraddin Hosseinzadeh<sup>c</sup>

<sup>a</sup> Department of Chemistry, Islamic Azad University, Tehran-North Branch, Zafar St, Tehran, Iran

<sup>b</sup> Drug Design & Development Research Center, Tehran University of Medical Sciences, Tehran, Iran

<sup>c</sup> Department of Medicinal Chemistry, Faculty of Pharmacy and Pharmaceutical Sciences Research Center, Tehran University of Medical Sciences, Tehran, Iran



### Synthesis of novel 2-benzylidenebenzofuran-3(2H)-one derivatives


pp 733-736

Shiva Masoomi<sup>a</sup>, Eskandar Alipour<sup>\*a</sup>, Mohammad Ali Rasouli<sup>b</sup>, Ali Reza Foromadi<sup>c</sup> and Abbas Shafiee<sup>c</sup>

<sup>a</sup>Faculty of chemistry, Islamic Azad University, Tehran-North branch, Tehran, Iran

<sup>b</sup>School of chemistry, College of science, University of Tehran, Tehran, Iran

<sup>c</sup>Department of Medicinal Chemistry, Faculty of Pharmacy and Pharmaceutical Sciences Research Center Tehran university of Medical sciences, Tehran, Iran.




### Dynamic <sup>1</sup>H NMR along with theoretical study around the carbon-carbon double bond in stable phosphorus ylide derived from triphenylphosphine

pp 737-740

Sayyed Mostafa Habibi Khorassani<sup>\*</sup>, MalekTaher Maghsoodlou, Ali Ebrahimi, Marzyeh Mohammadi and Mehdi Shahraki

Department of Chemistry, University of Sistan and Baluchestan, P. O. Box 98135-674, Zahedan, Iran




**A novel synthesis of oxazino derivatives from an efficient one-pot three-component reaction of isoquinoline and dimethylacetylene dicarboxylate (DMAD) in the presence of arylaldehyde derivatives**

pp 741-744

Manzarbanou Asnaashari Isfahani\* and Fereshteh Ziaei Rad

Department of Chemistry, North Tehran Branch, Islamic Azad University, Tehran, Iran



**Interaction between an organometallic mediator and thiosulfate as an inorganic salt; A new strategy in electrocatalytic investigation**

pp 745-750


Ali Massahi<sup>a\*</sup>, Hasan Karimi-Maleh<sup>b\*</sup>, Ali Mokhtari<sup>c</sup> and Hamed Ghaedi<sup>d</sup>

<sup>a</sup> Faculty of Engineering, Bushehr Branch, Islamic Azad University, Bushehr, Iran

<sup>b</sup> Department of Chemistry, Science and Research Branch, Islamic Azad University, Mazandaran, Iran

<sup>c</sup> Department of Engineering and Technology, Golestan University, Aliabad Katool, Iran

<sup>d</sup> Faculty of Engineering, Bushehr Branch, Islamic Azad University, Bushehr, Iran




**Extraction and identification of flavonoid in *Phlomis Olivieri* benth**

pp 751-753

Mohammad Reza Zardoost<sup>a\*</sup> and Siamak Gheibi<sup>b</sup>

<sup>a</sup> Department of Chemistry, Qaemshahr Branch, Islamic Azad University, Qaemshahr, Iran

<sup>b</sup> Guilan Science & Technology Park, Entezam Sq., Rasht, Iran



## Synthesis of quinoxalines in the presence of $\text{Mg}(\text{HSO}_4)_2$ as an efficient catalyst

pp 755-759

Abdolhamid Bamoniri<sup>a</sup>, BiBi Fatemeh Mirjalili<sup>b\*</sup> and Ali Akbari<sup>b</sup>

<sup>a</sup> Department of Organic Chemistry, Faculty of Chemistry, University of Kashan, Kashan, I. R. Iran

<sup>b</sup> Department of Chemistry, College of Science, Yazd University, Yazd. P.O.Box 8915813149

