

## Extend a ranking method of trapezoidal fuzzy numbers to all fuzzy numbers by weighting functions

Abolfazl Saeidifar <sup>a,\*</sup>,

<sup>a</sup>*Faculty of Science, Arak Branch, Islamic Azad University, Arak, Iran.*

Received 12 September 2016; accepted 20 February 2017

---

### Abstract

Recently Abbasbandy and Hajjari (Computers and Mathematics with Applications 57 (2009) 413-419) have introduced a ranking method for the trapezoidal fuzzy numbers. This paper extends their method to all fuzzy numbers, which uses from a defuzzification of fuzzy numbers and a general weighting function. Extended method is interesting for ranking all fuzzy numbers, and it can be applied for solving and optimizing engineering and economics problems in a fuzzy environment.

*Key words:* Fuzzy numbers; Ranking; Weighting function.

*2010 AMS Mathematics Subject Classification :* 65L05, 34A07.

---

\* Corresponding author's E-mail: saeidifar1349@yahoo.com, a-saeidifar@iau-arak.ac.ir (A. Saeidifar)

## 1 Introduction

Ranking fuzzy numbers plays an importance role in fuzzy decision making problems; therefore, deriving the final efficiency and powerful ranking are helpful to decision makers when solving fuzzy problems. Selecting a good ranking method can apply to choosing a desired criterion in a fuzzy environment. In recent years many ranking methods have been introduced by researchers; some of these ranking methods have been compared and reviewed by Bortolan and Degani [2]. Wang and Kerre [23,24] proposed some axioms as reasonable properties to determine the rationality of a fuzzy ranking method and systematically compared a wide array of existing fuzzy ranking methods. Almost all method, however have pitfalls in some aspect, such as inconsistency with human intuition, indiscrimination, and difficulty of interpretation. What seems to be clear is that there exists no uniquely best method for comparing fuzzy numbers and different methods may satisfy different criteria. Among the existing ranking methods of fuzzy numbers, a number of them are based on area measurements with the integral value of the membership function of fuzzy numbers. A commonly used ranking technique for fuzzy numbers is the centroid based ranking method. Some other methods use statistical techniques such as simulation and hypothesis and quadratic fuzzy regression.

In the following, we first introduce the developments of centroid- based fuzzy number ranking methods. Yager [28] proposed the centroid index ranking method with a weighting function. Chen and others have proposed a centroid index ranking method that calculates the distance between the centroid point of each fuzzy number and the original point to improve some of the ranking methods [4–6,26]. They also proposed a coefficient of variation (CV index) to improve Lee and Li's method [14]. Chu and Tsao [6] proposed a ranking method of fuzzy numbers by using the area between the centroid and original point. Chen and Chen [7] proposed a ranking index based on the centroid point and standard deviations to overcome some the drawbacks of previous centroid point indices. Lee [13] proposed a fuzzy number ranking method with user viewpoints. Yager and Filve [27] proposed a ranking method with parameterized valuation functions. Detyniecki and Yager [10] proposed a fuzzy number ranking

method with an  $\alpha$  weighting function. Tran and Duckstein [21] proposed a weighting function that represents the decision maker's attitude. Lee and Li [14] introduced a ranking method for the normalized trapezoidal fuzzy numbers (NTFNs). Tang [22] showed Lee and Li's method is inconsistent. Liu and Han [25] proposed a method to rank fuzzy numbers with preference weighting function expectation. Cheng [5] has proposed the distance method for ranking fuzzy numbers. Goetschel and Voxman [12] introduced a method for ranking fuzzy numbers: their definition for ordering fuzzy numbers was motivated by the desire to give less importance to the lower levels of fuzzy numbers. Deng, Zhenfu and Qi [8] introduced the ranking of fuzzy numbers by an area method using the radius of gyration (ROG). Wang et al. [26] improved the correct centroid formula for ranking fuzzy numbers that justified them from the viewpoint of analytical geometry. Seidifar [20] has applied a weighting function and a weighted mean to ranking fuzzy numbers. Therefore, the essential subject of paper is extend a ranking method of trapezoidal fuzzy numbers to all fuzzy numbers and for any arbitrary weighting function.

The rest of this paper is organized as follows. In section 2, we recall some of the basic definitions and notions. In section 3, we introduce a extend method for ranking fuzzy numbers by the weighting mean and its properties are mentioned. The last section (Section 4) is devoted to discussion and conclusion.

## 2 Basic definitions and notions

Let  $\mathbb{R}$  be the set of all real numbers. We assume a fuzzy number  $A$  that can be expressed for all  $x \in \mathbb{R}$  in the form

$$A(x) = \begin{cases} A_L(x) , & x \in [a, b], \\ 1 & , x \in [b, c] \\ A_R(x) , & x \in [c, d], \\ 0 & , otherwise, \end{cases} \quad (\text{I})$$

where  $a, b, c$  and  $d$  are real numbers such that  $a < b \leq c < d$ ,  $A_L$  is real-valued function that is increasing and right continuous and  $A_R$  is a real-valued function that is decreasing and left continuous. Notice that (I) is an **L-R** fuzzy number with strictly monotone shape function, as proposed by Dubois and Prade in 1981. Each fuzzy number  $A$  described by (I) has the following  $\gamma$ -level sets ( $\gamma$ -cuts)  
 $[A]_\gamma = [A_L^{-1}(\alpha), A_R^{-1}(\gamma)] = [\underline{a}(\gamma), \bar{a}(\gamma)]$  for all  $\gamma \in [0, 1]$ . The family of fuzzy numbers will be denoted by  $\mathcal{F}$ .

**Definition 2.1** A fuzzy number  $A = (a, b, c, d)$  is called a trapezoidal fuzzy number if its membership function  $A(x)$  has the following form:

$$A(x) = \begin{cases} \frac{x-a}{b-a}, & x \in [a, b], \\ 1 & , x \in [b, c] \\ \frac{d-x}{d-c}, & x \in [c, d], \\ 0 & , otherwise. \end{cases} \quad (\text{II})$$

**Definition 2.2** [1] Let  $A = (a, b, c, d)$  be a trapezoidal fuzzy number with  $\gamma$ -cut  $[A]_\gamma[\underline{a}(\gamma), \bar{a}(\gamma)] = [a + (b - a)\gamma, d - (d - c)\gamma]$ . The magnitude of the trapezoidal fuzzy number  $A$  is defined as

$$Mag(A) = \frac{1}{2} \int_0^1 (\underline{a}(\gamma) + \bar{a}(\gamma) + x_0 + y_0) f(\gamma) d\gamma, \quad (2.1)$$

where  $x_0 = b, y_0 = c$  and the function  $f(\gamma)$  is a non-negative and increasing function on  $[0, 1]$  with  $f(0) = 0, f(1) = 1$  and  $\int_0^1 f(\gamma) = \frac{1}{2}$ . Function  $f(\gamma)$  is considered as a weighting function.

Abbasbandy and Hajjari[1] have applied  $Mag(A)$  for ranking of trapezoidal fuzzy numbers and a especial function  $f(\gamma) = \gamma$ . Therefore for the two trapezoidal fuzzy numbers  $A$  and  $B$  they defined

$$Mag(A) > Mag(B) \Leftrightarrow A \succ B$$

### 3 Extend ranking method of trapezoidal fuzzy numbers

In this section, we will extend the above method to all fuzzy numbers and for any weighting function.

Suppose  $f = (\underline{f}, \overline{f}) : ([0, 1], [0, 1]) \rightarrow (\mathbb{R}, \mathbb{R})$  is a weighting function such that the functions  $\underline{f}, \overline{f}$  are non-negative, monotone increasing, and satisfy the following normalization condition

$$\int_0^1 \underline{f}(\gamma) d\gamma = 1, \int_0^1 \overline{f}(\gamma) d\gamma = 1.$$

Note that if  $g = (g, \overline{g}) : ([0, 1], [0, 1]) \rightarrow (\mathbb{R}, \mathbb{R})$  is a function non-negative, and monotone increasing, then we can consider

$$\underline{f}(\gamma) = \frac{g(\gamma)}{\int_0^1 g(\gamma) d\gamma}, \overline{f}(\gamma) = \frac{\overline{g}(\gamma)}{\int_0^1 \overline{g}(\gamma) d\gamma}.$$

Let  $f = (\underline{f}, \overline{f})$  is a weighting function. Then for any arbitrary fuzzy number  $A \in \mathcal{F}$  with set  $\alpha$ -cut  $[A]_\gamma = [\underline{a}(\gamma), \overline{a}(\gamma)]$ , we define the magnitude of the fuzzy number  $A$  as

$$Mag^f(A) = \frac{\int_0^1 [(\underline{a}(\gamma) + x_{01})\underline{f}(\gamma) + (\overline{a}(\gamma) + x_{02})\overline{f}(\gamma)] d\gamma}{2(\int_0^1 \underline{f}(\gamma) d\gamma + \int_0^1 \overline{f}(\gamma) d\gamma)}, \quad (3.1)$$

where

$$x_{01} = \min\{x | A(x) \geq \max(A(x))\}, \quad x_{02} = \max\{x | A(x) \geq \max(A(x))\}.$$

The function  $f(\gamma) = (\underline{f}(\gamma), \overline{f}(\gamma))$  is considered as a weighting function. In actual applications, function  $f(\gamma)$  can be chosen according to the actual situation. In this paper we can apply the different weighting functions for ranking fuzzy numbers. Obviously, the  $Magn^f(A)$  of a fuzzy number  $A$ , synthetically reflects the information on every membership degree, and meaning of this magnitude is visual and natural. We state the properties of  $Magn^f(A)$  by the following theorems.

**Theorem 3.1** Let  $A \in \mathcal{F}$  be a fuzzy number with  $[A]_\gamma = [\underline{a}(\gamma), \bar{a}(\gamma)]$  and  $f(\gamma) = (\underline{f}(\gamma), \bar{f}(\gamma))$  be a weighted function, and  $\lambda_1, \lambda_2$  be real numbers. Then

$$\text{Mag}^f(\lambda_1 A + \lambda_2) = \lambda_1 \text{Mag}^f(A) + \lambda_2. \quad (3.2)$$

**Proof.** Suppose  $A$  is a fuzzy number with  $[A]_\gamma = [\underline{a}(\gamma), \bar{a}(\gamma)]$ . Then for the real numbers  $\lambda_1 > 0$  and  $\lambda_2$ , we have

$[\lambda_1 A + \lambda_2]_\gamma = [\lambda_1 \underline{a}(\gamma) + \lambda_2, \lambda_1 \bar{a}(\gamma) + \lambda_2]$ , and also for any  $\gamma \in [0, 1]$ , we get

$$(\lambda_1 \underline{a}(\gamma) + \lambda_2) + (\lambda_1 x_{01} + \lambda_2) = \lambda_1 (\underline{a}(\gamma) + x_{01}) + 2\lambda_2,$$

$$(\lambda_1 \bar{a}(\gamma) + \lambda_2) + (\lambda_1 x_{02} + \lambda_2) = \lambda_1 (\bar{a}(\gamma) + x_{02}) + 2\lambda_2,$$

so

$$\begin{aligned} \text{Mag}^f(\lambda_1 A + \lambda_2) &= \\ &= \frac{\int_0^1 [(\lambda_1 (\underline{a}(\gamma) + x_{01}) + 2\lambda_2) \underline{f}(\gamma) + (\lambda_1 (\bar{a}(\gamma) + x_{02}) + 2\lambda_2) \bar{f}(\gamma)] d\gamma}{2(\int_0^1 \underline{f}(\gamma) d\gamma + \int_0^1 \bar{f}(\gamma) d\gamma)} \\ &= \frac{\lambda_1 \int_0^1 [(\underline{a}(\gamma) + x_{01}) \underline{f}(\gamma) + (\bar{a}(\gamma) + x_{02}) \bar{f}(\gamma)] d\gamma + 2\lambda_2 (\int_0^1 \underline{f}(\gamma) d\gamma + \int_0^1 \bar{f}(\gamma) d\gamma)}{2(\int_0^1 \underline{f}(\gamma) d\gamma + \int_0^1 \bar{f}(\gamma) d\gamma)} \\ &= \lambda_1 \text{Mag}^f(A) + \lambda_2. \end{aligned}$$

Similarly, the theorem for  $\lambda_1 < 0$  is hold.  $\square$

**Theorem 3.2** Let  $A, B \in \mathcal{F}$  be two fuzzy numbers with  $[A]_\gamma = [\underline{a}(\gamma), \bar{a}(\gamma)]$ , and  $[B]_\gamma = [\underline{b}(\gamma), \bar{b}(\gamma)]$ . And  $f(\gamma) = (\underline{f}(\gamma), \bar{f}(\gamma))$  be a weighted function. Then

$$\text{Mag}^f(A + B) = \text{Mag}^f(A) + \text{Mag}^f(B). \quad (3.3)$$

**Proof.** Suppose that

$$x_{01}^A = \min\{x | A(x) \geq \max(A(x))\},$$

$$x_{02}^A = \max\{x | A(x) \geq \max(A(x))\},$$

$$x_{01}^B = \min\{x | B(x) \geq \max(B(x))\},$$

$$x_{02}^B = \max\{x | B(x) \geq \max(B(x))\}.$$

Then for any  $\gamma \in [0, 1]$ , we have  $[A + B]_\gamma = [\underline{a}(\gamma) + \underline{b}(\gamma), \bar{a}(\gamma) + \bar{b}(\gamma)]$ ,

and also

$$\begin{aligned}
& [\underline{a}(\gamma) + \underline{b}(\gamma), \bar{a}(\gamma) + \bar{b}(\gamma)] + [x_{01}^A + x_{01}^B, x_{02}^A + x_{02}^B] \\
&= [\underline{a}(\gamma) + x_{01}^A + \underline{b}(\gamma) + x_{01}^B, \bar{a}(\gamma) + x_{02}^A + \bar{b}(\gamma) + x_{02}^B], \\
&= [\underline{a}(\gamma) + x_{01}^A, \bar{a}(\gamma) + x_{02}^A] + [\underline{b}(\gamma) + x_{01}^B, \bar{b}(\gamma) + x_{02}^B],
\end{aligned}$$

thus with replacement in Eq.2, implies that

$$Mag^f(A + B) = Mag^f(A) + Mag^f(B).$$

□

**Theorem 3.3** *Let  $A \in \mathcal{F}$  is a symmetrical fuzzy numbers with  $[A]_\gamma = [\underline{a}(\gamma), \bar{a}(\gamma)]$ , and  $\underline{f}(\gamma) = \bar{f}(\gamma) = f(\gamma)$  be a weighted function. Then we have  $Mag^f(A) = \bar{k}$ , where  $k = \frac{x_{01} + x_{02}}{2}$ .*

**Proof.** Since fuzzy number  $A$  with  $[A]_\gamma = [\underline{a}(\gamma), \bar{a}(\gamma)]$  is a fuzzy symmetrical fuzzy number, then for any  $\gamma \in [0, 1]$  we have  $\frac{\underline{a}(\gamma) + \bar{a}(\gamma)}{2} = k$ , and so

$$\begin{aligned}
Mag^f(A) &= \frac{\int_0^1 [(\underline{a}(\gamma) + x_{01})f(\gamma) + (\bar{a}(\gamma) + x_{02})f(\gamma)]d\gamma}{2(\int_0^1 f(\gamma)d\gamma + \int_0^1 f(\gamma)d\gamma)} \\
&= \frac{\int_0^1 [(\underline{a}(\gamma) + \bar{a}(\gamma))f(\gamma) + (x_{01} + x_{02})f(\gamma)]d\gamma}{4} = \frac{2k + 2k}{4} = k.
\end{aligned}$$

**Definition 3.1** [19] *Let  $A \in \mathcal{F}$  be a fuzzy number with  $[A]_\gamma = [\underline{a}(\gamma), \bar{a}(\gamma)]$  and  $f(\gamma) = (\underline{f}(\gamma), \bar{f}(\gamma))$  be a weighting function. Then the  $f$ -weighted mean of  $A$  is defined as*

$$\bar{M}_f(A) = \int_0^1 \frac{\underline{f}(\gamma)\underline{a}(\gamma) + \bar{f}(\gamma)\bar{a}(\gamma)}{2} d\gamma. \quad (3.4)$$

Saeidifar [19] has shown that  $\bar{M}_f(A)$  is a weighting mean of fuzzy number  $A$ , and it is also the nearest weighted point approximation to fuzzy number  $A$  which is unique. □

**Theorem 3.4** *Let  $A \in \mathcal{F}$  is a symmetrical fuzzy numbers with  $[A]_\gamma = [\underline{a}(\gamma), \bar{a}(\gamma)]$ , and  $\underline{f}(\gamma) = \bar{f}(\gamma) = f(\gamma)$  be a weighted function. Then we have  $Mag^f(A) = \bar{M}_f(A) = k$ .*

**Proof.** The proof is similar to the proof of theorem 3.  $\square$

**Proposition 3.1** *Let  $A = (a, b, c, d)$  be a trapezoidal fuzzy number and let  $f(\gamma) = (\underline{f}(\gamma), \overline{f}(\gamma))$  be a weighting function, then the following hold.*

(1) For  $f(\gamma) = (1, 1)$ , we have

$$Mag^f(A) = \frac{a + 3b + 3c + d}{8}. \quad (3.5)$$

(2) For  $f(\gamma) = (2\gamma, 2\gamma)$ ,

$$Mag^f(A) = \frac{a + 5(b + c) + d}{12}, \quad (3.6)$$

(3) For  $f(\gamma) = (n\gamma^{n-1}, n\gamma^{n-1})$ ,  $n \in \mathcal{N}$  (natural numbers),

$$Mag^f(A) = \frac{a + (b + c)(2n + 1) + d}{4(n + 2)}, \quad (3.7)$$

(4) For  $f(\gamma) = (m\gamma^{m-1}, n\gamma^{n-1})$ ,  $m, n \in \mathcal{N}$ ,

$$Mag^f(A) = \frac{a + (2m + 1)b}{4(m + 1)} + \frac{d + (2n + 1)c}{4(n + 1)}. \quad (3.8)$$

**Proof.** The proof is simple.

**Proposition 3.2** *Let  $A = (a, b, c, d)$  be a trapezoidal fuzzy number and let  $f(\gamma) = (m\gamma^{m-1}, n\gamma^{n-1})$ ,  $m, n \in \mathcal{N}$  be a weighting function. Then, for  $m, n \rightarrow \infty$*

$$Mag^f(A) = \frac{b + c}{2}$$

We now apply the  $Mag^f(A)$  for ranking fuzzy numbers as the following definition.

**Definition 3.2** *For two fuzzy numbers  $A, B \in \mathcal{F}$ , and the weighting function  $f$ , we define the ranking of  $A$  and  $B$  by  $Mag^f(A)$ , i.e.*



- $Mag^f(A) < Mag^f(B)$  if and only if  $A \prec B$
- $Mag^f(A) = Mag^f(B)$  if and only if  $A \sim B$
- $Mag^f(A) > Mag^f(B)$  if and only if  $A \succ B$ .

Then we formulate the order  $\preceq$  and  $\succeq$  as

$A \preceq B$  if and only if  $A \prec B$  or  $A \sim B$

$A \succeq B$  if and only if  $A \succ B$  or  $A \sim B$ .

We consider the following reasonable properties for the ordering approaches ( see [24]).

- $A_1$  : For an arbitrary finite subset  $\Gamma$  of  $\mathcal{F}$  and  $A \in \Gamma$ ,  $A \succeq A$ .
- $A_2$  : For an arbitrary finite subset  $\Gamma$  of  $\mathcal{F}$  and  $(A, B) \in \Gamma^2$ ,  $A \succeq B$  and  $B \succeq A$ , we should have  $A \sim B$ .
- $A_3$  : For an arbitrary finite subset  $\Gamma$  of  $\mathcal{F}$  and  $(A, B, C) \in \Gamma^3$ ,  $A \succeq B$  and  $B \succeq C$ , we should have  $A \succeq C$ .
- $A_4$  : For an arbitrary finite subset  $\Gamma$  of  $\mathcal{F}$  and  $(A, B) \in \Gamma^2$ ,  $inf sup(A) \geq inf sup(B)$  we should have  $A \succeq B$ .
- $A'_4$  : For an arbitrary finite subset  $\Gamma$  of  $\mathcal{F}$  and  $(A, B) \in \Gamma^2$ ,  $inf sup(A) > inf sup(B)$  we should have  $A \succ B$ .
- $A'_5$  : Let  $\Gamma$  and  $\Gamma'$  be two arbitrary finite subset of  $\mathcal{F}$ ; also,  $A$  and  $B$  are in  $\Gamma \cap \Gamma'$ . We obtain the ranking order  $A \succ B$  by  $Mag^f(\cdot)$  on  $\Gamma'$  if and only if  $A \succ B$  by  $Mag^f(\cdot)$  on  $\Gamma$ .
- $A_6$  : Let  $A, B, A+C$  and  $B+C$  be elements of  $\mathcal{F}$ . If  $A \succeq B$ , then  $A+C \succeq B+C$ .
- $A'_6$  : Let  $A, B, A+C$  and  $B+C$  be elements of  $\mathcal{F}$ . If  $A > B$ , then  $A+C \succ B+C$ .
- $A_7$  : For an arbitrary finite subset  $\Gamma$  of  $\mathcal{F}$  and  $A \in \Gamma$ ,  $Mag^f(A)$  must belong to its support.

**Theorem 3.5** *The function  $Mag^f(\cdot)$  has the properties  $A_1, A_2, \dots, A_7$ .*

**Proof.** It is easy to verify that the properties  $A_1 - A_6$  hold. For the proof of  $A_7$  we consider the fuzzy number  $[A]_\gamma = [\underline{a}(\gamma), \bar{a}(\gamma)]$ , and the weighting function  $f(\gamma) = (\underline{f}(\gamma), \bar{f}(\gamma))$ . For all  $\gamma \in [0, 1]$ , we have  $a \leq$

$\underline{a}(\gamma) \leq \bar{a}(\gamma) \leq d$ ; hence

$$\begin{aligned} \frac{\underline{f}(\gamma)(a+a) + \bar{f}(\gamma)(a+a)}{4} &\leq \frac{\underline{f}(\gamma)(\underline{a}(\gamma) + x_{01}) + \bar{f}(\gamma)(\bar{a}(\gamma) + x_{02})}{4} \\ &\leq \frac{\underline{f}(\gamma)(d+d) + \bar{f}(\gamma)(d+d)}{4}, \end{aligned}$$

so

$$\begin{aligned} \int_0^1 \frac{\underline{f}(\gamma)2a + \bar{f}(\gamma)2a}{4} d\gamma &\leq \int_0^1 \frac{\underline{f}(\gamma)(\underline{a}(\gamma) + x_{01}) + \bar{f}(\gamma)(\bar{a}(\gamma) + x_{02})}{4} d\gamma \\ &\leq \int_0^1 \frac{\underline{f}(\gamma)2d + \bar{f}(\gamma)2d}{4} d\gamma, \end{aligned}$$

or

$$\begin{aligned} \frac{2a}{4} \int_0^1 (\underline{f}(\gamma) + \bar{f}(\gamma)) d\gamma &\leq \int_0^1 \frac{\underline{f}(\gamma)(\underline{a}(\gamma) + x_{01}) + \bar{f}(\gamma)(\bar{a}(\gamma) + x_{02})}{4} d\gamma \\ &\leq \frac{2d}{4} \int_0^1 (\underline{f}(\gamma) + \bar{f}(\gamma)) d\gamma, \end{aligned}$$

and this implies that

$$a \leq \text{Mag}^f(A) \leq d.$$

□

**Example 3.1** Let  $A, B, C$  and  $D$  be four fuzzy numbers with the following membership functions (Fig.1) and  $f(\gamma) = (f_1(\gamma), f_2(\gamma))$  be a weighting function.

$$A(x) = \begin{cases} \frac{7-x}{4} & 3 \leq x \leq 7, \\ 0 & \text{otherwise,} \end{cases} \quad B(x) = \begin{cases} 1 - \frac{(x-5)^2}{4} & 3 \leq x \leq 7, \\ 0 & \text{otherwise,} \end{cases}$$

$$C(x) = \begin{cases} \frac{x-3}{2} & 3 \leq x \leq 5, \\ \frac{7-x}{2} & 5 \leq x \leq 7, \\ 0 & \text{otherwise,} \end{cases} \quad D(x) = \begin{cases} \frac{x-3}{4} & 3 \leq x \leq 7, \\ 0 & \text{otherwise.} \end{cases}$$

Then we have

$$\begin{aligned} [A]_\gamma &= [\underline{b}(\gamma), \bar{b}(\gamma)] = [3, 7 - 4\gamma], \\ [B]_\gamma &= [\underline{a}(\gamma), \bar{a}(\gamma)] = [5 - 2\sqrt{1-\gamma}, 5 + 2\sqrt{1-\gamma}], \\ [C]_\gamma &= [\underline{c}(\gamma), \bar{c}(\gamma)] = [3 + 2\gamma, 7 - 2\gamma], \\ [D]_\gamma &= [\underline{c}(\gamma), \bar{c}(\gamma)] = [3 + 4\gamma, 7], \quad \gamma \in (0, 1]. \end{aligned}$$

For the different weighting functions the results are given in the Table 1.




Fig. 1. Fuzzy numbers  $A, B, C$  and  $D$  from Example 1.

Table 1 : Comparative results of Example 1.

Fuzzy number	A	B	C	D
Abbasbandy and Hajjari method	3.333	not defined	5	6.5
result	$A \prec C \prec D$			
Extended method, $f(\gamma) = (1, 1)$	3.5	5	5	6.5
result	$A \prec B \sim C \prec D$			
Extended method, $f(\gamma) = (2\gamma, 2\gamma)$	3.333	5	5	6.667
Result	$A \prec B \sim C \prec D$			
Extended method, $f(\gamma) = (3\gamma^2, 2\gamma)$	3.333	5.038	5.042	6.75
Result	$A \prec B \prec C \prec D$			
Extended method, $f(\gamma) = (2\gamma, 3\gamma^2)$	3.25	4.962	3.958	6.667
Result	$A \prec C \prec B \prec D$			

This example shows that extended method is flexible and can solve his shortcoming by select a suitable weighting function, even it can rank symmetrical fuzzy numbers such that previously method (methods) cannot rank these fuzzy numbers, and this is an interesting property for extended metod. Also, one can see that with change the left and right weighting functions ( $\underline{f}, \bar{f}$ ) is may that change the ordering of fuzzy numbers, specially for symmetrical fuzzy numbers(see Table 1).




Fig. 2. Fuzzy numbers  $A_{0.5}, A_1, A_2$  from Example 2 We obtain that

**Example 3.2** Consider fuzzy number  $A_n$  (Fig.2) as follows:

$$A_n(x) = \begin{cases} \left(\frac{x}{10}\right)^n & , & x \in [0, 10], \\ 1 & , & x \in (10, 11], \\ (12 - x)^n & x \in (11, 12], \\ 0 & , & \text{otherwise.} \end{cases}$$

iff  $(\gamma) = (1, 1)$  then  $Mag^f(A_{0.5}) = 9, Mag^f(A_1) = 9.375, Mag^f(A_2) = 9.75$  and hence the ranking order is  $A_{0.5} \prec A_1 \prec A_2$ . If  $f(\gamma) = (2\gamma, 2\gamma)$  then  $Mag^f(A_{0.5}) = 9.375, Mag^f(A_1) = 9.75, Mag^f(A_2) = 10.05$ , and hence  $A_{0.5} \prec A_1 \prec A_2$ . If  $f(\gamma) = (3\gamma^2, 2\gamma)$  then  $Mag^f(A_{0.5}) = 9.625, Mag^f(A_1) = 9.958$  and  $Mag^f(A_2) = 10.193$ . So  $A_{0.5} \prec A_1 \prec A_2$ .

The above examples show that this method is extended and improved of ranking method of trapezoidal fuzzy numbers [1].

Note that decision makers can select the other suitable functions for ranking fuzzy numbers. Therefore extended method of ranking fuzzy numbers is interesting and flexible.

**Example 3.3** Consider the following sets (Fig.3); see [20].

- Set 1:  $A = (0.3, 0.4, 0.7, 0.9), B = (0.3, 0.7, 0.9), C = (0.5, 0.7, 0.9)$ ,
- Set 2:  $A = (0.3, 0.5, 0.7), B = (0.3, 0.5, 0.8, 0.9), C = (0.3, 0.5, 0.9)$ ,
- Set 3:  $A = (0, 0.4, 0.7, 0.8), B = (0.2, 0.5, 0.9), C = (0.1, 0.6, 0.8)$ .

To compare with other methods we refer the reader to Table 2.

Table 2 : Comparative results of Example 3.

Authors	Fuzzy number	set 1	set 2	set 3
Choobineh and Li	A	0.458	0.333	0.50
	B	0.583	0.4167	0.5833
	C	0.667	0.5417	0.6111
Results		$A \prec B \prec C$	$A \prec B \prec C$	$A \prec B \prec C$
Yager	A	0.5778	0.5	0.4336
	B	0.6333	0.6222	0.5353
	C	0.8571	0.6986	84
Results		$A \prec B \prec C$	$A \prec B \prec C$	$A \prec B \prec C$
Chen	A	0.4315	0.375	0.52
	B	0.5625	0.425	0.57
	C	0.625	0.55	0.625
Results		$A \prec B \prec C$	$A \prec B \prec C$	$A \prec B \prec C$
Baldwin and Guild	A	0.27	0.27	0.40
	B	0.27	0.37	0.42
	C	0.37	0.45	0.42
Results		$A \sim B \prec C$	$A \prec B \prec C$	$A \prec B \sim C$
Chu and Tsao	A	0.2847	0.25	0.24402
	B	0.32478	0.31526	0.26243
	C	0.350	0.27475	0.2619
Results		$A \prec B \prec C$	$A \prec C \prec B$	$A \prec C \prec B$
Cheng Distance	A	0.7577	0.7071	0.7106
	B	0.8149	0.8037	0.7256
	C	0.8602	0.7458	0.7241
Results		$A \prec B \prec C$	$A \prec C \prec B$	$A \prec C \prec B$

<i>Wang et al. Centroid</i>	<i>A</i>	0.2568	0.1778	0.1967
	<i>B</i>	0.2111	0.2765	0.1778
	<i>C</i>	0.2333	0.1889	0.1667
<i>Results</i>		$B \prec C \prec A$	$A \prec C \prec B$	$C \prec B \prec A$
<i>Wang Distance</i>	<i>A</i>	0.7289	0.6009	0.6284
	<i>B</i>	0.7157	0.7646	0.6289
	<i>C</i>	0.7753	0.6574	0.6009
<i>Results</i>		$B \prec A \prec C$	$A \prec C \prec B$	$C \prec A \prec B$
<i>Yao and Wu</i>	<i>A</i>	0.575	0.5	0.475
	<i>B</i>	0.65	0.625	0.525
	<i>C</i>	0.7	0.55	0.525
<i>Results</i>		$A \prec B \prec C$	$A \prec C \prec B$	$A \prec B \sim C$
<i>Deng et al. area method</i>	<i>A</i>	0.3169	0.2369	0.2523
	<i>B</i>	0.3240	0.3503	0.2495
	<i>C</i>	0.3240	0.2549	0.2473
<i>Results</i>		$A \prec B \sim C$	$A \prec C \prec B$	$C \prec B \prec A$
<i>Cheng CV uniform distribution</i>	<i>A</i>	0.0328	0.0133	0.0693
	<i>B</i>	0.0246	0.0304	0.0385
	<i>C</i>	0.0095	0.0275	0.0433
<i>Results</i>		$C \prec B \prec A$	$A \prec C \prec B$	$B \prec C \prec A$
<i>Cheng CV proportional distribution</i>	<i>A</i>	0.026	0.008	0.0471
	<i>B</i>	0.0146	0.0234	0.0236
	<i>C</i>	0.0057	0.0173	0.0255
<i>Results</i>		$C \prec B \prec A$	$A \prec C \prec B$	$B \prec C \sim A$




Fig. 3. Fuzzy numbers A,B,C in sets 1, 2, 3.

Authors	Fuzzy number	set 1	set 2	set 3
Goetschel and Voxman	A	0.5667	0.5	0.5
	B	0.6667	0.6333	0.5167
	C	0.7	0.5333	0.55
Results		$A \prec B \prec C$	$A \prec C \prec B$	$A \prec B \prec C$
Abbasbandy and Asady sign distance	A	1.15	1	0.95
	B	1.3	1.25	1.05
	C	1.4	1.1	1.05
Results		$A \prec B \prec C$	$A \prec C \prec B$	$A \prec B \sim C$
Asady and Zendehnam	A	0.575	0.5	0.475
	B	0.65	0.625	0.525
	C	0.7	0.55	0.525
Results		$A \prec B \prec C$	$A \prec C \prec B$	$A \prec B \sim C$
Saeidifar $f(\gamma) = (2\gamma, 2\gamma)$	A	0.5667	0.5	0.5
	B	0.6667	0.6333	0.5167
	C	0.7	0.5333	0.55
Results		$A \prec B \prec C$	$A \prec C \prec B$	$A \prec B \prec C$

<i>Abbasbandy and Hajjari</i>	A	0.5558	0.5	0.5250
	B	0.6334	0.6416	0.5084
	C	0.7	0.5166	0.5750
<i>Results</i>		$A \prec B \prec C$	$A \prec C \prec B$	$B \prec A \prec C$
<i>Extended method</i>	A	0.5604	0.5042	0.5333
	B	0.6917	0.6458	0.5146
	C	0.7042	0.5208	0.5854
<i>Results</i>		$A \prec B \prec C$	$A \prec C \prec B$	$B \prec A \prec C$

We surmise the ranking results that is agreement with the extended method as the following Table based on the percentage.

Table 3.

<i>Sets</i>	<i>Set 1</i>	<i>Set 2</i>	<i>Set 3</i>
<i>Extended method result</i>	$A \prec B \prec C$	$A \prec C \prec B$	$B \prec A \prec C$
<i>Agreement percentage</i>	67	78	11

Note that in set 3, the results of ranking fuzzy numbers are very different.

**Example 3.4** Let  $A = (0.1, 0.3, 0.3, 0.5)$ ,  $B = (0.2, 0.3, 0.3, 0.4)$ , and  $C = (1, 1, 1, 1)$ , and  $f = (2\gamma, 3\gamma^2)$ . Then,  $\text{Mag}^f(A) = 0.2958$ ,  $\text{Mag}^f(B) = 0.2979$ ,  $\text{Mag}^f(C) = 1$ . Therefore the ranking order is  $A \prec B \prec C$ .

**Example 3.5** Let  $A = (0.1, 0.3, 0.3, 0.5)$ ,  $B = (-0.5, -0.3, -0.3, -0.1)$ , and  $f = (2\gamma, 2\gamma)$ . Then,  $\text{Mag}^f(A) = 0.3$ ,  $\text{Mag}^f(B) = -0.3$ . Therefore,  $A \succ B$

The above examples results are the same as the method result of Phani Bushan Rao et al. [18].

**Example 3.6** [18] Consider four fuzzy numbers  $A_1 = (0.1, 0.2, 0.3)$ ,  $A_2 = (0.2, 0.5, 0.8)$ ,  $A_3 = (0.3, 0.4, 0.9)$  and  $A_4 = (0.6, 0.7, 0.8)$  which were ranked earlier by Yager [28], Fortemps and Roubens [11], Liou and Wang [16], and Chen and Lu [3] as shown in Table 4. It can be seen from Table 4 that none of the methods discriminates fuzzy numbers. Yager [28] and Fortemps and Roubens [11] methods failed to discriminate the fuzzy num-



bers  $A_2$  and  $A_3$ , whereas the methods of Liou and Wang [16] and Chen and Lu [3] cannot discriminate the fuzzy numbers  $A_2, A_3$  and  $A_1, A_4$ .

By Extended method, and weighting functions  $f_1 = (2\gamma, 2\gamma)$ , and  $f_2 = (2\gamma, 3\gamma^2)$ , we get

$Mag^{f_1}(A) = 0.2$ ,  $Mag^{f_1}(B) = 0.5$ ,  $Mag^{f_1}(C) = 0.4333$ ,  $Mag^{f_1}(D) = 0.7$

$Mag^{f_2}(A) = 0.1979$ ,  $Mag^{f_2}(B) = 0.4937$ ,  $Mag^{f_2}(C) = 0.4229$ ,  $Mag^{f_2}(D) = 0.6979$ .

Therefore the ranking order is  $A_4 \succ A_2 \succ A_3 \succ A_1$ . Observe that the our method result is the same as method of Phani Bushan Rao1 et al. [18].

Table 4: Comparison of various ranking methods.

Method		$A_1$	$A_2$	$A_3$	$A_4$	Ranking order
Ranking order Yager		0.20	0.50	0.50	0.70	$A_4 \succ A_2 \sim A_3 \succ A_1$
Fortemps and Roubens		0.20	0.50	0.50	0.70	$A_4 \succ A_2 \sim A_3 \succ A_1$
Liou and Wang	$\alpha = 1$	0.25	0.65	0.65	0.75	$A_4 \succ A_2 \sim A_3 \succ A_1$
	$\alpha = 0.5$	0.20	0.50	0.50	0.70	$A_4 \succ A_2 \sim A_3 \succ A_1$
	$\alpha = 0$	0.15	0.35	0.35	0.65	$A_4 \succ A_2 \sim A_3 \succ A_1$
Chen and Lu [37]	$\beta = 1$	-0.20	0.00	0.00	-0.20	$A_2 \sim A_3 \succ A_1 \sim A_4$
	$\beta = 0.5$	-0.20	0.00	0.00	-0.20	$A_2 \sim A_3 \succ A_1 \sim A_4$
	$\beta = 0$	-0.20	0.00	0.00	-0.20	$A_2 \sim A_3 \succ A_1 \sim A_4$
Phani Bushan Rao1 et al.		0.4591	0.6320	0.6146	0.8129	$A_4 \succ A_2 \succ A_3 \succ A_1$
Extended method	$f_1 = (2\gamma, 2\gamma)$	0.2	0.5	0.4333	0.7	$A_4 \succ A_2 \succ A_3 \succ A_1$
Extended method	$f_2 = (2\gamma, 3\gamma^2)$	0.1979	0.4937	0.4229	0.6979	$A_4 \succ A_2 \succ A_3 \succ A_1$

## 4 Conclusion

In this paper, we develops a ranking method of trapezoidal fuzzy numbers to all fuzzy numbers which uses from a weighting function and a defuzzification for ranking fuzzy numbers. The properties of defuzzification ( $Mag^f(.)$ ) are given by theorems and propositions. The flexibility is one of the most important properties of extended ranking method, because decision makers can select the different weighting functions as  $f = (\underline{f}, \bar{f})$  such that functions  $\underline{f}, \bar{f} : [0, 1] \rightarrow R$  are weighting functions for the lower and upper  $\gamma$ -cuts sets of a fuzzy number, respectively. This

means that the functions  $\underline{f}(\gamma)$  and  $\overline{f}(\gamma)$  can be treated as the subjective weights indicating neutral, optimistic, or pessimistic preferences of the decision maker. Therefore, our method is more general and interesting for ranking fuzzy numbers. Also, the maximum entropy of weighting function ( $f(\gamma)$ ) is discussed in [15]. It can be used to choose a suitable weighting function.

## References

- [1] S. Abbasbandy and T. Hajjari, A new approach for ranking of trapezoidal fuzzy numbers, *Computers and Mathematics with Applications* 57 (2009) 413-419.
- [2] G. Bortolan and R. Degani, A review of some methods for ranking fuzzy numbers, *Fuzzy Sets and Systems* 15 (1985) 1-19.
- [3] L.-H. Chen and H.-W. Lu, An approximate approach for ranking fuzzy numbers based on left and right dominance, *Computers and Mathematics with Applications*, vol. 41, no. 12, pp. 1589-1602, 2001.
- [4] S. Chen, Ranking fuzzy numbers with maximizing set and minimizing set, *Fuzzy Sets and Systems* 17 (1985) 113-129.
- [5] C.H. Cheng, A new approach for ranking fuzzy numbers by distance method, *Fuzzy Sets and Systems* 95 (1998) 307-317.
- [6] T. Chu and C. Tsao, Ranking fuzzy numbers with an area between the centroid point and original point, *Computers and Mathematics with Applications*. 43 (2002) 11-117.
- [7] S.J. Chen and S.M. Chen, A new method for handling multicriteria fuzzy decision-making problems using FN-IOWA operators, *Cybernetics and Systems* 34 (2003) 109-137.
- [8] Y. Deng, Z. Zhenfu, L. Qi, Ranking fuzzy numbers with an area method using radius of gyration, *Computers and Mathematics with Applications* 51 (2006) 1127-1136.
- [9] D. Dubois and H. Prade, The mean value of a fuzzy number, *Fuzzy Sets and Systems*. 24 (1987) 279-300.

- [10] M. Detynieck and R.R. Yager, ranking fuzzy numbers using  $\alpha$ - weighted, International journal of uncertainty Fuzziness and Knowledg-Based Systems 8 (2000) 573-591.
- [11] P. Fortemps and M. Roubens, Ranking and defuzzification methods based on area compensation, Fuzzy Sets and Systems, vol. 82, no. 3, pp. 319330, 1996.
- [12] R. Goetschel and W. Voxman, Elementary calculus, Fuzzy Sets and Systems 18 (1986) 31-43.
- [13] H. Lee,k.wang and J.-H. Lee, A method rankinr for fuzzy numbers and its application to dicision-making, IEEE Transactions on Fuzzy Systems, 7 (1999) 677-685.
- [14] E.S. Lee, R.J. Li, Comparation of fuzzy numbers based on the probability measure of fuzzy event, Computers and Mathematics with Applications 15(1988) 887-896.
- [15] X. Liu, On the maximum entropy parameterizd interval approximation of fuzzy numbers, Fuzzy sets and Systems 157 (2006) 869-878.
- [16] T.-S. Liou and M.-J. Wang, Ranking fuzzy numbers with integral value, Fuzzy Sets and Systems, vol. 50, no. 3, pp. 247 255, 1992.
- [17] Ming Ma, A. Kandel, M. Friedman, A new approach for defuzzification, Fuzzy sets and Systems 111 (2000) 351-356.
- [18] P. Phani Bushan Rao1 and N. Ravi Shankar, Ranking fuzzy Numbers with a distanceMethod using circumcenter of centroids and an index of modality, Advances in Fuzzy Systems Volume 2011, Article ID 178308, 7 pages.
- [19] A. Saeidifar and E. Pasha, The possibilistic moments of fuzzy numbers and their applications, Journal of Computational Applied Mathematics, 223 (2009) 1028-1042.
- [20] A. Saeidifar, Application of weighting functions to the ranking of fuzzy numbers, Computers and Mathematics with Applications 62 (2011) 2246-2258.
- [21] L. Tran, L. Duckstein, Comparison of fuzzy numbers using a fuzzy distance measure, Fuzzy Sets and Systems 130 (2002) 331-341.

- [22] H.C. Tang, Inconsistent Property of Lee and Li fuzzy ranking method, Computers and Mathematics with Applications 45 (2003) 709-713.
- [23] X. Wang and E.E. Kerre, Reasonable properties for the ordering of fuzzy quantities (I), Fuzzy Sets and Systems 118 (2001) 375-385.
- [24] X. Wang and E.E. Kerre, Reasonable properties for the ordering of fuzzy quantities (II), Fuzzy Sets and Systems 118 (2001) 387-405.
- [25] X. Wang Liu and S. Lina Han, Ranking fuzzy numbers with preference weighting function expectations, Computers and Mathematics with Applications 49 (2005) 1731-1753.
- [26] Y. M. Wang, J. B. Yang, D. L. Xu and K. S. chin, On the centroids of fuzzy numbers, Fuzzy Sets and Systems 157 (2006) 919-926.
- [27] R.R. Yager and D.P. Filev, Parameterized and like and or-like OWA operators, International journal of General Systems,22 (1994) 297-316.
- [28] R.R. Yager, A procedure for ordering fuzzy subsets of the unit interval, Information Sciences 24 (1981) 143-161.