

## Gibberellic Acid Application and Wheat Production

MOHAMMAD HASHEM AZIZI<sup>1\*</sup>, ALI SOLEYMANI<sup>1,2\*</sup>, AND HAMID REZA JAVANMARD<sup>1</sup>

<sup>1</sup>Department of Agronomy and Plant Breeding, Isfahan (Khorasgan) Branch, Islamic Azad University, Isfahan, Iran; <sup>2</sup>Plant Improvement and Seed Production Research Center, Isfahan (Khorasgan) Branch, Islamic Azad University, Isfahan, Iran

\* Corresponding Author E-mail: a.soleymani444@gmail.com

Received: 25 October 2023

Accepted: 10 January 2024

### ABSTRACT

Gibberellic acid (GA<sub>3</sub>) is known as important plant growth regulators which are also considered to induce various physiological responses in plants, which is unduly suited for improving and stimulating, photosynthetic activity, and plant growth. Gibberellic acid, a tetracyclic di-terpenoid compound which is known as a plant hormone stimulating plant growth and development. Gibberellins may have an important function in different metabolic pathways influencing several characteristics, such as nitrogen redistribution, nitrogen metabolism, translocation of assimilates, chlorophyll production and degradation. This review article aims to provide an overview of the effects of gibberellic acid on yield and yield components of wheat and survey of different mechanisms of action of gibberellic acid in different plants. The keywords searches for gibberellic acid, wheat, root growth, gibberellin, plant hormone, and plant growth were performed by using PubMed, Google scholar, Scopus, and Web of Science.

**Keywords:** Benzyl adenine, Gibberellic acid, Grain protein, Irrigation, Salicylic acid, Superoxide dismutase, Wheat.

### INTRODUCTION

It has been reported that gibberellic acid (GA<sub>3</sub>) has significant ability to reduce the adverse impacts of salinity by promoting accumulation of osmolytes, antioxidant enzyme activity, and boosting vigor (Shahrajabian *et al.*, 2011; Seo *et al.*, 2019). GA<sub>3</sub> can significantly promote the

stimulation of various non-enzymatic and enzymatic antioxidants, and the accumulation of osmolytes in plants (Jaleel *et al.*, 2007; Misratia *et al.*, 2015; Shahrajabian *et al.*, 2021). The most important functions of GAs in plants are breaking of seed dormancy, seed germination and seedling growth, promotion of cell division and stem elongation, induction of fruit enlargement, regulation of sex expression in some plants, promotion of flowering in long day plants, prevention of genetic dwarfism, and tolerance to chilling (Gupta and Chakrabarty, 2013; Shahrajabian *et al.*, 2023). Its application can boost the development of plant by revealing the fact that they improve the amino acid concentration in embryo and promote the syntheses of hydrolytic enzyme needed for digestion of endospermic starch when renew growth at seed germination (Soleymani *et al.*, 2010; Soleymani *et al.*, 2011; Ghanbary *et al.*, 2020; Sun and Shahrajabian, 2023; Sun *et al.*, 2023). GA signaling positively regulates the resistance of rice to saline-alkali, and promotes resistance to saline-alkali stress by increasing the uptake of ammonium in rice (Li *et al.*, 2024). Bekaardt *et al.* (2004) reported that application of GA<sub>3</sub> can result in the highest seed germination in guayule, and Araujo *et al.* (2009) also showed that GA<sub>3</sub> application can increase the emergence of seeds of Jua tree. Its application can reduce the negative impacts of phytotoxic impacts, as it can restored the mobilization of starch and protein reserves from the endosperm to seedling roots during germination of barley plants (Amri *et al.*, 2016; Pan *et al.*, 2017). GA<sub>3</sub> can increase plant height, enhance yield, increase in chlorophyll contents, boost antioxidant enzymes, enhance P, K, Fe, and Mn in plants, reduce reactive oxygen species (ROS) production, and decrease in Zn concentrations (Araujo *et al.*, 2009; Rodrigues *et al.*, 2012; Liu *et al.*, 2013; Uslu *et al.*, 2013; Gokdere and Ates, 2014; Goel *et al.*, 2023). It has been reported that the foliar application of GA can significantly influence both wheat physiology and morphology, and foliar application of GA increased wheat growth and nutritional quality with positive impact on wheat irrespective of nanoparticles (NPs) combinations or alone supply (Al-Huqail *et al.*, 2023). In many experiments, it has been reported that GA<sub>3</sub> treatment can increase stem elongation through changing the orientation of cellulose microfibrils, regulate cambial cell division, promote xylem differentiation, and lead to formation of secondary xylem fibers (Sauter and Kende, 1993; Bjorklund *et al.*, 2007; Hamayun *et al.*, 2010; Yazdpour *et al.*, 2012; Cuiying *et al.*, 2013; Ogbaji *et al.*, 2013; Wang *et al.*, 2015). GA<sub>3</sub> application can delay ripening stage and increase shelf life of sweet cherry (Kondo and Danjo, 2001), and it can also improve and stimulate both bolting and flowering stage (Ingram *et al.*, 1986; Singh and Usha, 2003; Hara *et al.*, 2012; Peng *et al.*, 2014; Zhang *et al.*, 2018; Jung *et al.* 2020). This article aimed to assess the impacts of different plant growth regulators and irrigation treatments on different growth and development of winter wheat. The goal of this article is to study the impacts of gibberellic acid on different plants, especially wheat by considering different mechanism of action. In this manuscript, the presented information gathered from randomized control experiments, review articles, and analytical

studies and observations which were obtained from PubMed, Science Direct, Scopus, and Google Scholar. The keywords used were the common and Latin names of different agronomical plants, wheat, gibberellic acid, gibberellin, reactive oxygen species, plant hormone, and plant growth regulator.




Figure 1. Effects of gibberellic acid on growth, photosynthesis and antioxidant defence of crops.

### ***Gibberellic acid ( $GA_3$ ) and the Mechanism of Action***

$GA_3$  is a biologically form of Gibberellins (GAs) (O'Neill et al., 2000; Zhou et al., 2009), which can promote cell elongation, and increase cell numbers, and the are the main parameter involved in stem elongation (Srivastava and Handa, 2005; Kato *et al.*, 2011; Shen *et al.*, 2020).  $GA_3$  is important in reducing abiotic stress-induced perturbations in different plants by modulating various physio-biochemical and molecular processes, and around 140 different GAs can be discovered in higher plants and gibberellin-producing fungi, and most of them serve as precursors or catabolites, with only a few having independent biological characteristics (El-Tohamy *et al.*, 2023). GA are essential plant regulators for different plant development processes, such as seed germination, leaf extension, stem elongation, flowering induction, and pollen maturation, and mutant plants which are deficient in GA have a dwarf phenotype and late flowering, treating the plants with GA restores normal growth (Cornea-Cipcigan *et al.*, 2020; Shahzad *et al.*, 2021; Elahi *et al.*, 2023).  $GA_3$  content in the stems is the key parameter for the change in internode diameter and length of soybean internodes in response to shading (Zhang *et al.*, 2011; Hedden and Thomas, 2012; Wu *et al.*, 2017; Sun *et al.*, 2021a,b,c).  $GA_3$  offers an ideal molecule platform for chemical modifications, as it is a tetracyclic diterpenoid compound including two hydroxyl groups, a lactone ring, and a carboxyl group (Phuoc *et al.*,

2008), and GA<sub>3</sub> derivatives featuring diverse structures could show various biological roles from parent molecule GA<sub>3</sub> (Mander, 2003; Murase *et al.*, 2008), for instance, 7-homo-GA<sub>3</sub> and 16,17-dihydro-GA<sub>5</sub> inhibited stem elongation, while 16,17-dichloromethano-dihydro-GA<sub>3</sub> has an inhibitory impact on both stem elongation and flowering of many plants (Berger *et al.*, 1982; Mander *et al.*, 1998). Moreover, some gibberellin derivatives characterized by an  $\alpha,\beta$ -unsaturated ketone moiety even revealed anti-tumor characteristics (Chen *et al.*, 2009; Zhang *et al.*, 2012). The functional groups in the ring A of GA<sub>3</sub> were quite sensitive under many reaction conditions (Kirkwood *et al.*, 1980; Oh *et al.*, 2012).

The increase in GA<sub>3</sub> and GA<sub>7</sub> levels caused by the interaction of high temperature and low light which can promote the elongation of soybean hypocotyls (Bawa *et al.*, 2020; Zhang *et al.*, 2020). It can also increase the length of internodes and the overall height of plants and decreases mechanical strength (Beal *et al.*, 1996; Yuan and Yang, 2007; Martinez-Garcia *et al.*, 2010). After application of GA<sub>3</sub> to soybean the contents of jasmonic acid, GA<sub>1</sub>, and GA<sub>4</sub> were shown to have increased, while the abscisic acid and salicylic acid contents decreased (Ingram *et al.*, 1986; Martin *et al.*, 1996; Hamayun *et al.*, 2010; Larson *et al.*, 2023). GA<sub>3</sub> usage is an effectual way to change plant height, and the current researches in different parts of the world are about the usage of GA<sub>3</sub> involve the spraying of leaves or soaking of seeds (Shan *et al.*, 2021; Shahrajabian and Sun, 2023a,b,c). Zalnierius *et al.* (2022) reported that the exogenous GA<sub>3</sub> application result in incomplete seed development, and a consequent reduce in variability and germination, and the maximum GA<sub>3</sub> dose resulted in significantly reduction in propagation of Sosnowsky' hogweed through seeds in the field, which shows the importance of GA<sub>3</sub> as a promising approach to the control of the spread of some weed species. GA<sub>3</sub> application can increase and its concentration in stems is the major parameter for the change in internode diameter and length of soybean (Zhang *et al.*, 2011; Zhang *et al.*, 2020; Bawa *et al.*, 2020). The usage of GA<sub>3</sub> is effective in increasing seedling growth and length (Shahrajabian *et al.*, 2021; Sharajabian and Sun, 2023g,h,i; Sun and Shahrajabian, 2024; Sun *et al.*, 2024). It has been reported that the treatment of dwarf varieties of rye with GAs increased the activity of  $\alpha$ -amylase in the germinating caryopses (Zwar and Chandler, 1995), and treating seeds with GA<sub>3</sub> increased  $\alpha$ -amylase expression through binding of a transcription factor to the gene, which activated it (Wang *et al.*, 2016). Dwarf rye varieties are capable to synthesize GA<sub>3</sub> under certain conditions, and the normal growth of these varieties can be restored by exposure to exogenous GA<sub>3</sub> (Pavlista *et al.*, 2014; Shahrajabian and Sun, 2023d,e,f). Skalicky *et al.* (2020) reported that usage of GA<sub>3</sub> to the seeds had no impact on the transition of the growing point to the double-ridge generative stage of wheat plants, and the priming effects of GA<sub>3</sub> on seeds of common wheat varieties with various vernalization and photoperiod requirements as it influenced the transition from vegetative to generative stage.

It is also reported that GA<sub>3</sub> can help in the stimulation of various non-enzymatic and enzymatic antioxidants and the accumulation of osmolytes in plants, and the usage of exogenous GA<sub>3</sub> proliferates plant development by leading to the intensification of amino acid synthesis of hydrolytic enzymes as prerequisite for the breakdown of endospermic starch when plant seeds renovate development at the seed propagation stage (Jaleel *et al.*, 2007; Tuna *et al.*, 2008; Moumita *et al.*, 2019). The modified GA<sub>3</sub> derivatives characterized by amide groups could provide wonderful bioactivities to the target components, and the synthesis of a series of novel C3-OH substituted GA<sub>3</sub> derivatives bearing an amide moiety (Zhu *et al.*, 2012; Zong *et al.*, 2014; Gao *et al.*, 2016). It has been reported that gibberellic acid leads to cell enlargement and division, produces new tissues, increases protein biosynthesis, and promotes the absorption of nutrients and water, and it is an effective method in improving grape cultivar berries, which could be also used in wide range orchards (Abu-Zahra and Salameh, 2012; Dimovska *et al.*, 2014; Abu-Zinada, 2015; Alshallash *et al.*, 2023). For many species with physiological dormancy, it is usually possible to preemptively break seed dormancy by adding GA to the seed, which increases the endogenous levels of GA relative to ABA (Xu *et al.*, 2017; Larson *et al.*, 2023), and the addition of GA to the seed may also promote the cell expansion, activation of enzymes, and the mobilization of stored reserves, which collectively facilitate the transition from dormancy to germination (Gupta and Chakrabarty, 2013; Gonzalez-Lopez *et al.*, 2014; Shu *et al.*, 2018). Mimi *et al.* (2023) reported that gibberellins act to overcome dormancy and increase the germination rate of seeds of several species, including genus *Annona*, and GA<sub>3</sub> concentrations (500 and 1000 mg L<sup>-1</sup>) led to increases of 25%, and 20%, respectively, in the germination rate, intensification of lipid degradation in seeds with primary root emission, and a decrease in sugar concentration until the 5th days. In another experiment the application of 10<sup>-4</sup> M GA<sub>3</sub> exceeded the acceptable threshold for use in hydroponics production systems of lettuce, and GA<sub>3</sub> stimulated plant growth and increased the yield, such as leaf expansion, stomatal conductance, biomass accumulation, nitrogen use efficiency, and water use efficiency (Miceli *et al.*, 2019). Foliar application of gibberellic acid negatively influenced vegetative growth of kenaf, and exogenous gibberellic acid delayed flower and bud initiation in kenaf cultivars (Muniandi *et al.*, 2018). Gibberellins biosynthesis pathway is presented in Figure 2.




Figure 2. Gibberellins biosynthesis pathway; residing in three different cellular compartments (cytoplasm, endoplasmic reticulum, and plastid). *ent*-CDP: *ent*-copalyl diphosphate; GGDP: geranylgeranyl diphosphate; KS: *ent*-kaurene synthase; CPS: *ent*-copalyl diphosphate synthase; KAO: *ent*-kaurenoic acid oxidase, KO: *ent*-kaurene oxidase.

### ***Gibberellic acid (GA<sub>3</sub>) and Wheat Production***

Gibberellic acid (GA<sub>3</sub>) are known as important plant growth regulators which are also considered to induce various physiological responses in plants (Azizi *et al.*, 2023), which is unduly suited for improving and stimulating, photosynthetic activity, and plant growth (Shayanfar *et al.*, 2011; Soleymani *et al.*, 2013; Soleymani and Shahrajabia, 2017). GA<sub>3</sub> is a biologically active form of gibberellins (GAs) (Srivastava and Handa, 2005), and GAs promote cell numbers, and increase cell elongation, which has shown in importance in stem elongation (Hedden and Thomas, 2012; Shen *et al.*, 2020). GAs can stimulate root and stem elongation, flowering, leaf expansion, seed germination, fruit senescence, and seed dormancy (Hooley, 1994; Kato *et al.*, 2011; Hedden and Sponsel, 2015; Cui *et al.*, 2023), as they can lead to better cell division, and cell elongation during growth, and improve the expression of hydrolytic enzymes which have function in the conversion of starch into sugar (Akiyama and Suzuki, 1980; Broumand *et al.*, 2010; Khoshkharam *et al.*, 2010; Riaziat *et al.*, 2012; Sayed *et al.*, 2020; Aliahmadi *et al.*, 2021). The application of gibberellic acid can cause a rapid stimulation of growth of wheat plants, reaching a maximum withing four weeks, and it can also cause yellowing of the herbage and stem elongation up to the first mowing, six weeks after application (Scott, 1959; Horre *et al.*, 1989; Hartung *et al.*, 2010; Matzer *et al.*, 2010; Rebetzke *et al.*, 2012; Gupta and Chakrabarty, 2013).

Pfleger *et al.* (2011) reported that gibberellic acid 30x had significant influence on wheat seedling development, and the wheat growth bio assay can be appropriate tool for further

experiments on homeopathic dilutions of gibberellic acid. The positive increase of grains of winter wheat was also reported after application of gibberellic acid ( $10^{-30}$ ) (Endler *et al.*, 2015). Priming with GA<sub>3</sub> ( $150 \text{ mgL}^{-1}$ ) was very effective in increasing salicylic acid concentration in wheat cultivars under salt stress, and its treatment decreased leaf free putrescine and spermidine concentrations in the plants of wheat cultivars (Iqbal and Ashraf, 2013). Foliar application of GA increased wheat growth and nutritional quality when exposed to nanoparticles (NPs), minimized the oxidative stress, enhanced the growth of NPs-stressed wheat, and showed positive impacts on wheat irrespective of NPs combinations or alone supply (Hou *et al.*, 2008; Al-Huail *et al.*, 2023). GA increased the wheat growth at all doses of TiO<sub>2</sub> NPs, increased the photosynthetic pigments and antioxidant enzyme activities at all doses of TiO<sub>2</sub> NPs, and boosted the nutritional values of wheat grains under TiO<sub>2</sub> NPs (Alharby *et al.*, 2021; Alybayev *et al.*, 2024). Exogenous application of GA increased seed germination, and plant growth traits under Ni stress, it could lower the Ni fixation inside the root cell wall and hemicellulose, and regulated the antioxidant activities, glyoxalase systems to scavenge the ROS, and modulated the phytochelatin and antioxidant-related gene expression level for sequestration and efflux (Bossen *et al.*, 1991; Mathur *et al.*, 1992; Bhat *et al.*, 2023). GAs are synthesized by the action of terpene cyclases, cytochrome P450 mono-oxygenases and 2-oxoglutarate-dependent dioxygenases localized, respectively, in plastids, cytosol, and the endomembrane system, and GAs are synthesized in actively growing organs on plants like roots, leaves, fruits, and developing seeds (Akiyama and Suzuki, 1981; Saluja *et al.*, 1987; Saluja *et al.*, 1989). Chattopadhyay *et al.* (1994) showed that GA<sub>3</sub> induced elongation of wheat coleoptile sections and the effect of GA was reduced to 1/4 when the endogenous methyleneoxindole (MeOx), an oxidation product of IAA, was made unavailable to GA<sub>3</sub> during its action.

## CONCLUSION

Plant growth regulators are synthetic or natural substances which influence the growth and development of many crops, especially wheat. They control one or more specific biochemical and physiological characteristics possibly by their effects on enzyme and gene activities. In fact, the organic chemical component other than nutrients which are active in low concentration in inhibiting, promoting or modifying growth and development should be called growth regulators. Plant growth regulator which comes through photosynthesis supplies and respiration can control the growth and development of plant. Phytohormones are organic components produced naturally by the plants in low concentration with very low concentration. GAs are the second noticeable growth hormones found in plants, and they are natural or endogenous hormones which can be produced by some tissues in plants. The physiological processes like growth and development of the plant, flower differentiation, enhancement of the fruit color,

tissue growth, fruit ripening, etc., are controlled by the suitable applications of GAs. GAs are synthesized in actively growing organs by the action of terpene cyclases, cytochrome P450 mono-oxygenases and 2-oxoglutarate-dependent dioxygenases localized, respectively in the cytosol, the endomembrane, and in plastids. In wheat production, exogenous application of GA<sub>3</sub> can promote seed germination and subsequent seedling growth under different stresses. Its application can ensure the growth of wheat plants.

### **Disclosure statement**

The authors declare that they do not have any conflict of interest

### **Author contributions**

All authors have contributed equally to this manuscript.

### **Funding**

This research was part of Ph-D thesis of the first author, and some part of the research was supported by Department of Agronomy and Plant Breeding, Faculty of Agriculture, Islamic Azad University of Isfahan (Khorasgan), Isfahan, Iran.

### **REFERENCES**

- Abu-Zahra TR, Salameh M. 2012. N influence of gibberellic acid and cane girdling on berry size of black magic grape cultivar. *Middle-East Journal of Scientific Research*, 11:718-722.
- Abu-Zinada IA. 2015. Effect of GA<sub>3</sub>, girdling or pruning on yield and quality of Parletta seedless grape. *American Journal of Agriculture and Forestry*, 3:230.
- Akiyama T, Suzuki H. 1980. The release of acid phosphatase in wheat half-seed treated with gibberellic acid. *Biochemie und Physiologie der Pflanzen*, 175(5):436-446.
- Akiyama T, Suzuki H. 1981. Inhibitory effects of abscisic acid on gibberellic acid-induced acid phosphatase activity in wheat half-seeds. *Plant Science Letters*, 21(2):137-143.
- Alharby HF, Rizwan M, Iftikhar A, Hussaini KM, Rehman MZU, Bamagoos AA, Alharbi BM, Asrar M, Yasmeen T, Ali S. 2021. Effect of gibberellic acid and titanium dioxide nanoparticles on growth, antioxidant defense system and mineral nutrient uptake in wheat. *Ecotoxicology and Environmental Safety*, 221:112436.
- Al-Huqial AA, Alshehri D, Nawaz R, Irshad MA, Iftikhar A, Hussaini KM, Rizwan M, Alghanem SMS, Abeer AHA. 2023. The effect of gibberellic acid on wheat growth and nutrient uptake under combined stress of cerium, zinc, and titanium dioxide nanoparticles. *Chemosphere*, 336:139199.
- Aliahmadi Z, Mohadesi A, Ranjbar M, Javanshah A. 2021. Preparation and evaluation of Ca/Mg-layered double hydroxide as a novel modifier for electrochemical determination of gibberellic acid. *Journal of Molecular Structure*, 1246:131200.



- Alshallash KS, Fahmy MA, Tafeeq AM, Baghdady GA, Abdrabboh GA, Hamdy AE, Kabsha EA. 2023. GA<sub>3</sub> and hand thinning improves physical, chemical characteristics, yield and decrease bunch compactness of Sultanina grapevines (*Vitis vinifera* L.). Horticulturae, 9(2):160.
- Alybayev S, Smekenov I, Kuanbay A, Sarbassov D, Bissenbaev A. 2024. Gibberellic-acid-dependent expression of α-amylase in wheat aleurone cells is mediated by target of rapamycin (TOR) signaling. Current Plant Biology, 37:100312.
- Amri B, Khamassi K, Ali MB, Silva JATD, Kaab LBB. 2016. Effects of gibberellic acid on the process of organic reserve mobilization in barley grains germinated in the presence of cadmium and molybdenum. South African Journal of Botany, 106:35-40.
- Araujo PC, Freitas RMOD, Praxedes SC, Ribeiro MCC. 2009. Gibberellic acid enhances the emergence of *Zizyphus joazeiro* seeds. Journal of New Seeds. 10(4):277-281.
- Azizi MH, Soleymani A, Javanmard HR. 2023. Wheat (*Triticum aestivum* L.) biochemical and nutritional properties affected by plant growth regulators under field conditions. Cereal Research Communications. <https://doi.org/10.1007/s42976-023-0041401>
- Bawa G, Feng L, Chen G, Chen H, Hu Y, Pu T, Cheng YJ, Shi Y, Xiao T, Zhou WG, Yong T, Sun X, Yang F, Yang W, Wang X. 2020. Gibberellins and auxin regulate soybean hypocotyl elongation under low light and high-temperature interaction. Physiologia Plantarum, 170:345-356.
- Beal FD, Yeung EC, Pharis RP. 1996. Far-red light stimulates internode elongation, cell division, cell elongation and gibberellin levels in bean. Canadian Journal of Botany, 74:743-752.
- Bekaardt CR, Pieterse PJ, Coetzee JH, Agenbag GA. 2004. The breaking of seed dormancy of guayule (*Parthenium argentatum* Gray). By treatment with gibberellic acid and smoke water solution. South of African Journal of Plant and Soil, 21(2):114-116.
- Bergner C, Lischewski M, Adam G, Sembdner G. 1982. Biological activity of gibberellin analogues. Planta, 155:231-237.
- Bhat JA, Basit F, Alyemeni MN, Mansoor S, Kaya C, Ahmad P. 2023. Gibberellic acid mitigates nickel stress in soybean by cell wall fixation and regulating oxidative stress metabolism and glyoxalase system. Plant Physiology and Biochemistry, 198:107678.
- Bjorklund S, Antti H, Uddestrand I, Mortiz T, Sundberg B. 2007. Cross-talk between gibberellin and auxin in development of *Populus* wood: Gibberellin stimulates polar auxin transport and has a common transcriptome with auxin. The Plant Journal, 52:499-511.
- Bossen ME, Tretyn A, Kendrick RE, Vredenberg WJ. 1991. Comparison between swelling of etiolated wheat (*Triticum aestivum* L.) protoplasts induced by phytochrome and α-naphthaleneacetic acid, benzylaminopurine, gibberellic acid, abscisic acid, and acetylcholine. Journal of Plant Physiology, 137(6):706-710.
- Broumand P, Rezaei A, Soleymani A, Shahrajabian MH, Noory A. 2010. Influence of forage clipping and top dressing of nitrogen fertilizer on grain yield of cereal crops in dual purpose cultivation system. Research on Crops, 11(3):603-613.
- Chattopadhyay KK, Bhattacharyya RN, Basu PS. 1984. Relation of gibberellic acid-induced growth of wheat coleoptiles with indole acetic acid. Biochemie und Physiologie der Pflanze, 179(1-2):123-128.
- Chen J, Sun Z, Zhang Y, Zeng X, Qing C, Liu J, Li L, Zhang H. 2009. Synthesis of gibberellin derivatives with anti-tumor bioactivities. Bioorganic and Medicinal Chemistry Letters, 19:5496-5499.

- Cornea-Cipcigan M, Pamfil D, Sisea CR, Margaoan R. 2020. Gibberellic acid can improve seed germination and ornamental quality of selected *Cyclamen* species grown under short and long days. *Agronomy*, 10(4):516.
- Cui H, Shahrajabian MH, Kuang Y, Zhang H, Sun W. 2023. Heterologous expression and function of cholesterol oxidase: A review. *Protein and Peptide Letters*, 30(7):531-540.
- Cuiying W, Bao Y, Wang QQ, Zhang HX. Introduction of the rice CYP714D1 gene into *Populus* inhibits expression of its homologous genes and promotes growth, biomass production and xylem fibre length in transgenic trees. *Journal of Experimental Botany*, 64:2847-2857.
- Dimovska V, Petropoulos VI, Salamovska A, Ilieva F. 2014. Flame seedless grape variety (*Vitis vinifera* L.) and different concentration of gibberellic acid (GA<sub>3</sub>). *Bulgarian Journal of Agricultural Science*, 20:127-132.
- Elahi NN, Raza S, Rizwan MS, Albalawi BFA, Ishaq MZ, Ahmed HM, Mehmood S, Imtiaz M, Farooq U, Rashid M, Ditta A. 2023. Foliar application of gibberellin alleviates adverse impacts of drought stress and improves growth, physiological and biochemical attributes of canola (*Brassica napus* L.). *Sustainability*, 15(1):78.
- El-Tohamy WA, Dasgan HY, Gruda NS. 2023. Impact of gibberellic acid on water status, growth, and development of cape gooseberry in newly reclaimed sandy lands within arid regions. *Horticulturae*, 9(12):1283.
- Endler PC, Scherer-Pongratz W, Lothaller H, Stephen S. 2015. Wheat and ultra high diluted gibberellic acid-further experiments and re-analysis of data. *Homeopathy*, 104(4):257-262.
- Gao F, Hu T, Tan W, Yu C, Li Z, Zhang L, Duan L. 2016. Photoprotectant improves photostability and bioactivity of abscisic acid under uv radiation. *Journal of Photochemistry and Photobiology B: Biology*, 158:99-104.
- Ghanbary E, Asiabani Z, Hosseini N, Kiaie SH, Kaki S, Ghasempour H, Babakhanian A. 2020. The development of a new modified graphite pencil electrode for quantitative detection of gibberellic acid (GA<sub>3</sub>) herbal hormone. *Microchemical Journal*, 157:105005.
- Goel B, Tripathi N, Bhardwaj N, Kumar A, Jian SK. 2023. Iodine-mediated one-step synthesis of ipomone from gibberellic acid. *Natural Product Research*, 37(19):3237-3244.
- Gokdere M, Ates S. 2014. Extractive fermentation of gibberellic acid with free and immobilized *Gibberella fujikuroi*. *Preparative Biochemistry and Biotechnology*, 44(1):80-89.
- Gonzalez-Lopez O, Casquer PA. 2014. Effects of GA<sub>3</sub> pregerminative treatment on *Gentiana lutea* L. var. *Aurantiaca* germination and seedlings morphology. *The Scientific World Journal*, 2014:751279.
- Gupta R, Chakrabarty SK. 2013. Gibberellic acid in plant: Still a mystery unresolved. *Plant Signaling and Behavior*, 8: e25504.
- Gupta R, Chakrabarty SK. 2013. Gibberellic acid in plant: Still a mystery unresolved. *Plant Signaling and Behavior*, 8:e25504.
- Hamayun M, Khan SA, Khan AL, Shin JH, Ahmad B, Shin DH, Lee IJ. 2010. Exogenous gibberellic acid reprogram soybean to higher growth and salt stress tolerance. *Journal of Agricultural and Food Chemistry*, 58:7226-7232.
- Hara M, Furukawa J, Sato A, Mizoguchi T. 2012. Abiotic stress and role of salicylic acid in plants. In *Abiotic Stress Responses in Plants*, 235-251. Springer, New York, NY.

- Hartung H, Schiestl S, Matzer W, Endler PC. 2010. Wheat germination (20H) and extremely diluted gibberellic acid ( $10^{-30}$ ): Explorative experiments on a fundamental homeopathy research model. *European Journal of Integrative Medicine*, 2(4):244-245.
- Hedden P, Thomas SG. 2012. Gibberellin biosynthesis and its regulation. *Biochemical Journal*, 444:11-25.
- Hedden P, Sponsel V. 2015. A century of gibberellin research. *Journal of Plant Growth Regulation*, 34:740-760.
- Hooley R. 1994. Gibberellins: Perception, transduction and responses. *Plant Molecular Biology*, 26:1529-1555.
- Horre BA, Jamesonll PE, Bannister P. 1989. Response of juvenile *Pseudopanax crassifolius* to gibberellic acid and paclobutrazol. *New Zealand Journal of Botany*, 27(4):591-594.
- Hou S, Zhu J, Ding M, Lv, G. 2008. Simultaneous determination of gibberellic acid, indole-3-acetic acid and abscisic acid in wheat extracts by solid-phase extraction and liquid chromatography-electrospray tandem mass spectrometry. *Talanta*, 76(4):798-802.
- Ingram TJ, Reid JB, Macmillan J. 1986. The quantitative relationship between gibberellin A1 and internode growth in *Pisum sativum* L. *Planta*, 168:414-420.
- Iqbal M, Ashraf M. 2013. Gibberellic acid mediated induction of salt tolerance in wheat plants: Growth, ionic partitioning, photosynthesis, yield and hormonal homeostasis. *Environmental and Experimental Botany*, 86:76-85.
- Jaleel CA, Gopi R, Manivannan P, Sankar B, Kishorekumar A, Panneerselvam R. 2007. Antioxidant potentials and ajmalicine accumulation in *Catharanthus roseus* after treatment with gibberellic acid. *Colloids and Surfaces B: Biointerfaces*, 60:195-200.
- Jaleel CA, Gopi R, Manivannan P, Sankar V, Kishorekumar A, Panneerselvam R. 2007. Antioxidant potentials and ajmalicine accumulation in *Catharanthus roseus* after treatment with gibberellic acid. *Colloids and Surfaces B: Biointerfaces*, 60:195-200.
- Jung H, Jo SH, Jung WY, Park HJ, Lee A, Moon JS, Seong SY, Kim JK, Kim YS, Cho HS. 2020. Gibberellin promotes bolting and flowering via the floral integrators RsFT and RsSOC1-1 under marginal vernalization in radish. *Plants*, 9:594.
- Kato F, Araki M, Miyazawa Y, Fujii N, Takeda K, Suge H, Takahashi H. 2011. Factors responsible for deep-sowing tolerance in wheat seedlings: Varietal differences in cell proliferation and the co-ordinated synchronization of epidermal cell expansion and cortical cell division for the gibberellin-mediated elongation first internodes. *Annals of Botany*, 108:439-447.
- Khoshkharam M, Rezaei A, Soleymani A, Shahrajabian MH. 2010. Effects of tillage and residue management on yield components and yield of maize in second cropping after barley. *Research on Crops*, 11(3):659-666.
- Kirkwood PS, Macmillan J, Sinnott ML. 1980. Rearrangement of the lactone ring of gibberellin-A<sub>3</sub> in aqueous alkali-Participation of the ionized 3-hydroxy-group in an anti-S<sub>N</sub>2' reaction. *Journal of the Chemical Society, Perkin Transactions 1*, 1:2117-2121.
- Kondo S, Danjo C. 2001. Cell wall polysaccharide metabolism during fruit development in sweet cherry Satohnishiki as affected by gibberellic acid. *Journal of the Japanese Society for Horticultural Science*, 70(2),178-184.

- Larson AJS, Cartwright MM, Jones, WD, Luce K, Chen M-Y, Petersen K, Nelson SV, Michaelis DJ, Madsen MD. Slow release of GA<sub>3</sub> hormone from polymer coating overcomes seed dormancy and improves germination. *Plants*, 12(24):4139.
- Li Z, Chen H, Guan Q, Li L, Xuan YH. 2024. Gibberellic acid signaling promotes resistance to saline-alkaline stress by increasing the uptake of ammonium in rice. *Plant Physiology and Biochemistry*, 207:108424.
- Liu C, Zhu L, Yin X, Xu Z, Li Q. 2013. Study on the gibberellic acid residues in brewing. *Journal of the American Society of Brewing Chemists*, 71(2):76-82.
- Mander LN, Sherburn M, Camp D, King RW, Evans LT, Pharis RP. 1998. Effects of D-ring modified gibberellins on flowering and growth in *Lolium temulentum*. *Phytochemistry*, 49:2195-2206.
- Mander LN. 2003. Twenty years of gibberellin research electronic supplementary information (ESI) available: Functionality patterns for naturally occurring gibberellins (gan). *Natural Product Reports*, 20:49-69.
- Martin DN, Proebsting WM, Parks TD, Dougherty WG, Lange T, Lewis JM, Gaskin P, Hedden P. 1996. Feed-back regulation of gibberellin biosynthesis and gene expression in *Pisum sativum* L. *Planta*, 200:159-166.
- Martinez-Garcia JF, Santes CM, Garcia-Martinez JL. 2010. The end-of-day far-red irradiation increases gibberellin A1 content in cowpea (*Vigna sinensis*) epicotyls by reducing its inactivation. *Physiologia Plantarum*, 108:426-434.
- Mathur M, Satpathy M, Sachar RC. 1992. Phytohormonal regulation of S-adenosylmethionine synthetase by gibberellic acid in wheat aleurones. *Biochimica et Biophysica Acta (BBA)-Molecular Cell Research*, 1137(3):338-348.
- Matzer W, Hartmann M, Endler PC. 2010. Wheat growth (7 days) and extremely diluted gibberellic acid (10e-30): Repetitive experiments on a fundamental homoeopathy research model. *European Journal of Integrative Medicine*, 2(4):245.
- Meldal M, Tornoe CW. 2008. Cu-catalyzed azide-alkyne cycloaddition. *Chemical Reviews*, 108:2952-3015.
- Miceli A, Moncada A, Sabatino L, Vetrano. 2019. Effect of gibberellic acid on growth, yield and quality of leaf lettuce and rocket grown in a floating system. *Agronomy*, 9(7): 382.
- Mimi CO, Sousa MC, Correa PLC, Chacon IDIC, Boaro CSF, Ferreira G. Impact of GA<sub>3</sub> on sugar and lipid degradation during *Annona x atemoya* Mabb. seed germination. *Horticulturae*, 9(3):388.
- Misratia KM, Islam MR, Ismail MR, Oad FC, Hanafi MM, Puteh A. Interactive effects of gibberellic acid (GA<sub>3</sub>) and salt stress on growth and ion accumulation of two rice (*Oryza sativa* L.) varieties differing in salt tolerance. *Journal of Food, Agriculture and Environment*, 13:66-70.
- Moumita, Al-Mahmud J, Biswas PK, Nahar K, Fujita M, Hasanuzzaman M. 2019. Exogenous application of gibberellic acid mitigates drought-induced damage in spring wheat. *Acta Agrobotanica*, 72:1776.
- Muniandi SKM, Hossain MdA, Abdullah MP, Shukor NAAb. 2018. Gibberellic acid (GA<sub>3</sub>) affects growth and development of some selected kenaf (*Hibiscus cannabinus* L.) cultivars. *Industrial Crops and Products*, 118:180-187.
- Murase K, Hirano Y, Sun TP, Hakoshima T. 2008. Gibberellin-induced DELLA recognition by the gibberellin receptor gid1. *Nature*, 456:U459-U463.

- Ogbaji PO, Shahrajabian MH, Xue X. 2013. Changes in germination and primarily growth of three cultivars of tomato under diatomite and soil materials in auto-irrigation system. *International Journal of Biology*, 5(3):80-84.
- Oh K, Yamada K, Asami T, Yoshizawa Y. 2012. Synthesis of novel brassiosteroid biosynthesis inhibitors based on the ketoconazole scaffold. *Bioorganic and Medicinal Chemistry Letters*, 22:1625-1628.
- O'Neill DP, Ross JJ, Reid JB. 2000. Changes in gibberellin A1 levels and response during de-etiolation of pea seedlings. *Plant Physiology*, 124:805-812.
- Pan B, Zhong T, Zhao G. 2017. Promoting deep-sowing germinability of corn (*Zea mays* L.) by seed soaking with gibberellic acid. *Archives of Agronomy and Soil Science*, 63(9):1314-1323.
- Pavlista AD, Baltensperger DD, Santra DK, Hergert GW, Knox S. 2014. Gibberellic acid promotes early growth of winter wheat and rye. *American Journal of Plant Sciences*, 05:2984-2996.
- Peng D, Chen X, Yin Y, Lu KL, Yang WB, Tang YH, Wang ZL. 2014. Lodging resistance of winter wheat (*Triticum aestivum* L.): lignin accumulation and its related enzymes activities due to the application of paclobutrazol or gibberellin acid. *Field Crops Research*, 157:1-7.
- Pfleger A, Hofacker J, Scherer-Pongratz W, Lothaller H, Reich C, Endler PC. 2011. The effect of extremely diluted agitated gibberellic acid (10e-30) on wheat stalk growth- A two researcher pilot study. *Complementary Therapies in Medicine*, 19(3):164-169.
- Phuoc LT, Mander LN, Koshioka M, Oyama-Okubo N, Nakayama M, Ito A. 2008. Confirmation of structure and synthesis of three new 11 $\beta$ -OC C<sub>20</sub> gibberellins from loquat fruit. *Tetrahedron*, 64:4835-4851.
- Rebetzke GJ, Bonnett DG, Ellis MH. 2012. Combining gibberellic acid-sensitive and insensitive dwarfing genes in breeding of higher-yielding, sesqui-dwarf wheats. *Field Crops Research*, 127:17-25.
- Riazat A, Soleymani A, Shahrajabian MH. 2012. Changes in seed yield and biological yield of six wheat cultivars on the basis of different sowing dates. *Journal of Food, Agriculture and Environment*, 10(1):467-469.
- Rodrigues C, Vandenberghe LPDS, Oliveira JD, Soccol CR. 2012. New perspectives of gibberellic acid production: a review. *Critical Reviews in Biotechnology*, 32(3):263-273.
- Saluja D, Berry M, Sachar RC. 1987. Inorganic phosphate mimics the specific action of gibberellic acid in regulating the activity of monophenolase in embryo-less half-seeds of wheat. *Phytochemistry*, 26(3):611-614.
- Saluja D, Ansari AI, Sood A, Sachar RC. 1989. Early response to gibberellic acid of monophenolase (tyrosinase) activity in de-embryonated half-seeds of wheat. *Phytochemistry*, 28(2):341-344.
- Sauter M, Kende H. 1993. Internodal elongation and orientation of cellulose microfibrils and microtubules in deepwater rice. *Planta*, 190:354-362.
- Sayed AEDH, AbdAllah EA, Hamed M, Soiliman HAM. 2020. Hepatonephrotoxicity in late juvenile of *Oreochromis niloticus* exposed to gibberellic acid: Ameliorative effects of *Spirulina platensis*. *Pesticide Biochemistry and Physiology*, 167:104600.
- Scott RS. 1959. Effect of gibberellic acid and nitrogen on winter growth of pasture. *New Zealand Journal of Agricultural Research*, 2(6):1203-1210.
- Seo Y, Gudz A, Lowe JM, Gagne MR. 2019. Selective deoxygenation of gibberellic acid with fluoroarylborane catalysts. *Tetrahedron*, 75(49):130712.

- Shahrajabian MH, Soleymani A, Naranjani L. 2011. Grain yield and forage characteristics of forage sorghum under different plant densities and nitrogen levels in second cropping after barley in Isfahan, Iran. *Research on Crops*, 12(1):68-78.
- Shahrajabian MH, Chaski C, Polyzos N, Petropoulos SA. 2021. Biostimulants application: A low input cropping management tool for sustainable farming of vegetables. *Biomolecules*, 11(5):698.
- Shahrajabian MH, Chaski C, Polyzos N, Tzortzakis N, Petropoulos SA. 2021. Sustainable agriculture systems in vegetable production using chitin and chitosan as plant biostimulants. *Biomolecules*, 11(6):819.
- Shahrajabian MH, Sun W. 2023a. Great health benefits of essential oils of pennyroyal (*Mentha pulegium* L.): A natural and organic medicine. *Current Nutrition and Food Science*, 19(4):340-345.
- Shahrajabian MH, Sun W. 2023b. Survey on multi-omics, and multi-omics data analysis, integration and application. *Current Pharmaceutical Analysis*, 19(4):267-281.
- Shahrajabian MH, Sun W. 2023c. The importance of salicylic acid, humic acid and fulvic acid on crop production. *Letters in Drug Design and Discovery*, 2023(20):1-16.
- Shahrajabian MH, Sun W, 2023d. Various techniques for molecular and rapid detection of infectious and epidemic diseases. *Letters in Organic Chemistry*, 20(9):779-801.
- Shahrajabian MH, Sun W. 2023e. The golden spice for life: Turmeric with the pharmacological benefits of curcuminoids components, including curcumin, bisdemethoxycurcumin, and demethoxycurcumin. *Current Organic Synthesis*, 20.
- Shahrajabian MH, Sun W. 2023f. Five important seeds in traditional medicine, and pharmacological benefits. *Seeds*, 2(3):290-308.
- Shahrajabian MH, Sun W. 2023g. Study of different types of fermentation in wine-making process and considering aromatic substances and organic acid. *Current Organic Synthesis*, 20.
- Shahrajabian MH, Sun W, 2023h. Iranian traditional medicine (ITM) and natural remedies for treatment of the common cold and flu. *Reviews on Recent Clinical Trials*, 18:1-10.
- Shahrajabian MH, Sun W. 2023i. The importance of traditional Chinese medicine in the intervention and treatment of HIV while considering its safety and efficacy. *Current HIV Research*, 21(6):1-16.
- Shahrajabian MH, Petropoulos SA, Sun W. 2023. Survey of the influences of microbial biostimulants on horticultural crops: Case studies and successful paradigms. *Horticulturae*, 9(193):1-24.
- Shahzad K, Hussain S, Arfan M, Hussain S, Waraich EA, Zamir S, Saddique M, Rauf A, Kamal KY, Hano C, El-Esawi, MA. 2021. Exogenously applied gibberellic acid enhances growth and salinity stress tolerance of maize through modulating the morpho-physiological, biochemical and molecular attributes. *Biomolecules*, 11(7):1005.
- Shakirova FM, Sakhabutadinova AR, Bezrukova MV, Fatkhutdinova RA, Fatkhutdinova DR. 2003. Changes in the hormonal status of wheat seedlings induced by salicylic acid and salinity. *Plant Science*, 164(3):317-322.
- Shan F, Zhang R, Zhang J, Wang C, Lyu X, Xin T, Yan C, Dong S, Ma C, Gong Z. 2021. Study on the regulatory effects of GA3 on soyeabn internode elongation. *Plants*, 10(8):1737.
- Shayanfar M, Soleymani A, Shahajabian MH. 2011. Effect of plant densities on solar radiation absorption and depreciation, leaf area index and crop growth rate of different cultivars of sunflower. *Research on Crops*, 12(3):728-730.

- Shen C, Wang XJ, Guo FT, Zhou WQ, Wang GL. 2020. Gibberellin and the plant growth retardant Paclobutrazol altered fruit shape and ripening in tomato. *Protoplasma*, 257:853-861.
- Shu K, Zhou W, Chen F, Luo X, Yang W. 2018. Abscisic acid and gibberellins antagonistically mediate plant development and abiotic stress responses. *Frontiers in Plant Science*, 9:416.
- Singh B, Usha K. 2003. Salicylic acid induced physiological and biochemical changes in wheat seedlings under water stress. *Plant Growth Reg*, 39:137-141.
- Skalicky M, Kubes J, Vachova P, Hajjhashemi S, Martinkova J, Hejnak V. 2020. Effect of gibberellic acid on growing-point development of non-vernalized wheat plants under long-day conditions. *Plants*, 9(12):1735.
- Soleymani A, Shahri MM, Shahrajabian MH, Naranjani L. 2010. Responses of cultivars of canola to sulfur fertilizer and plant densities under climatic condition of Gorgan region, Iran. *Journal of Food, Agriculture and Environment*, 8(3&4):298-304.
- Soleymani A, Shahrajabian MH, Naranjani L. 2011. The effects of nitrogen as starter fertilizer on ash percentage, important elements and solar radiation absorption of berseem clover cultivars intercropped by corn. *Journal of Food, Agriculture and Environment*, 9(1):342-344.
- Soleymani A, Shahrajabian MH, Naranjani L. 2013. Effect of planting dates and different levels of nitrogen seed yield and yield components of nuts sunflower (*Helianthus annuus* L.). *African Journal of Agricultural Research*, 8(46):5802-5805.
- Soleymani A, Shahrajabian MH. 2017. Effects of planting dates and row distance on sugar content, root yield, and solar radiation absorption in sugar beet at different plant densities. *Romanian Agricultural Research*, 34(34):1-11.
- Srivastava A, Handa AK. 2005. Hormonal regulation of tomato fruit development: A molecular perspective. *Journal of Plant Growth Regulation*, 24:67-82.
- Sun W, Shahrajabian MH, Cheng Q. 2021a. Natural Dietary and medicinal plants with anti-obesity therapeutics activities for treatment and prevention of obesity during lock down and in post-Covid-19 era. *Applied Sciences*, 11(17):7889.
- Sun W, Shahrajabian MH, Cheng Q. 2021b. Archaea, bacteria and termite, nitrogen fixation and sustainable plants production. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, 49(2):1-32.
- Sun W, Shahrajabian MH, Cheng Q. 2021c. Organic waste utilization and urban food waste composting strategies in China- A review. *Notulae Scientia Biologicae*, 13(2):1-16.
- Sun W, Shahrajabian MH. 2023. The application of arbuscular mycorrhizal fungi as microbial biostimulant, sustainable approaches in modern agriculture. *Plants*, 12(17):3101.
- Sun W, Shahrajabian MH, Petropoulos SA, Shahrajabian N. 2023. Developing sustainable agriculture systems in medicinal and aromatic plant production by using chitosan and chitin-based biostimulants. *Plants*, 12(13):2469.
- Sun W, Shahrajabian MH. 2024. Survey on nitrogenase evolution by considering the importance of nitrogenase, its structure, and mechanism of nitrogenase. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, 52(1):13157.
- Sun W, Shahrajabian MH, Kuang Y, Wang N. 2024. Amino acids biostimulants and protein hydrolysates in agricultural sciences. *Plants*, 13(210):1-22.
- Sun W, Shahrajabian MH, Soleymani A. 2024. The roles of plant-growth-promoting rhizobacteria (PGPR)-based biostimulants for agricultural production systems. *Plants*, 13(613):1-38.

- Tuna AL, Kaya C, Dikilitas M, Higgs D. 2008. The combined effects of gibberellic acid and salinity on some antioxidant enzyme activities, plant growth parameters and nutritional status in maize plants. *Environmental and Experimental Botany*, 62:1-9.
- Uslu H. 2013. Extraction of gibberellic acid from aqueous solution by trioctyl amine (TOA). *Separation Science and Technology*, 48(3):487-492.
- Wang GL, Que F, Xu ZS, Wang F, Xiong AS. 2015. Exogenous gibberellin altered morphology, anatomic and transcriptional regulatory networks of hormones in carrot root and shoot. *BMC Plant Biology*, 15:290-302.
- Wang LL, Chen, XY, Yang Y, Wang Z, Xiong F. 2016. Effects of exogenous gibberellic acid and abscisic acid on germination, amylases, and endosperm structure of germinating wheat seeds. *Seed Science and Technology*, 44:64-76.
- Wu Y, Gong W, Yang W. 2017. Shade inhibits leaf size by controlling cell proliferation and enlargement in soybean. *Scientific Reports*, 7:9259.
- Xu D, Cao H, Fang W, Pan J, Chen J, Zhang J, Shen W. 2017. Linking hydrogen-enhanced rice Aluminum tolerance with reestablishment of GA/ABA balance and MiRNA-modulated gene expression: A case study on germination. *Ecotoxicology and Environmental Safety*, 145:303-312.
- Yazdpour H, Shahri MM, Soleymani A, Shahrajabian MH. 2012. Effects of harvesting time and harvesting height on grain yield and agronomical characters in rice ratoon (*Oryza sativa* L.). *Journal of Food, Agriculture and Environment*, 10(1):438-440.
- Yuan Y, Yang WY. 2007. Effects of shading on Endogenous hormone of *Cardiocrinum giganteum*. *North Horticulture*, 8:123-125.
- Zalnierius T, Sveikauskas V, Aphalo PJ, Gaveliene V, Buda V, Jurkoniene S. 2022. Gibberellic acid (GA<sub>3</sub>) applied to flowering *Heracleum sosnowskyi* decreased seed variability even it seed development is not inhibited. *Plants*, 11(3):314.
- Zhang J, Smith DL, Liu W. 2011. Effects of shade and drought stress on soybean hormones and yield of main-stem and branch. *African Journal of Biotechnology*, 10:14392-14398.
- Zhang Y, Zhang H, Chen J, Zhao H, Zeng X, Zhang H, Qing C. 2012. Antitumor and antiangiogenic effects of ga-13315, a gibberellin derivative. *Investigational New Drugs*, 30:8-16.
- Zhang HX, Wang HH, Zhu Q, Gao YB, Wang HY, Zhao LZ, Wang, YS, Xi FH, Wang WF, Yang YQ, Lin C, Gu L. 2018. Transcriptome characterization of moso bamboo (*Phyllostachy edulis*) seedlings in response to exogenous gibberellin applications. *BMC Plant Biology*, 18:125-140.
- Zhang R, Shan FX, Wang C, Yan C, Dong SK, Xu Y, Gong ZP, Ma CM. 2020. Internode elongation pattern, internode diameter and hormone changes in soybean (*Glycine max*) under different shading conditions. *Crop and Pasture Science*, 7:679-688.
- Zhou F, Ran J, Tan W, Li Z, Duan L. 2013. Synthesis and biological activity of aryamido-cyclopropanecarboxylic acid. *Chinese Journal of Pesticide Science*, 5:490-495.
- Zhou R, Wang XZ, Chen HF, Zhang XJ, Shan ZH, Wu XJ, Cai SP. 2014. QTL analysis of lodging and related traits in soybean. *Acta Agronomica Sinica*, 35:57-65.
- Zwar JA, Chandler PM. 1995.  $\alpha$ -Amylase production and leaf protein synthesis in a gibberellin-responsive dwarf mutant of Himalaya barley (*Hordeum vulgare* L.). *Planta*, 197:39-48.