



Trends in Phytochemical Research (TPR)

Journal Homepage: <https://sanad.iau.ir/journal/tptr>



Original Research Article

Antioxidant properties and anticancer activity of *Olea europaea* L. olive and *Ficus carica* fruit extracts against pancreatic cancer cell lines

NADIA DEKDOUK¹, SOUAD AMEDDAH¹, CHAWKI BENSOUICI², MEGHBOUN IBTISSEM², MENAD AHMED¹, AND FÁTIMA MARTEL^{3,4}

¹Laboratoire de Biologie et Environnement, Faculté des Sciences de la Nature et de la Vie, Université des Frères Mentouri Constantine1, Algérie

²Biotechnology Research Center (CRBt), Ali Mendjli, Constantine, Algérie

³Unit of Biochemistry, Department of Biomedicine, Faculty of Medicine, University of Porto, Porto, Portugal

⁴Instituto de Investigação e Inovação em Saúde (i3S), University of Porto, Porto, Portugal

ABSTRACT

Olea europaea (OFE) and *Ficus carica* (FFE) extracts have shown anticancer activity. In the current report, we aim to evaluate the phytochemical composition, antioxidant properties and anticancer activity of OFE and FFE against PANC-1 and AsPC-1 pancreatic cancer cell lines. OFE possessed higher phenolic, flavonoid content and antioxidant ability than FFE. OFE and FFE induced a cytotoxic effect and their combination resulted in an increased cytotoxicity. OFE showed an antiproliferative effect while FFE had no effect. Combination of the two extracts resulted in an antiproliferative effect similar to that of OFE. OFE showed an anti-migratory effect, while FFE exhibited a promigratory effect. Interestingly, combination of the two extracts resulted in an anti-migratory effect similar to that caused by OFE. In view of the findings of this study, these extracts could be considered potential sources of natural compounds with antioxidant and anti-pancreatic cancer effects. The combination of these extracts should be further investigated in the context of pancreatic cancer.

ARTICLE HISTORY

Received: 16 November 2023

Revised: 25 May 2024

Accepted: 12 June 2024

ePublished: 28 June 2024

KEYWORDS

Antioxidant capacity
Anti-pancreatic cancer effect
Antiproliferative
Cytotoxic
Fig extract
Olive extract
Phytochemical composition

1. Introduction

Medicinal plants have shown great promise in the fight against cancer, with many studies demonstrating their potent anticancer properties. Several naturally occurring plant-based compounds, such as curcumin, resveratrol, and quercetin, have exhibited anticancer effects and are being investigated as potential adjuvant chemotherapy agents (Greenwell and Rahman, 2015; Khan et al., 2019; Mahdavi and Mohammadhosseini 2022). These plant-derived compounds often have lower toxicity towards healthy cells and a tendency to target abnormal or cancerous cells specifically. Many of the drugs currently on the market are structurally similar to natural compounds found in medicinal plants (Khan et al., 2019; Obafemi and Besong, 2023). Herbal compounds possess various beneficial qualities, including antioxidant, anti-inflammatory, antimutagenic, and apoptosis-

inducing activities, which aid in the early diagnosis and treatment of cancer. Adequate dietary intake of these herbal products may help prevent and treat cancer by causing cell cycle arrest, inducing apoptosis, regulating carcinogen metabolism and oncogenic expression, inhibiting cell adhesion, proliferation, and migration, and blocking signaling pathways crucial for cancer progression (Obafemi and Besong, 2023). Plants produce a wide variety of secondary metabolites, such as tannins, alkaloids, terpenoids, flavonoids, pigments, and pigment-like molecules, which have been shown to have biological effects on hematopoietic cells, lipids, and cardiovascular systems, as well as anti-inflammatory, anticancer, and contraceptive effects (Mohammadhosseini et al., 2019a; Mohammadhosseini et al., 2019b; Mohammadhosseini et al., 2021). Pancreatic cancer is one of the most aggressive cancers with poor prognosis and survival despite surgical and adjuvant therapies. Currently, it constitutes the fourth

✉ Corresponding author: Fátima Martel

Tel: 351 22 0426654; Fax: 351 22 5513624

E-mail address: fmartel@med.up.pt, [doi:](https://doi.org/)

cause of cancer mortality in occidental countries (Siegel et al., 2016). Treatment with gemcitabine, a pyrimidine agonist, has been the standard of care for metastatic pancreatic cancer for more than a decade. However, survival for most patients treated with this drug has not improved beyond 6 months (Moore et al., 2007; Park et al., 2021). One of the main problems posed with anticancer drugs is that they target not only tumor cells, but also other cell types, leading to toxicity to both abnormal and normal cells (Baldo and Pham, 2014). Another problem related to anticancer therapy is the development of resistance of cancer cells against anticancer treatments (Bar-Zeevet et al., 2017). This is the case of pancreatic cancer cells, which are highly resistant to current chemotherapy and radiotherapy treatments (Vendrey et al., 2017). Therefore, finding new therapeutic options is urgently needed to help manage this deadly cancer.

Nutrition and diet are good candidate agents for protection against various types of cancer (Kazeminia et al., 2022). Epidemiological studies have shown a reduction in the risk of pancreatic cancer through increased consumption of fruits and vegetables (Zheng et al., 2017; Gianfredi et al., 2022). The Mediterranean diet has been recognized as a healthy diet with a preventive effect against chronic diseases associated with oxidative damage, including cardiovascular disease, diabetes, and cancer (Schwingshackl and Hoffmann, 2014; Saha et al., 2017; Mentella et al., 2019; Guasch-Ferré and Willett, 2021). These effects seem to be mediated by many individual components of this diet, including polyphenolic phytochemicals present in fruits such as figs and olive oil (Owen et al., 2000b; Solomon et al., 2006; Lopez-Miranda, 2010).

The olive tree, *Olea europaea*, produces the olive fruit and is one of the vital components of the Mediterranean diet. Olive and its products have many health benefits, including cancer (Markellos et al., 2022; Romani et al., 2019). Preclinical *in vitro* and *in vivo* studies have demonstrated an antitumoral effect of olive fruit extracts (OFE) against hepatic and colorectal cancer (Maalejet et al., 2017; Shirazi et al., 2018; Celano et al., 2019). The components of olives and olive oil that have been most consistently associated with a protective effect on carcinogenesis are vitamin E, squalene, and phenolic compounds (Guo et al., 2018). The most abundant phenolic compounds in olive oil are the lignans (+)-1-acetoxypinoresinol and (+)-pinoresinol, followed by the oleuropein-derived simple phenols hydroxytyrosol and tyrosol and the secoiridoids oleuropein, ligstroside aglycone and their respective decarboxylated dialdehyde derivatives (Owen et al., 2004). In contrast, polyphenols such as the flavonoid luteolin, the phenolic acid gallic acid, the phenolic alcohols hydroxytyrosol and tyrosol and the secoiridoids oleuropein and verbascoside are most abundant in olive fruits (Owen et al., 2004; Dekdouk et al., 2015). Previous studies have clearly evidenced an antitumor effect of these phenolic compounds against certain types of cancers, namely breast (Han et al., 2009; González et al., 2013; Carrera- Akl et al. 2014; Sepporta et al., 2014), colorectal (Owen et al., 2000a; Carrera-González et al., 2013; Hormozi et al., 2020), gastric (Kountouri et al.,

2009), hepatic (Pei et al., 2016), skin cancer (Carrera-González et al., 2013) and leukemia (Fabiani et al., 2002), and also pancreatic cancer (LeGendre et al., 2015; Goldsmith et al., 2018).

Ficus carica is a species characteristic of the Mediterranean region where its cultivation and use constitute an ancient tradition. Figs are consumed both fresh and dried (Baby and Justin, 2011). Figs are an excellent source of minerals, vitamins, carbohydrates, and dietary fiber, they are fat- and cholesterol-free and rich in several amino acids (Cruz et al., 2022; Olaoluwa et al., 2022; Ijoma et al., 2023; Popwo Tameye et al., 2023). Figs have traditionally been used for their medicinal benefits as laxative and cardiovascular, respiratory, antispasmodic and anti-inflammatory remedies (Guarrera, 2005). However, figs also exhibit anticancer properties (Morovatiet al., 2022). Namely, preclinical *in vitro* and *in vivo* models revealed anticancer activity of *F. carica* latex or fruit extracts (FFE) in relation to several types of cancers, including breast (Ghandehari et al., 2018), colorectal (Soltana et al., 2019), gastric (Hashemi et al., 2011), hepatic (Purnamasariet al., 2019) and, recently, also in relation to pancreatic cancer (Ou et al., 2022). An important organic compound in figs is 2-benzhydrylsulfinyl-*N*-hydroxyacetamide-Na, which is less abundant in other fruits and plants, and exhibits anticancer activity *in vitro* (Al Salman et al., 2020). Moreover, *Ficus carica* is a rich source of polyphenols. Quercetin and quercetin-related compounds are the main phenolic compounds present in this fruit (Khadhraoui et al., 2019; Arvaniti et al., 2019), and are well known to possess antitumor activity against several types of cancer, including pancreatic cancer (Rauf et al., 2018).

Several *in vitro* and *in vivo* preclinical studies have demonstrated varying degrees of efficacy of natural products against pancreatic cancer (Singh et al., 2015), and numerous dietary and pharmacological agents have been proposed as alternative strategies for the prevention and/or treatment of this disease (Dhar et al., 2018; Selvarajoo et al., 2022).

To the best of our knowledge, no report on the effects of a combination of OFE and FFE on pancreatic cancer cells is available in the literature. Keeping this point in view, the present study was conducted to evaluate the *in vitro* antioxidant activities and the cytotoxic, antiproliferative and antimigratory effects of OFE and FFE, alone and in combination, on two pancreatic cancer cell lines.

2. Experimental

2.1. Chemicals and reagents

³H-Thymidine ([methyl-³H]-thymidine, 79 Ci/mmol) was purchased from GE Healthcare GmbH (Freiburg, Germany). Dimethylsulfoxide (DMSO), hydrochloric acid (HCl), triton X-100, trichloroacetic acid (TCA) were purchased from Merck (Darmstadt, Germany). Aluminium chloride (AlCl₃), 1,1-diphenyl-2-picrylhydrazyl (DPPH), gallic acid, Folin-Ciocalteu's phenol reagent, ascorbic acid, sodium carbonate (Na₂CO₃), potassium ferricyanide [K₃Fe(CN)₆], 2,2-azinobis-3-ethylbenzothiazoline-6-sulfonic acid (ABTS), potassium persulfate (K₂S₂O₈),



2,6-di-tert-butyl-4-hydroxy toluene (BHA), butylated hydroxytoluene (BHT), α -tocopherol, iron (III) chloride (FeCl_3), 3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide (MTT), antibiotic/antimicrobial solution (100 U/mL penicillin, 100 $\mu\text{g}/\text{mL}$ streptomycin, and 0.25 $\mu\text{g}/\text{mL}$ amphotericin B), Dulbecco's Modified Eagle's Medium-high glucose, fetal bovine serum, N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid (HEPES), RPMI 1640 medium, sodium hydroxide (NaOH) and trypsin-ethylenediamine-tetraacetic acid (EDTA) solution were purchased from Sigma-Aldrich (St. Louis, MO, USA).

2.2. Extraction and identification of phenolic compounds

2.2.1. Plant material

Fruits of the *F. carica* 'Mkaousi' cultivar were harvested in the Constantine region (Algeria) in September 2017. Olives of the *Chemlal* cultivar of *Olea europaea* were harvested in Algeria in December 2017. The olives were handpicked during the harvest season and were immediately frozen in liquid nitrogen to block enzyme activities. Healthy fruits (100 g), without any physical damage or type of infection, were selected, and lyophilized. A reference specimen for each sample is deposited at the Herbarium of the Faculty of Natural and Life Sciences (Université des Frères Mentouri Constantine, Algeria). The plant samples were authenticated by Mendil, M, Director General of Technical Institute of Fruit Arboriculture and Vine (ITAF), Algeria.

2.2.2. Extraction procedure

The lyophilized fruits of *O. europaea* (100 g) were defatted with hexane, and then extracted for 48 h with CHCl_3 , and MeOH, successively, to give 28.33, 2.7, and 5.37 g of the respective residues. The polyphenols from lyophilized fruits of *F. carica* were extracted by maceration using ethyl acetate to give 2.07 g. It has been demonstrated that this solvent is better than others to extract phenolics (Fki et al., 2005). Solvent was renewed 3 times and the obtained extracts were dried using a rotary evaporator.

2.2.3. Determination of total phenolic content (TPC)

The TPC of crude OFE and FFE was estimated spectrophotometrically using Folin-Ciocalteu method (Le et al., 2007). In brief, 200 μL of each extract were added to 1 mL of Folin-Ciocalteu reagent (10% (w/v)). After 4 min, 800 μL of sodium carbonate solution (75 g/L) was added. The mixture was allowed to react for 2 h in darkness at room temperature.

After incubation, the absorbance was recorded at 765 nm using UV-Vis spectrophotometer (VIS-7220G). Gallic acid (5-200 $\mu\text{g}/\text{mL}$) was used as reference to establish the calibration curve from which the concentration of polyphenols was calculated, and the results are expressed as TPC (mg EGA/100 g extract).

2.2.4. Determination of total flavonoid content (TFC)

The quantification of TFC was performed by the trichloroaluminum method (Turkoglu et al., 2007). Briefly, 1 mL of AlCl_3 (2.0 (w/v)) solution was added to 1 mL of OFE or FFE. The mixture was vigorously stirred and incubated for 10 min at room temperature, and then the absorbance of each sample was read at 430 nm. Quercetin (1.25-25 $\mu\text{g}/\text{mL}$) was used to establish the calibration curve used to estimate the concentration of flavonoids found in the crude extracts. The results are given as TFC (mg EQ/100 g extract).

2.3. Antioxidant activity of OFE and FFE

2.3.1. DPPH[•] (1,1-Diphenyl-2-picrylhydrazil) scavenging assay

The antiradical activity of OFE and FFE was evaluated by the free radical DPPH[•] assay (Blois, 1958). In brief, 25 μL of different dilutions of the extracts or standards BHA (butylated hydroxy anisole) and BHT (butylhydroxytoluene) were added to 975 μL of DPPH (0.025 mg/mL) prepared in methanol. The obtained mixture was incubated for 30 min in obscurity. Then, the absorbances were measured using a UV-Vis spectrophotometer at 517 nm and the percentage of DPPH[•] free radical-scavenging activity of each extract was calculated as follows, where I(%) is the percentage of inhibition and A_c and A_s are the absorbances of the control and the test sample, respectively (Eqn. 1):

$$I(\%) = [(A_c - A_s) / A_c] \times 100 \quad (\text{Eqn. 1})$$

The results were compared to the standard antioxidant BHA and BHT.

2.3.2. ABTS^{•+} (2,2'-Azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt) scavenging assay

The ABTS^{•+} scavenging activity of OFE and FFE was performed spectrophotometrically (Re et al., 1999). 2 mL of the diluted ABTS cation solution was added to 1 mL of sample solution: Extracts or standards (BHT and BHA) at different concentrations (0.0156-1 mg/mL). After 30 min of incubation, the absorbance was measured at 734 nm and the percentage of ABTS scavenging effect of each extract was calculated using Eqn. 1, where I(%) is the percentage of inhibition and A_c and A_s are the absorbances of the control and the test sample, respectively. The results were compared to the standard antioxidant BHA and BHT.

2.3.3. Cupric acid reducing antioxidant capacity (CUPRAC)

Fifty μL of a solution of copper(II) chloride (10 mM) were added to 50 μL of the neocuprine solution (7.5 mM) and 60 μL of ammonium acetate buffer solution (1 M, pH = 7.0). Different concentrations of OFE, FFE and standards (BHT and BHA) were added to the initial mixture to make a final volume of 200 μL . The samples were shielded from light and the absorbance was measured at 450 nm after 1 h of incubation. The results were calculated as A 0.5 ($\mu\text{g}/\text{mL}$) (the absorbance whose inhibition is equivalent to 50%), and the reduction capacity of the extracts was compared with those of BHA and BHT as

standards (Apak et al., 2004).

2.3.4. Reducing power assay

The reducing power of OFE and FFE were examined according to the previous method (Oyaizu, 1986). Accordingly, 100 μ L of sample solution (extracts or references) prepared at different concentrations were added to 0.5 mL of phosphate buffer (0.2 M pH 6.6) and 0.5 mL of potassium ferricyanide (1.0 (w/v)). The mixture was incubated at 50 °C for 20 min. Then, 500 μ L of CCl_3COOH (10% (w/v)) was added to the mixture, which was centrifuged (10 min/3000 rpm). A volume of 500 μ L of the supernatant solution was mixed with 500 μ L of distilled water and 125 μ L of FeCl_3 (1.0% (w/v)) freshly prepared. The absorptions were read at 700 nm and the results were calculated as A 0.5 (μ g/mL) (the absorbance whose inhibition is equivalent to 50%). The reducing power of the various extracts was compared with those of ascorbic and tannic acids as standards.

2.4. Anticancer effect OFE and FFE

2.4.1. Cell culture and treatments

PANC-1 cell line (human pancreatic epithelioid carcinoma cell line) and AsPC-1 cell line (human pancreatic ascites adenocarcinoma cell line) were obtained from the American Type Culture Collection (ATCC) and used between passage numbers 36-38. Cells were maintained in a humidified atmosphere of 5% CO_2 -95% air and were grown in Dulbecco's modified Eagle's medium supplemented with 4.5 g/L glucose, 2 mM L-glutamine, 10 mM sodium bicarbonate, 10 (v/v%) heat-inactivated fetal bovine serum and 1.0 (v/v%) antibiotic/antimycotic (PANC-1 cells) or in RPMI 1640 supplemented with 2 mM of L-glutamine, 10 mM of sodium bicarbonate, 1 mM of sodium pyruvate, 10 (v/v%) heat-inactivated fetal bovine serum and 1.0 (v/v%) antibiotic/antimycotic (AsPC-1 cells). Culture medium was changed every 2-3 days and the culture was split every 7 days. For sub-culturing, the cells were removed enzymatically (0.25 (w/v%) trypsin-EDTA, 37 °C), split 1:4 (PANC-1) or 1:3 (AsPC-1) and sub-cultured in plastic culture dishes (21 cm^2 ; \varnothing 60 mm, TPP®, Trasadingen, Switzerland). For determination of migration, viability and cell proliferation, cells were seeded on 24-well plastic cell culture dishes (2 cm^2 ; \varnothing 16 mm; TPP®, Trasadingen, Switzerland) and were used after 7 (PANC-1 cells) or 11 days (AsPC-1 cells) (90% confluence). The OFE and FFE to be tested were dissolved in DMSO. The final concentration of the solvent in the culture medium was 1.0 (v/v%). Controls for the extracts were run in the presence of this solvent.

2.4.2. Determination of cell viability

Cell viability was determined with the MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) assay. Briefly, cells were exposed to OFE and/or FFE (or to the solvent) for 24 h. After the treatment period, 30 μ L of MTT solution (5 mg/mL) was added to each well and the cells were further incubated for 3

h at 37 °C. Afterwards, the MTT solution was removed and the cells were lysed by the addition of 200 μ L of DMSO followed by plate shaking for 10 min at room temperature. Optical density (OD) for the solution in each well was determined at both 540 and 660 nm. The OD at 660 nm corresponds to unspecific light absorption and was subtracted from the OD at 540 nm to give the OD value specific to formazan crystals derived from MTT cleavage, which is proportional to the number of viable cells with active mitochondria.

2.4.3. Determination of cell proliferation

Cell proliferation rate was determined by the ^3H -thymidine incorporation assay. Briefly, cells were exposed to OFE and/or FFE (or the solvent) for 24 h, being incubated with ^3H -thymidine 0.025 μ Ci/mL in the last 5 h of treatment. After removal of excess ^3H -thymidine by a wash with 300 μ L of 10% TCA (w/v) for 1 h at 4 °C, drying for 30 min and addition of 1 M NaOH (280 μ L/well), the lysate was neutralized with 5 M HCl prior to the addition of scintillation fluid. The radioactivity of the samples was then quantified by liquid scintillometry (LKB Wallac 1209 Rackbeta, Turku, Finland). Cellular DNA synthesis rate was expressed as incorporation of ^3H -thymidine (μ Ci/mg protein).

2.4.4. Determination of cell migration rates

Cell migration rates were determined by a scratch injury assay. Briefly, the cell monolayers were scratched with a 10 μ L pipette tip and were afterwards treated for 24 h with OFE and/or FFE (or the solvent). Images were obtained at 0 and 24 h after the scratch, and quantification was performed using the ImageJ software (NIH, Bethesda, MD, USA).

2.4.5. Protein determination

The protein content of cell monolayers was determined as described by Bradford (1976), using human serum albumin as standard.

2.5. Calculations and statistics

Arithmetic means are given with standard error of the mean (SEM) or standard deviation (SD). The value of n indicates the number of replicates of at least 2 or 3 different replicates.

The extract/drug concentration for 50% inhibition (IC_{50}) was estimated by plotting percentage inhibition with respect to control against treatment concentration *in vitro* tests. $A_{0.5}$ values were determined from absorbance curves at the corresponding wavelength.

Statistical significance of the difference between various groups was evaluated by analysis of variance test followed by the Student-Newman-Keuls test. For comparison between two groups, Student's t test was used. Differences were considered significant when $p < 0.05$.

3. Results and Discussion



3.1. Total polyphenol content (TPC) and flavonoid contents (TFC) of OFE and FFE

The total phenol and flavonoid content of OFE and FFE extracts is shown in Table 1. OFE (174.7 ± 6.0 mg EGA/100 g extract) was proven to have a higher total polyphenol content when compared to FFE (83.2 ± 4.4 mg EGA/100 g extract); similarly, the total flavonoid content of OFE (59.9 ± 2.9 mg EQ/100 g extract) was higher than that of FFE (27.6 ± 1.2 mg EQ/100 g extract). The TPC of OFE reported in the present work (174.7 mg EGA/100 g) agrees with a previous determination of TPC in olive fruits of seven different cultivars of *Olea europaea*, collected in Italy (*Coratina*, *Frantoio*, *Leccino*, *Maiatica*, and *Ogliarola* cultivars) and Algeria (*Sigoise* and *Chemlal* cultivars), which ranged from 147.1 to 290.2 EGA/100 g. However, the TPC was slightly lower than the previously determined in the same olive fruit variety (272.8 EGA/100 g) (Dekdouk et al., 2015). This value is also similar to those observed in the *Chetoui* variety of olive fruits, from Tunisia, which were 313.6, 309.8, 36.1 and 83.3 mg hydroxytyrosol equivalents/100 g during the extraction with hexane, dichloromethane, ethyl acetate and methanol, respectively, although a different standard (hydroxytyrosol equivalents) was used (Brahmi et al., 2015). This variation in TPC may be related to the solvent system used (Brahmi et al., 2015) and the varieties of olive fruits (Dekdouk et al., 2015).

In the case of figs, the TPC content was 83.2 mg EGA/100 g extract, which is lower than that reported by Khadhraoui et al. (2019) for dark dried fig of *Saoudi Douiret* cultivar, but similar to that of green dried fig of *Bayoudhi Douiret* cultivar from Tunisia (TPC of 201.8 and 73.7 mg EGA/100 g, respectively). This value is also lower than those reported for dark colored (*Aberkane*, *Azandjar* and *Bouankik* varieties) and light colored (*Abiarous*, *Azegzaw*, *El-bakour*, *Taamriwth*, *Tahyounte* and *Taghanimt* varieties) dried figs from northern Algeria: the median for dark and light varieties of figs were 619 and 515 mg EGA/100g, respectively (Bachir Bey and Louaileche, 2015). Moreover, they are also lower than the TPC estimated for Iranian (1120-2682 mg EGA/100 g; Pourghayoumi et al., 2017), Indian (331.9 mg EGA/100 g; Vijaya Kumar Reddy et al. (2010) and Turkish (169.4 mg EGA/100 g; dried figs (Capanoglu, 2014). These differences in TPC could be due to many parameters such as the geographical origin, varieties, extraction conditions, and postharvest storage conditions (Bachir Bey et al., 2013; Hoxha et al., 2015).

Flavonoids, a class of polyphenolic secondary metabolites, are one of the most diverse and widespread groups of phytochemicals. These compounds constitute a potential therapeutic alternative for various diseases (Mutha et al., 2021). We verified that the TFC of OFE and FFE were 59.9 ± 2.9 and 27.6 ± 1.2 mg EQ/100 g, respectively. The flavonoid content of olive and Fig fruits are well documented in the literature. The TFC values of OFE obtained is lower than that previously reported for olive extracts of the *Chetoui* variety, from Tunisia, which were 844.6 and 131.2 mg catechin equivalents (EC)/100 g, by dichloromethane and hexane extraction, respectively (Brahmi et al. 2015). Similarly, the TFC of FFE is also slightly lower than those obtained by Bachir

Bey and Louaileche (2015), which indicated TFC of dark and light dried fig varieties of 126.6 mg EQ/100 g and 87.2 mg EQ/100 g, respectively. Khadhraoui et al. (2019) reported that the TFC of dried fig extracts from South-Eastern and Middle Eastern Tunisia ranged from 58.0 mg to 112.3 mg EQ/100 g. The variation of TPC could be due to differences in Fig varieties/genotype (Caliskan and Polat, 2012), drying methods (Khadhraoui et al., 2019), solvent used and could also be related to different unit used (catechin or quercetin equivalents).

3.2. Antioxidant activity of OFE and FFE

The antioxidant activity of OFE and FFE was determined using different *in vitro* methods: DPPH[•] free radical scavenging assay, ABTS^{•+} scavenging assay, CUPRAC assay and the Reducing Power assay. The results of all the tests, expressed in terms of IC₅₀ (µg/mL) or A_{0.5}, are presented in Table 2. The effectiveness of antioxidant properties is inversely correlated with their A_{0.5} and IC₅₀ values.

DPPH[•] or ABTS^{•+} can accept an electron or a hydrogen atom from the antioxidant compound and will thus be converted to a more stable molecule; this reduction is observed as a color switch for DPPH[•] (from purple to yellow) and as a decolorization for the radical ABTS^{•+} (Re et al., 1999; Magalhães et al., 2006). Table 2 showed that OFE was more effective in DPPH[•] and ABTS^{•+} scavenging assays (IC₅₀ value of 29.6 ± 0.8 and 32.4 ± 0.5 µg/mL, respectively) than FFE (IC₅₀ > 100 µg/mL in both assays). In the Reducing power assay, the presence of reductants converts the oxidized form of iron (Fe³⁺) in ferric chloride to ferrous (Fe²⁺), which can be monitored by absorbance measurement at 700 nm (Oyaizu, 1986). The CUPRAC assay is based on the reduction of the neocuproine-copper complex, resulting in the formation of a chromogenic complex of copper-neocuproine (Cu(II)-Nc), which absorbs at 450 nm (Saci et al., 2020). The antioxidant capacity of the extracts evaluated with the CUPRAC and the Reducing power assays is presented as A_{0.5} values, which were determined from absorbance curves at the corresponding wavelength (700 and 450 nm, respectively). This reduction was found to be concentration-dependent (results not shown). OFE exhibited a strong ability to reduce iron (A_{0.5} value: 129.8 ± 3.7 µg/mL) and a weaker ability to reduce copper (A_{0.5} value: 29.8 ± 0.9 µg/mL). FFE showed a lower capacity to reduce both iron and copper (A_{0.5} values were > 200 µg/mL and > 100 µg/mL, respectively) (Table 2).

As expected, BHA and BHT, the standard references for the DPPH[•], ABTS^{•+} and CUPRAC assays, showed a strong scavenger and reducing ability. Ascorbic acid and tannic acids, which were used as standards for the Reducing Power assay, showed also strong reducing activities (Table 2).

Due to the variable responses exerted by a specific antioxidant in various testing systems, it is important to use diverse antioxidant methods to take in account the mechanism of action of each compound (Moukette et al., 2015). The potent antioxidant activity of polyphenols and flavonoids resumes in their ability to act as reducing agents, hydrogen donors and singlet oxygen quenchers (Kasote et al., 2015). The methanolic extract conclusion,

Table 1

Total phenolic content (TPC) and total flavonoid content (TFC) of *Olea europaea* (OFE) and *Ficus carica* (FFE) fruit extracts. Total phenolics content is expressed as mg of gallic acid equivalents per 100 mg of extract (mg EGA/100 mg) and total flavonoid content as mg quercetin equivalents per 100 mg of extract (mg EQ/100 g)^a.

Extract	TPC (mg EGA/ 100 g extract)	TFC (mg EQ/100 g extract)
OFE	174.74 ± 6.03	59.88 ± 2.91
FFE	83.19 ± 4.42	27.64 ± 1.22

^a Results are expressed as arithmetic means ± SD of three measurements.

Table 2

Antioxidant activity of *Olea europaea* (OFE) and *Ficus carica* (FFE) fruit extracts. Results are expressed as IC₅₀ (µg/mL) (DPPH[•] and ABTS^{•+} assays) or A_{0.5} (µg/mL) (CUPRAC and reducing power assays).

Sample	DPPH [•] (IC ₅₀)	ABTS ^{•+} (IC ₅₀)	CUPRAC (A _{0.5})	Reducing Power (A _{0.5})
OFE	29.55 ± 0.77	32.40 ± 0.45	29.83 ± 0.91	129.84 ± 3.72
FFE	>100	>100	>100	>200
BHT	22.32 ± 1.19	1.29 ± 0.30	9.62 ± 0.87	nt
BHA	5.73 ± 0.41	1.81 ± 0.10	3.64 ± 0.19	nt
Ascorbic acid	nt	nt	nt	6.77 ± 1.15
Tannic acid	nt	nt	nt	5.39 ± 0.91

The A_{0.5} (Reducing power and CUPRAC assays) and the IC₅₀ values (DPPH[•] and ABTS^{•+} assays) were calculated by linear regression analysis. Results are expressed as arithmetic means ± SD of three measurements. BHA: Butylated hydroxyl anisole, BHT: Butylhydroxytoluene. nt: Not tested.




Fig. 1. Effects of *Olea europaea* (OFE) and/or *Ficus carica* (FFE) fruit extracts (24 h) on the viability of PANC-1 and AsPC-1 cells. (A-B) Effect of distinct concentrations of OFE and FFE (n=6); (C-D) Effect of the combination of OFE (0.5 mg/mL) and FFE (0.5 mg/mL) (n=8). Results are presented as arithmetic means ± SEM of two measurements. (A-B) * $p < 0.05$ (Student's t test); (C-D) * $p < 0.05$ vs control and # $p < 0.05$ vs each other (ANOVA followed by Student-Newman-Keuls test).




Fig. 2. Effects of *Olea europaea* (OFE) and/or *Ficus carica* (FFE) fruit extracts (24 h) on proliferation of PANC-1 and AsPC-1 cells. (A-B) Effect of distinct concentrations of OFE and FFE (n=4); (C-D) Effect of combination of OFE (0.5 mg/mL) and FFE (0.5 mg/mL) (n=6). Results are presented as arithmetic means \pm SEM of two measurements. (A-B) * $p < 0.05$ (Student's t test); (C-D) * $p < 0.05$ vs control and # $p < 0.05$ vs each other (ANOVA followed by Student-Newman-Keuls test).

of olive fruit (OFE) was found to exhibit a more potent activity as a scavenger of free radicals and as a reducing agent than Fig extract (FFE). Olive fruits are known to have a potent antioxidant activity (Brahmi et al., 2015; Dekdouk et al., 2015), and the major contributors to their antioxidant activity appear to be hydroxytyrosol, tyrosol, *p*-hydroxybenzoic acid, and verbascoside (Dekdouk et al., 2015). So, the more potent antioxidant activity of OFE is most probably correlated with its higher phenolic content (Dekdouk et al., 2015; Brahmi et al., 2015).

3.3. Anticancer activity

3.3.1. Effects of OFE and/or FFE on cell viability

As shown in Fig. 1, OFE and FFE showed promising cytotoxicity against PANC-1 and AsPC-1 cells (Fig. 1A-B). Combination of the extracts clearly increase the cytotoxic effect against both pancreatic cell lines (Fig. 1C-D).

3.3.2. Effect of OFE and/or FFE on cell proliferation

OFE exhibited a marked concentration-dependent antiproliferative effect in both cell lines. In contrast, FFE caused no significant changes in proliferation rates (Fig. 2A-B). When the two extracts were combined, the inhibitory effect was equal to that caused by OFE, in both cell lines (Fig. 2C-D).

A general consensus exists on the importance of dietary

habits, including the quantity and quality of food ingested, and their impact on health. Epidemiological studies have clearly established a protective effect of the Mediterranean diet against several types of cancer (Schwingshackl and Hoffmann, 2014; Saha et al., 2017; Mentella et al., 2019; Guasch-Ferré and Willett, 2021), including colorectal (Bamia et al., 2013), gastric (Buckland et al., 2010) and breast (Buckland et al., 2013) cancers.

A positive relationship between olive oil consumption and a reduction in cancer risk has been reported (Romani et al., 2019; Markellos et al., 2022). In agreement with this, our study revealed that OFE possesses a concentration-dependent cytotoxic and antiproliferative effect in both AsPC-1 and PANC-1 pancreatic cancer cell lines. These findings agree with previous reports showing a cytotoxic and/or antiproliferative effect of OFE against other cancer lines, namely hepatic (HepG2) and colorectal (Caco-2) (Maalej et al., 2017; Celano et al., 2019) and a protective effect in a rat model of hepatocellular carcinoma (Shirazi et al., 2018). Moreover, an antiproliferative effect of extracts from *Corregiola* and *Frantoio* varieties of *Olea europaea* L. leaves were reported in the MIA PaCa-2 pancreatic cancer cell line (Goldsmith et al., 2015). The underlying mechanisms by which olive oil exerts a cancer preventive effect have been attributed to its content in various bioactive natural products, specifically polyphenols. Indeed, polyphenols are known to possess several anticancer mechanisms, namely antioxidant, anti-inflammatory, anti-mutagenic, anti-proliferative, pro-apoptotic and anti-angiogenic

(Gorzynik-Debicka et al., 2018). In support of this the main phenolic compounds present in olive extracts (tyrosol, hydroxytyrosol, oleocanthal, oleuropein, rutin, quercetin and glucoside forms of luteolin) have been shown to have anticancer effects on several types of cancers. For instance, hydroxytyrosol induces apoptosis of the LS180 human colorectal carcinoma cell line (Hormozi et al., 2020). Additionally, oleocanthal has antiproliferative and cell cycle inhibitory effects in hepatocarcinoma cells *in vitro* and suppresses tumor growth in an orthotopic hepatocarcinoma model (Pei et al., 2016). It also inhibits invasiveness and tumor growth in breast cancer models (Akl et al., 2014). As to oleuropein, it decreases the number of breast cancer MCF-7 cells by inhibiting the rate of proliferation and inducing cell apoptosis (Han et al., 2009). Importantly, oleocanthal was reported to induce death of the BxPC-3 pancreatic cancer cell line (LeGendre et al., 2015) and Goldsmith et al. (2018) reported a potential antitumoral effect of oleuropein and hydroxytyrosol through reduction of proliferation, influence of cell cycle and induction of apoptosis in MIA PaCa-2 pancreatic cancer cells. However, these compounds showed no effect on other pancreatic cancer cell lines (BxPC-3 and CFPAC-1) (Goldsmith et al., 2018).

Our current study evidenced that FFE possesses a cytotoxic effect against both AsPC-1 and PANC-1 pancreatic cancer cell lines, although no significant antiproliferative effect was verified. The methanolic extract of *Ficus carica* leaves and fruits was previously shown to possess cytotoxic, pronecrotic and proapoptotic effects in Huh7it cancer liver cells (Purnamasari et al., 2019). Very interestingly, a recent study also showed that a *F. carica* fruit extract reduced viability and effectively suppressed migration, metastasis, invasion, and colony formation of PANC-1 and QGP-1 pancreatic cancer cell lines, by inducing apoptotic cell death and autophagy (Ou et al., 2022). However, in contrast to the present study, antiproliferative effects of fig extracts were observed in other cancer cell types. Namely, *F. carica* leaves extract induced anti-growth effects on triple-negative breast cancer MDA-MB-231 cells by promoting apoptosis and cell-cycle arrest at the S phase (Zhang et al., 2018), an antiproliferative effect of fig tree latex was observed on a stomach cancer cell line (Hashemi et al., 2011), and an antiproliferative and proapoptotic effect of fig leaves, whole fruit and latex extracts was observed on HCT-116 and HT-29 colorectal cancer cell lines (Soltana et al., 2019).

A synergic effect has been widely investigated between polyphenols and therapeutical drugs. For instance, quercetin promoted cisplatin-induced apoptosis *in vitro* in human oral squamous cell carcinoma cell lines by downregulating NF- κ B and promoted the apoptotic effect of cisplatin *in vivo* in a mouse xenograph model (Li et al., 2019). Moreover, this same combination had a synergic inhibitory effect on cervical cancer cells via inhibition of proliferation, migration, invasion, and stimulation of apoptosis (Xu et al., 2021). In the current study, we were interested in verifying the effect of combination of the two extracts against AsPC-1 and PANC-1 cell lines. Our findings show a potent additive cytotoxic and antiproliferative effect of combination

of OFE and FFE against these two cell lines. This result may be explained by the additive effect of phenolic compounds present in the two extracts, as previously reported when combining quercetin and sulforaphane (Srivastava et al., 2011), isoflavone and curcumin (Wang et al., 2008) and ferulic acid and δ -tocotrienol (Eitsuka et al., 2014).

3.3.3. Effect of OFE and/or FFE on cell migration

As shown in Fig. 3, OFE evidenced a concentration-dependent antimigratory effect in PANC-1 cells, but no effect on AsPC-1 migration rates was found. As for FFE, it exhibited a marked and concentration-dependent stimulatory effect on the migration rates of both cell lines (Fig. 3A-B). More interestingly, combination of the two extracts led to a complete abolition of the stimulatory effect of FFE in the two cell lines; instead, an effect very similar to that caused by OFE was observed (Fig. 3C-D).

Directional cell migration plays a key role in pivotal steps that promote tumor metastasis, such as cellular migration and invasion into the surrounding stroma. Given the fact that current chemotherapy drugs for pancreatic cancer only offer an increase in survival of up to six months (Moore et al., 2007; Park et al., 2021), the search for novel agents with an antimigratory effect on pancreatic cancer cells while displaying limited toxicity towards normal cells is thus paramount. The present data clearly demonstrate that:

- (1) OFE showed a concentration-dependent antimigratory effect towards PANC-1 cells only.
- (2) OFE presented a marked stimulatory effect on the migration of both cell lines.
- (3) Very interestingly, combination of the two extracts resulted in a complete abolition of the stimulatory effect of FFE and instead an effect comparable to that caused by OFE was found. Although we have no evidence as to the mechanism involved in this effect of OFE, components in OFE appear to be able to abolish the stimulatory effect of FFE on cell migration.

Several studies have tested the ability of crude extracts rich in phenolic compounds or isolated phenolic compounds in inhibiting migration of pancreatic cancer cells *in vitro* and *in vivo*. Fahrioglu et al. (2016) reported that ferulic acid exhibits an anticancer effect by inhibiting migration of MIA PaCa-2 pancreatic cancer cells. Similarly, kaempferol effectively inhibited the migratory activity of MIA PaCa-2, PANC-1, and SNU-213 human pancreatic cancer cell lines at relatively low concentrations (Lee and Kim, 2016). From our findings, we believe that the use of OFE, either alone or in combination with pro-migratory agents, can inhibit pancreatic cell migration, which is very interesting.

4. Concluding remarks

Collectively, this report demonstrates that OFE, which is very rich in bioactive polyphenolic compounds, possesses anticancer effects against PANC-1 and AsPC-1 pancreatic cancer cell lines. It exhibits cytotoxic, antiproliferative and antimigratory effects. In contrast, FFE exhibits a cytotoxic effect in these two cell lines,




Fig. 3. Effects of *Olea europaea* (OFE) and/or *Ficus carica* (FFE) fruit extracts (24 h) on the migration capacity of PANC-1 and AsPC-1 cells. (A-B) Effect of distinct concentrations of OFE and FFE (n=6); (C-D) Effect of combination of OFE (0.5 mg/mL) and FFE (0.5 mg/mL) (n=8). Results are presented as arithmetic means \pm SEM of two measurements. (A-B) * $p < 0.05$ (Student's t test); (C-D) * $p < 0.05$ vs control and # $p < 0.05$ vs each other (ANOVA followed by Student-Newman-Keuls test).

but it does not have an antiproliferative effect and even stimulates the migration of these cells. The combination of OFE and FFE led to an increase in the cytotoxic effect in both cell lines and the disappearance of the promigratory effect of FFE. Thus, the combination of these extracts in fact exerts a more interesting anticancer effect than either of the extracts alone.

This report also demonstrates that OFE and FFE exhibit different antioxidant properties in response to different models, namely DPPH⁺, ABTS⁺, CUPRAC and Reducing Power models, which may be related to their different polyphenol and flavonoid content. More importantly, the higher phenolic and flavonoid content and antioxidant ability of OFE, in relation to FFE, correlates with a more prominent anticancer effect. Because the biological activity of polyphenols is strongly related to their antioxidant properties (Gorzynik-Debicka et al., 2018), we suggest that the anticancer effect of the extracts results from their antioxidant properties.

This study could lead to new research on pancreatic cancer and the consumption of olive and fig products, and place OFE and FFE/ phenolic compounds as good candidates for further molecular studies aimed at finding a better, and more effective treatment approach for resistant pancreatic cancer. In future work, it would be very interesting to confirm the results of this study in other human pancreatic cancer cell lines, and also in

animal pre-clinical models. Moreover, the mechanisms involved in the positive interaction between OFE and FFE should be further explored.

List of abbreviations

ABTS: 2,2-Azinobis-3-ethylbenzothiazoline-6-sulfonic acid; **AsPC-1 Cell Line:** Human Pancreatic Ascites Adenocarcinoma Cell Line; **ATCC:** American Type Culture Collection; **BHA:** 2,6-Di-*tert*-butyl-4-hydroxy toluene; **BHT:** Butylated Hydroxytoluene; **CUPRAC:** Cupric Acid Reducing Antioxidant Capacity; **DMSO:** Dimethylsulfoxide; **DPPH⁺:** (1,1-Diphenyl-2-picrylhydrazil) Scavenging Assay; **FFE:** *Ficus carica* Extract; **HEPES:** *N*-2-Hydroxyethylpiperazine-*N'*-2-ethanesulfonic Acid; **MTT:** 3-(4,5-Dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide; **OFE:** *Olea europaea* Extract; **PANC-1 Cell Line:** Human Pancreatic Epithelioid Carcinoma Cell Line; **SD:** Standard Deviation; **SEM:** Standard Error of the Mean; **TFC:** Total Flavonoid Content; **TPC:** Total Phenolic Content.

Author contribution statement

All authors contributed to the study's conception and

design. Nadia Dekdouk, Souad Ameddah, Chawki Bensouici, Meghboun Ibtissem and Menad Ahmed performed material preparation, experiments, data collection, and analysis. Nadia Dekdouk and Fátima Martel wrote the first draft of the manuscript, and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Conflicts of interest

The authors declare that there is no conflict of interest.

Acknowledgments

Nadia Dekdouk thanks very much the Department de Microbiologie et Biochimie-Faculté des Sciences de la Nature et de la Vie-Université Chahid Mostafa Benboulaïd-Batna2, Algérie, for the internship granted to the University of Porto, Porto, Portugal.

References

Akl, M.R., Ayoub, N.M., Mohyeldin, M.M., Busnena, B.A., Foudah, A.I., Liu, Y.Y., Sayed, K.A.E., 2014. Olive phenolics as c-Met inhibitors:(-)-Oleocanthal attenuates cell proliferation, invasiveness, and tumor growth in breast cancer models. *PLoS One* 9, e97622.

Al-Salman, H.N.K., Ali, E.T., Jabir, M., Sulaiman, G.M., Al-Jadaan, S.A.S., 2020. 2-Benzhydrylsulfonyl-N-hydroxyacetamide-Na extracted from fig as a novel cytotoxic and apoptosis inducer in SKOV-3 and AMJ-13 cell lines viaP53 and caspase-8 pathway. *Eur. Food Res. Technol.* 246, 1591-1608.

Apak, R., Güçlü, K., Özyürek, M., Çelik, S.E., 2008. Mechanism of antioxidant capacity assays and the CUPRAC (cupric ion reducing antioxidant capacity) assay. *Microchim. Acta* 160, 413-419.

Arvaniti, O.S., Samaras, Y., Gatidou, G., Thomaidis, N.S., Stasinakis, A.S., 2019. Review on fresh and dried figs: Chemical analysis and occurrence of phytochemical compounds, antioxidant capacity and health effects. *Food Res. Int.* 119, 244-267.

Baby, J., Justin, S.R., 2011. Pharmacognostic and phytochemical properties of *Ficus carica*. *Linn. Int. J. Pharm. Tech. Res.* 3, 8-12.

Bachir Bey, M., Louaileche, H., 2015. A comparative study of phytochemical profile and *in vitro* antioxidant activities of dark and light dried fig (*Ficus carica* L.) varieties. *J. Phytopharmacol.* 4, 41-48.

Baldo, B.A., Pham, N.H., 2013. Adverse reactions to targeted and non-targeted chemotherapeutic drugs with emphasis on hypersensitivity responses and the invasive metastatic switch. *Cancer Metastasis Rev.* 32, 723-761.

Bamia, C., Lagiou, P., Buckland, G., Grioni, S., Agnoli, C., Taylor, A.J., Dahm, C.C., Overvad, K., Olsen, A., Tjønneland, A., Cottet, V., Boutron-Ruault, M.-C., Morois, S., Grote, V., Teucher, B., Boeing, H., Buijssse, B., Trichopoulos, D., Adarakis, G., Tumino, R., Naccarati, A., Panico, S., Palli, D., Bueno-de-Mesquita, H.B., van Duijnhoven, F.J.B., Peeters, P.H.M., Engeset, D., Skeie, G., Lund, E., Sánchez, M.-J., Barricarte, A., Huerta, J.-M., Ramón Quirós, J.,

Dorronsoro, M., Ljuslinder, I., Palmqvist, R., Drake, I., Key, T.J., Khaw, K.-T., Wareham, N., Romieu, I., Fedirko, V., Jenab, M., Romaguera, D., Norat, T., Trichopoulou, A., 2013. Mediterranean diet and colorectal cancer risk: results from a European cohort. *Eur. J. Epidemiol.* 28, 317-328.

Bar-Zeev, M., Livney, Y.D., Assaraf, Y.G., 2017. Targeted nanomedicine for cancer therapeutics: Towards precision medicine overcoming drug resistance. *Drug Resist. Updat.* 31, 15-30.

Basha, R., Connelly, S.F., Sankpala, U.T., Purnachandra, N.G., Patel, H., Vishwanathaa, J.K., Shelakea, S., Tabor-Simecka, L., Shoji, M., Simecka, J.W., El-Rayes, B., 2016. Small molecule tolfenamic acid and dietary spice curcumin treatment enhances antiproliferative effect in pancreatic cancer cells via suppressing Sp1, disrupting NF- κ B translocation to nucleus and cell cycle phase distribution. *J. Nutr. Biochem.* 31, 77-87.

Blois, M.S., 1958. Antioxidants determination by the use of a stable free radical. *Nature.* 26, 1199-1200.

Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72, 248-254.

Brahmi, F., Mechri, B., Dhibi, M., Hammami, M., 2015. Antioxidant capacity and phenolic content of olive fruits from Tunisia, extracted with different solvents. *J. Plant Sci.* 3, 8-12.

Buckland, G., Travier, N., Cottet, V., González, C.A., Luján-Barroso, L., Agudo, A., Trichopoulou, A., Lagiou, P., Trichopoulos, D., Peeters, P.H., May, A., Bueno-de-Mesquita, H.B., Bvan Duijnhoven, F.J., Key, T.J., Allen, N., Khaw, K.T., Wareham, N., Romieu, I., McCormack, V., Boutron-Ruault, M., Clavel-Chapelon, F., Panico, S., Agnoli, C., Palli, D., Tumino, R., Vineis, P., Amiano, P., Barricarte, A., Rodríguez, L., Sanchez, M.J., Chirlaque, M.D., Kaaks, R., Teucher, B., Boeing, H., Bergmann, M.M., Overvad, K., Dahm, C.C., Tjønneland, A., Olsen, A., Manjer, J., Wirfält, E., Hallmans, G., Johansson, I., Lund, E., Hjartaker, A., Skeie, G., Vergnaud, A.C., Norat, T., Romaguera, D., Riboli, E., 2013. Adherence to the mediterranean diet and risk of breast cancer in the European prospective investigation into cancer and nutrition cohort study. *Int. J. Cancer* 132, 2918-2927.

Buckland, G., Agudo, A., Luján, L., Jakszyn, P., Bueno-de-Mesquita, H.B., Palli, D., Boeing, H., Carneiro, F., Krogh, V., Sacerdote, C., Tumino, R., Panico, S., Nesi, G., Manjer, J., Regnér, S., Johansson, I., Stenling, R., Sanchez, M.-J., Dorronsoro, M., Barricarte, A., Navarro, C., Quirós, J.R., Allen, N.E., Key, T.J., Bingham, S., Kaaks, R., Overvad, K., Jensen, M., Olsen, A., Tjønneland, A., Peeters, P.H.M., Numans, M.E., Ocké, M.C., Clavel-Chapelon, F., Morois, S., Boutron-Ruault, M.-C., Trichopoulou, A., Lagiou, P., Trichopoulos, D., Lund, E., Couto, E., Boffeta, P., Jenab, M., Riboli, E., Romaguera, D., Mouw, T., González, C.A., 2010. Adherence to a mediterranean diet and risk of gastric adenocarcinoma within the European Prospective Investigation into Cancer and Nutrition (EPIC) cohort study. *Am. J. Clin. Nutr.* 91, 381-390.

Caliskan, O., Polat, A.A., 2012. Effects of genotype and harvest year on phytochemical and fruit quality properties of Turkish fig genotypes. *Span. J. Agric. Res.* 10, 1048-1058.



- Capanoglu, E., 2014. Investigating the antioxidant potential of Turkish dried fruits. *Int. J. Food Prop.* 17, 690-70.
- Carrera-González, M.P., Ramírez-Expósito, M.J., Mayas, M.D., Martínez-Martos, J.M., 2013. Protective role of oleuropein and its metabolite hydroxytyrosol on cancer. *Trends Food Sci. Technol.* 31, 92-99.
- Celano, M., Maggisano, V., Lepore, S.M., Russo, D., Bulotta, S., 2019. Secoiridoids of olive and derivatives as potential adjuvant drugs in cancer: A critical analysis of experimental studies. *Pharmacol. Res.* 142, 77-86.
- Cruz, J.M.A., Corrêa, R.F., Lamarão, C.V., Kinupp, V.F., Sanches, E.A., Campelo, P.H., Bezerra, J.A., 2022. *Ficus* spp. fruits: Bioactive compounds and chemical, biological and pharmacological properties. *Food Res. Int.* 152, 110928.
- Dekdouk, N., Malafrente, N., Russo, D., Faraone, I., De Tommasi, N., Ameddah, S., Severino, L., Milella, L., 2015. Phenolic compounds from *Olea europaea* L. possess antioxidant activity and inhibit carbohydrate metabolizing enzymes *in vitro*. *Evid. Based Complementary Altern. Med.* 2015, 684925.
- Dhar, D., Raina, K., Agarwal, R., 2018. Mechanisms and drug targets for pancreatic cancer chemoprevention. *Curr. Med. Chem.* 25, 2545-2565.
- Eitsuka, T., Tatewaki, N., Nishida, H., Kurata, T., Nakagawa, K., Miyazawa, T., 2014. Synergistic inhibition of cancer cell proliferation with a combination of δ -tocotrienol and ferulic acid. *Biochem. Biophys. Res. Commun.* 453, 606-11.
- Fabiani, R., De Bartolomeo, A., Rosignoli, P., Servili, M., Montedoro, G.F., Morozzi, G., 2002. Cancer chemoprevention by hydroxytyrosol isolated from virgin olive oil through G1 cell cycle arrest and apoptosis. *Eur. J. Cancer Prev.* 11, 351-358.
- Fahrioglu, U., Dodurga, Y., Elmas, L., Seçme, M., 2016. Ferulic acid decreases cell viability and colony formation while inhibiting migration of MIA PaCa-2 human pancreatic cancer cells *in vitro*. *Gene* 576, 476-82.
- Fki, I., Bouaziz, M., Sahnoun, Z., Sayadi, S., 2005. Hypocholesterolemic effects of phenolic-rich extracts of Chemlali olive cultivar in rats fed a cholesterol-rich diet. *Bioorg. Med. Chem.* 13, 5362-70.
- Ghandehari, F., Fatemi, M., Avicenna, J., 2018. The effect of *Ficus carica* latex on 7,12-dimethylbenz (a) anthracene-induced breast cancer in rats. *Avicenna Phytomed.* 8, 286-295.
- Gianfredi, V., Ferrara, P., Dinu, M., Nardi, M., Nucci, D., 2022. Diets, dietary patterns, single foods and pancreatic cancer risk: an umbrella review of meta-analyses. *Int. J. Environ. Res. Public Health* 19, 14787.
- Goldsmith, C.D., Bond, D.R., Jankowski, H., Weidenhofer, J., Stathopoulos, C.E., Roach, P.D., Scarlett, C.J., 2018. The olive biophenols oleuropein and hydroxytyrosol selectively reduce proliferation, influence the cell cycle, and induce apoptosis in pancreatic cancer cells. *Int. J. Mol. Sci.* 19, 1937.
- Goldsmith, C.D., Vuong, Q.V., Sadeqzadeh, E., Stathopoulos, C.E., Roach, P.D., Scarlett, C.J., 2015. Phytochemical properties and anti-proliferative activity of *Olea europaea* L. leaf extracts against pancreatic cancer cells. *Molecules* 20, 12992-13004.
- Gorzynik-Debicka, M., Paulina Przychodzen, P., Cappello, F., Kuban-Jankowska, A., Gammazza, A.M., Knap, N., Wozniak, M., Gorska-Ponikowska, M., 2018. Potential health benefits of olive oil and plant polyphenols. *Int. J. Mol. Sci.* 19, 686.
- Greenwell, M., Rahman, P.K.S.M., 2015. Medicinal plants: Their use in anticancer treatment. *Int. J. Pharm. Sci. Res.* 6, 4103.
- Guarrera, P.M., 2005. Traditional phytotherapy in central Italy (Marche, Abruzzo, and Latium). *Fitoterapia* 76, 1-25.
- Guasch-Ferré, M., Willett, W.C., 2021. The Mediterranean diet and health: a comprehensive overview. *J. Intern. Med.* 290, 549-566.
- Guo, Z., Jia, X., Zheng, Z., Lu, X., Zheng, Y., Zheng, B., Xiao, J., 2018. Chemical composition and nutritional function of olive (*Olea europaea* L.): A review. *Phytochem. Rev.* 17, 1091-1110.
- Han, J., Talorete, T.P., Yamada, P., Isoda, H., 2009. Anti-proliferative and apoptotic effects of oleuropein and hydroxytyrosol on human breast cancer MCF-7 cells. *Cytotechnology* 59, 45-53.
- Hashemi, S.A., Abediankenari, S., Ghasemi, M., Azadbakht, M., Yousefzadeh, Y., Dehpour, A.A., 2011. The effect of fig tree latex (*Ficus carica*) on stomach cancer line. *Iran Red Crescent Med. J.* 13, 272-5.
- Hormozi, M., Marzijerani, A.S., Baharvand, P., 2020. Effects of hydroxytyrosol on expression of apoptotic genes and activity of antioxidant enzymes in LS180 cells. *Cancer Manag. Res.* 12, 7913-7919.
- Ijoma, K.I., Ajiwe, V.I.E., Odinma, S.C., 2023. The organic extracts from the leaves of *Ficus thonningii* Blume, *Jatropha tanjorensis* J.L Ellis and *Justicia carnea* Lindley as potential nutraceutical antioxidants and functional foods. *Trends Phytochem. Res.* 7(1), 76-85.
- Kasote, D.M., Katyare, S.S., Hegde, M.V., Bae, H., 2015. Significance of antioxidant potential of plants and its relevance to therapeutic applications. *Int. J. Biol. Sci.* 11, 982-991.
- Kazemina, M., Mehrabi, A., Mahmoudi, R., 2022. Chemical composition, biological activities, and nutritional application of Asteraceae family herbs: A systematic review. *Trends Phytochem. Res.* 6(3), 187-213.
- Khadhraoui, M., Bagues, M., Artés, F., Ferchichi, A., 2019. Phytochemical content, antioxidant potential, and fatty acid composition of dried Tunisian fig (*Ficus carica* L.) cultivars. *J. Appl. Bot. Food Qual.* 92, 143-150.
- Khan, T., Ali, M., Khan, A., Nisar, P., Jan, S.A., Afridi, S., Shinwari, Z.K., 2019. Anticancer plants: A review of the active phytochemicals, applications in animal models, and regulatory aspects. *Biomolecules* 10, 47.
- Kountouri, A.M., Kaliora, A.C., Koumbi, L., Andrikopoulos, N.K., 2009. *In-vitro* gastric cancer prevention by a polyphenol-rich extract from olives through induction of apoptosis. *Eur. J. Cancer Prev.* 18, 33-39.
- Le, K., Chiu, F., Ng, K., 2007. Identification and quantification of antioxidants in *Fructus lycii*. *Food Chem.* 105, 353-563.
- Lee, J., Kim, J.H., 2016. Kaempferol inhibits pancreatic cancer cell growth and migration through the blockade of EGFR-related pathway *in vitro*. *PLoS ONE* 11, e0155264.

- LeGendre, O., Breslin, P.A.S., Foster, D.A., 2015. (-)-Oleocanthal rapidly and selectively induces cancer cell death via lysosomal membrane permeabilization. *Mol. Cell. Oncol.* 2, e1006077.
- Li, X., Guo, S., Xiong, X.K., Peng, B.Y., Huang, J.M., Chen, M.F., Wang, F.Y., Wang, J.N., 2019. Combination of quercetin and cisplatin enhances apoptosis in OSCC cells by downregulating XIAP through the NF- κ B pathway. *J. Cancer* 10(19), 4509-4521.
- Lopez-Miranda, J., Perez-Jimenez, F., Ros, E., De Caterina, R., Badimón, L., Covas, M.I., Escrich, E., Ordovás, J.M., Soriguer, F., Abiá, R., Alarcón de la Lastra, C., Battino, M., Corella, D., Chamorro-Quirós, J., Delgado-Lista, J., Giugliano, D., Esposito, K., Estruch, R., Fernandez-Real, J.M., Gaforio, J.J., La Vecchia, C., Lairon, D., López-Segura, F., Mata, P., Menéndez, J.A., Muriana, F.J., Osada, J., Panagiotakos, D.B., Paniagua, J.A., Pérez-Martinez, P., Perona, J., Peinado, M.A., Pineda-Priego, M., Poulsen, H.E., Quiles, J.L., Ramírez-Tortosa, M.C., Ruano, J., Serra-Majem, L., Solá, R., Solanas, M., Solfrizzi, V., de la Torre-Fornell, R., Trichopoulou, A., Uceda, M., Villalba-Montoro, J.M., Villar-Ortiz, J.R., Visioli, F., Yiannakouris, N., 2010. Olive oil and health: Summary of the II international conference on olive oil and health consensus report, Jaen and Cordoba (Spain) 2008. *Nutr. Metab. Cardiovasc. Dis.* 20, 284-294.
- Maalej, A., Bouallagui, Z., Hadrich, F., Isoda, H., Sayadi, S., 2017. Assessment of *Olea europaea* L. fruit extracts: Phytochemical characterization and anticancer pathway investigation. *Biomed. Pharmacother.* 90, 179-186.
- Magalhães, L.M., Santos, M., Segundo, M.A., Reis, S., Lima, J.L.F.C., 2006. Automatic method for determination of total antioxidant capacity using 2,2-diphenyl-1-picrylhydrazyl assay. *Anal. Chim. Acta.* 558, 310-318.
- Mahdavi, B., Mohammadhosseini, M., 2022. Antioxidant, antimicrobial and anti-prostate cancer activity of the extracts from different parts of *Etlingera velutina* (Ridl.) R. M. Sm (Zingiberaceae). *Trends Phytochem. Res.* 6(4), 353-362.
- Markellos, C., Ourailidou, M.E., Gavriatopoulou, M., Halvatsiotis, P., Sergentanis, T.N., Psaltopoulou, T., 2022. Olive oil intake and cancer risk: a systematic review and meta-analysis. *PLoS One* 17, e0261649.
- Mentella, M.C., Scaldaferrri, F., Ricci, C., Gasbarrini, A., Miggiano, G.A.D., 2019. Cancer and Mediterranean diet. *Nutrients.* 11, 2059.
- Mohammadhosseini, M., Frezza, C., Venditti, A., Akbarzadeh, A., 2019a. Ethnobotany and phytochemistry of the genus *Eremostachys* Bunge. *Curr. Org. Chem.* 23, 1828-1842.
- Mohammadhosseini, M., Frezza, C., Venditti, A., Sarker, S., 2021. A systematic review on phytochemistry, ethnobotany and biological activities of the genus *Bunium* L. *Chem. Biodivers.* 18(11), e2100317.
- Mohammadhosseini, M., Venditti, A., Akbarzadeh, A., 2019b. The genus *Perovskia* Kar.: Ethnobotany, chemotaxonomy and phytochemistry: A review. *Toxin Rev.* 40(4), 484-505.
- Moore, M.J., Goldstein, D., Hamm, J., Figer, A., Hecht, J.R., Gallinger, S., Au, H.J., Murawa, P., Walde, D., Wolff, R.A., Campos, D., Lim, R., Ding, K., Clark, G., Voskoglou-Nomikos, T., Ptasynski, M., Parulekar, W.; National Cancer Institute of Canada Clinical Trials Group., 2007. Erlotinib plus gemcitabine compared with gemcitabine alone in patients with advanced pancreatic cancer: a phase III trial of the National Cancer Institute of Canada Clinical Trials Group. *J. Clin. Oncol.* 25, 1960-1966.
- Morovati, M.R., Ghanbari-Movahed, M., Barton, E.M., Farzaei, M.H., Bishayee, A., 2022. A systematic review on potential anticancer activities of *Ficus carica* L. with focus on cellular and molecular mechanisms. *Phytomedicine* 105, 154333.
- Moukette, B.M., Pieme, C.A., Njimou, J.R., Biapa, C.P.N., Marco, B., Ngogang, J.Y., 2015. *In vitro* antioxidant properties, free radicals scavenging activities of extracts and polyphenol composition of a non-timber forest product used as spice: *Monodora myristica*. *Evidence-Based Complement. Alternat. Med.* 48, 1-17.
- Mutha, R.E., Tatiya, A.U., Surana, S.J., 2021. Flavonoids as natural phenolic compounds and their role in therapeutics. *Futur. J. Pharm. Sci.* 7, 25.
- Nguyen, L.T., Lee, Y.-H., Sharma, A.R., Park, J.-B., Jagga, S., Sharma, G., et al., 2017. Quercetin induces apoptosis and cell cycle arrest in triple-negative breast cancer cells through modulation of Foxo3a activity. *Korean J. Physiol. Pharmacol.* 21, 205-213.
- Obafemi, F.A., Besong, E.E., 2023. A review of plants with anticancer properties. *J. Clin. Med. Res.* 5, 138-161.
- Olaoluwa, O., Taiwo, O., Nahar, L., Sarker, S.D., 2022. Ethnopharmacology, phytochemistry and biological activities of selected African species of the genus *Ficus*. *Trends Phytochem. Res.* 6, 46-69.
- Ou, A., Zhao, X., Lu, Z., 2022. Autophagy is involved in *Ficus carica* fruit extract-induced anti-tumor effects on pancreatic cancer. *Biomed. Pharmacother.* 150, 112966.
- Owen, R.W., Giacosa, A., Hull, W.E., Haubner, R., Spiegelhalder, B., Bartsch, H., 2000a. The antioxidant/anticancer potential of phenolic compounds isolated from olive oil. *Eur. J. Cancer.* 36, 1235-1247.
- Owen, R.W., Giacosa, A., Hull, W.E., Haubner, R., Würtele, G., Spiegelhalder, B., Bartsch, H., 2000b. Olive oil consumption and health: The possible role of antioxidants. *Lancet Oncol.* 1, 107-112.
- Owen, R.W., Haubner, R., Würtele, G., Hull, E., Spiegelhalder, B., Bartsch, H., 2004. Olives and olive oil in cancer prevention. *Eur. J. Cancer Prev.* 13, 319-326.
- Oyaizu, M., 1986. Studies on products of browning reaction reactions: antioxidative activities of browning reaction prepared from glucosamine. *Jap. J. Nutr.* 44, 307-315.
- Park, W., Chawla, A., O'Reilly, E.M., 2021. Pancreatic cancer: a review. *JAMA* 326, 851-862.
- Pei, T., Meng, Q., Han, J., Sun, H., Li, L., Song, R., Sun, B., et al., 2016. (-)-Oleocanthal inhibits growth and metastasis by blocking activation of STAT3 in human hepatocellular carcinoma. *Oncotarget.* 7, 43475-43491.
- Popwo Tameye, S.C., Ndom, J.C., Nguemfo, E.L., Wansi, J.D., Vardamides, J.C., Azebaze, A.G.B., 2020. Ficusanol, a new cinnamic acid derivative and other constituents from the roots of *Ficus exasperata* Vahl. (Moraceae) with antioxidant and cytotoxic activities. *Trends Phytochem. Res.* 4, 3-8.
- Pourghayoumi, M., Bakhshi, D., Rahemi, M., Noroozisharaf, A., Jafari, M., Salehi, M., Chamane, R., Hernandez, F., 2017. Phytochemical attributes of some dried fig (*Ficus carica* L.) fruit cultivars grown in Iran. *Agric. Conspec. Sci.* 3, 161-166.



- Purnamasari, R., Winarni, D., Permasari, A.A., Agustina, E., Hayaza, S., Win, D., 2019. Anticancer activity of methanol extract of *Ficus carica* leaves and fruits against proliferation, apoptosis, and necrosis in Huh7it cells. *Cancer Inform.* 18, 1-7.
- Rauf, A., Imran, M., Khan, I.A., Ur-Rehman, M., Gilani, S.A., Mehmood, Z., Mubarak, M.S., 2018. Anticancer potential of quercetin: A comprehensive review. *Phytother. Res.* 32, 2109-2130.
- Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M., Rice-Evans, C., 1999. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Rad. Biol. Med.* 26, 1231-1237.
- Romani, A., Ieri, F., Urciuoli, S., Noce, A., Marrone, G., Nediani, C., Bernini, R., 2019. Health effects of phenolic compounds found in extra-virgin olive oil, by-products, and leaf of *Olea europaea* L. *Nutrients* 11, 1776.
- Saci, F., Bachir Bey, M., Louaileche, H., Gali, L., Bensouici, C., 2020. Changes in anticholinesterase, antioxidant and related bioactive compounds of carob pulp (*Ceratonia siliqua* L.) during ripening stage. *J. Food Measur. Character.* 14, 937-945.
- Saha, S.K., Lee, S.B., Won, J., Choi, H.Y., Kim, K., Yang, G.-M., Dayem, A.A., Cho, S.-G., 2017. Correlation between oxidative stress, nutrition, and cancer initiation. *Int. J. Mol. Sci.* 18, 1544.
- Schwingshackl, L., Hoffmann, G., 2014. Adherence to Mediterranean diet and risk of cancer: A systematic review and meta-analysis of observational studies. *Int. J. Cancer.* 135, 1884-1897.
- Selvarajoo, N., Stanslas, J., Islam, M.K., Sagineedu, S.R., Lian, H.K., Lim, J.C.W., 2022. Pharmacological modulation of apoptosis and autophagy in pancreatic cancer treatment. *Mini Rev. Med. Chem.* 22, 2581-2595.
- Sepporta, M.V., Fuccelli, R., Rosignoli, P., Ricci, G., Servili, M., Morozzi G., Fabiani, R., 2014. Oleuropein inhibits tumour growth and metastases dissemination in ovariectomised nude mice with MCF-7 human breast tumour xenografts. *J. Funct. Foods.* 8, 269-273.
- Shirazi, F.H., Piri, M., Keshavarz, S., Gholamib, S., Hosseini, S.H., Noubaranib, M., Andalibb, S., Kamalinejad, M., Adiban, H., Eskandari, M.R., 2018. Olive fruit (*Olea europaea* L.): Chemopreventive effect in the rat model of hepatocellular carcinoma. *PharmaNutrition* 6, 207-214.
- Siegel, R.L., Miller, K.D., Jemal, A., 2016. Cancer statistics. *CA Cancer J. Clin.* 66, 7-30.
- Singh, D., Upadhyay, G., Srivastava, R.K., Shankar, S., 2015. Recent advances in pancreatic cancer: Biology, treatment, and prevention. *Biochim. Biophys. Acta.* 1856, 13-27.
- Solomon, A.S., Golubowicz, Z., Yablowicz, S., Grossman, M., Bergman, H.E., Gottlieb, A., Altman, Z., Kerem, M.A., 2006. Antioxidant activities and anthocyanin content of fresh fruits of common fig (*Ficus carica* L.). *J. Agric. Food Chem.* 20, 7717-7723.
- Soltana, H., Pinon, A., Limami, Y., Zaid, Y., Khalki, L., Zaid, N., Salah, D., Sabitaliyevich, U.Y., Simon, A., Liagre, Bertrand., Hammami, M., 2019. Antitumoral activity of *Ficus carica* L. on colorectal cancer cell lines. *Cell. Mol. Biol.* 65, 6-11.
- Srivastava, R.K., Tang, S.N., Zhu, W., Meeker, D., Shankar, S., 2011. Sulforaphane synergizes with quercetin to inhibit self-renewal capacity of pancreatic cancer stem cells. *Front. Biosci.* 3, 515-28.
- Turkoglu, A., Duru, M.E., Mercan, N., Kivrak, I., Gezer, K., 2007. Antioxidant and antimicrobial activities of *Laetiporus sulphureus* (Bull.) Murrill. *Food Chem.* 101, 267-273.
- Vendrey, V., Peuchant, E., Buscail, E., Moranvillier, I., Rousseau, B., Bedel, A., Brillac, A., de Verneuil, H., Moreau-Gaudry, F., Dabernat, S., 2017. Resveratrol and capsaicin used together as food complements reduce tumor growth and rescue full efficiency of low dose gemcitabine in a pancreatic cancer model. *Cancer Lett.* 390, 91-102.
- Vijaya Kumar Reddy, C., Sreeramulu, D., Raghunath, M., 2010. Antioxidant activity of fresh and dry fruits commonly consumed in India. *Food Res. Int.* 43, 285-288.
- Wang, Z., Desmoulin, S., Banerjee, S., Kong, D., Li, Y., Deraniyagala, R.L., Abbruzzese, J., Sarkar, F.H., 2008. Synergistic effects of multiple natural products in pancreatic cancer cells. *Life Sci.* 83, 293-300.
- Xu, W., Xie, S., Chen, X., Pan, S., Qian, H., Zhu, X., 2021. Effects of quercetin on the efficacy of various chemotherapeutic drugs in cervical cancer cells. *Drug Des. Devel. Ther.* 15, 577-588.
- Zhang, Y., Wan, Y., Huo, B., Li, B., Jin, Y., Hu, X., 2018. Extracts and components of *Ficus carica* leaves suppress survival, cell cycle, and migration of triple-negative breast cancer MDA-MB-231 cells. *OncoTargets Ther.* 11, 4377-4386.
- Zheng, J., Guinter, M.A., Merchant, A.T., Wirth, M.D., Zhang, J., Stolzenberg-Solomon, R.Z., Steck, S.E., 2017. Dietary patterns and risk of pancreatic cancer: A systematic review. *Nutr. Rev.* 75, 883-908.